



## Scientific Comment

# Diarrhea after autologous stem cell transplantation in low-middle income countries: is *Clostridium difficile* the most prevalent infectious etiology?☆

Renato Cunha<sup>a,b,\*</sup>

<sup>a</sup> Hospital das Clínicas da Faculdade de Medicina de Ribeirão Preto da Universidade de São Paulo (HC FMRP USP), Ribeirão Preto, SP, Brazil

<sup>b</sup> Experimental Transplantation and Immunology Branch, National Cancer Institute/National Institutes of Health, Bethesda, USA

Diarrhea is a major cause of morbidity during high-dose chemotherapy followed by autologous stem cell transplantation (ASCT). It occurs in almost half of recipients of ASCT and generally, it is attributed to the effects of high-dose chemotherapy on the gastrointestinal mucosa and the effects of broad-spectrum antibiotic regimens for the prophylaxis and treatment of neutropenia. However, other etiologies might be considered, such as viral, bacterial and parasitic infections.<sup>1,2</sup>

Despite being the most common etiology of diarrhea after ASCT, the conditioning regime is seldom implicated in severe complications, with some exceptions. Cytarabine-containing regimens, high-dose melphalan ( $\geq 200$  mg/m<sup>2</sup>), and regimens containing multiple alkylating agents may cause more severe diarrhea.<sup>3</sup>

Conversely, infectious causes of diarrhea are less prevalent than damage from conditioning therapy. *Clostridium difficile* is considered the leading cause of infectious diarrhea among hospitalized patients and is a major concern in immunosuppressed individuals.<sup>4</sup> Viruses such as cytomegalovirus (CMV) and some serotypes of adenovirus might also be implicated in diarrhea after ASCT, but rarely cause life-threatening complications. Diarrhea secondary to intestinal parasite infection, such as *Cryptosporidium*, *Giardia lamblia*, *Entamoeba histolytica* and helminths, is rare among patients who come to

transplant without diarrhea.<sup>5,6</sup> *Strongyloides* infection and non-tuberculous mycobacterial have also been described after ASCT in patients from endemic areas.<sup>2</sup>

In this issue of Hematology, Transfusion and Cell Therapy, Castro et al. investigated the role of enteric pathogens in 47 recipients of ASCT between May 2011 and May 2013.<sup>7</sup> Thirty-nine recipients (83%) met the criteria for the diagnosis of diarrhea. Among them, conditioning regime toxicity was the most prevalent cause (35%). Interesting, a high prevalence of infectious etiology was identified. Through an investigational platform based on parasitological stool examination, stool cultures and the identification of A and B *C. difficile* toxins, 30% of the recipients had an infectious cause as the etiology with *Coccidia* being the most common. Unexpectedly, *C. difficile* presented a low incidence in comparison with currently published data<sup>4</sup> but, in accordance with the frequency and microbiological causes of diarrhea in low-middle income countries.<sup>8</sup>

These results revealed some potential different etiologies of infectious diarrhea in Brazil. However, due to the low number of patients evaluated from a single center, conclusions should be taken with caution. Nevertheless, these data might justify a multicenter study with the aim to clarify etiologies of diarrhea among Brazilian patients. A careful analysis of the

DOI of original article: <https://doi.org/10.1016/j.htct.2017.10.002>.

☆ See paper by Marcelo Dias de Castro et al. on pages [132–135].

\* Correspondence to: ETIB, NCI, NIH, 10 Center Drive, Bldg 10-CRC, Rm 3-3224, Bethesda, MD 20892, USA.

E-mail address: [renatolg@gmail.com](mailto:renatolg@gmail.com)

<https://doi.org/10.1016/j.htct.2018.02.002>

2531-1379/© 2018 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

causative agents could lead to more accurate management, early intervention and prevention of severe complications.

---

### Conflicts of interest

The author declares no conflicts of interest.

---

### REFERENCES

1. van Kraaij MG, Dekker AW, Verdonck LF, van Loon AM, Vinje J, Koopmans MP, et al. Infectious gastro-enteritis: an uncommon cause of diarrhoea in adult allogeneic and autologous stem cell transplant recipients. *Bone Marrow Transplant.* 2000;26(3):299-303.
2. Tuncer HH, Rana N, Milani C, Darko A, Al-Homsi SA. Gastrointestinal and hepatic complications of hematopoietic stem cell transplantation. *World J Gastroenterol.* 2012;18(16):1851-60.
3. Cox GJ, Matsui SM, Lo RS, Hinds M, Bowden RA, Hackman RC, et al. Etiology and outcome of diarrhea after marrow transplantation: a prospective study. *Gastroenterology.* 1994;107(5):1398-407.
4. Alonso CD, Dufresne SF, Hanna DB, Labbe AC, Treadway SB, Neofytos D, et al. *Clostridium difficile* infection after adult autologous stem cell transplantation: a multicenter study of epidemiology and risk factors. *Biol Blood Marrow Transplant.* 2013;19(10):1502-8.
5. Stein A, Voigt W, Jordan K. Chemotherapy-induced diarrhea: pathophysiology, frequency and guideline-based management. *Ther Adv Med Oncol.* 2010;2(1):51-63.
6. Bhat V, Vira H, Khattry N, Toshniwal M. *Cryptococcus laurentii* diarrhea post hematopoietic stem cell transplant. *Transpl Infect Dis.* 2017;19(2).
7. de Castro MD, Chebli JM, Costa LJ, Alves KR, Atalla A, Hallack Neto AE. Infectious diarrhea in autologous stem cell transplantation: high prevalence of coccidia in a South American center. *Hematol Transfus Cell Ther.* 2018;40(2): 132-5.
8. Pala C, Kaynar L, Buyukoglan R, Kurnaz F, Metan G, Yazar S, et al. Diarrhea in peripheral stem cell transplant recipients: a developing country's experience. *J Infect Dev Ctries.* 2014;8(5):635-41.