



Resistant and refractory hypertension: two sides of the same disease?

Hipertensão resistente e refratária: duas faces de uma mesma doença?

Authors

Elizabeth Silaid Muxfeldt^{1,2} 

Bernardo Chedier^{1,2}

Cibele Isaac Saad Rodrigues³ 

¹ Universidade Federal do Rio de Janeiro, Faculdade de Medicina, Pós-Graduação em Clínica Médica, Rio de Janeiro, RJ, Brasil.

² Universidade Estácio de Sá, Curso de Medicina, Rio de Janeiro, RJ, Brasil.

³ Pontifícia Universidade Católica de São Paulo, Faculdade de Ciências Médicas e da Saúde, Departamento de Medicina, Sorocaba, SP, Brasil.

ABSTRACT

Refractory hypertension (RfH) is an extreme phenotype of resistant hypertension (RH), being considered an uncontrolled blood pressure besides the use of 5 or more antihypertensive medications, including a long-acting thiazide diuretic and a mineralocorticoid antagonist. RH is common, with 10-20% of the general hypertensives, and its associated with renin angiotensin aldosterone system hyperactivity and excess fluid retention. RfH comprises 5-8% of the RH and seems to be influenced by increased sympathetic activity. RH patients are older and more obese than general hypertensives. It is strongly associated with diabetes, obstructive sleep apnea, and hyperaldosteronism status. RfH is more frequent in women, younger patients and Afro-americans compared to RFs. Both are associated with increased albuminuria, left ventricular hypertrophy, chronic kidney diseases, stroke, and cardiovascular diseases. The magnitude of the white-coat effect seems to be higher among RH patients. Intensification of diuretic therapy is indicated in RH, while in RfH, therapy failure imposes new treatment alternatives such as the use of sympatholytic therapies. In conclusion, both RH and RfH constitute challenges in clinical practice and should be addressed as distinct clinical entities by trained professionals who are capable to identify comorbidities and provide specific, diversified, and individualized treatment.

Keywords: Resistant Hypertension; Refractory Hypertension; Sympathetic Nervous System; Hyperaldosteronism.

RESUMO

A Hipertensão Arterial Refratária (HARf) representa um fenótipo extremo da hipertensão arterial resistente (HAR), sendo considerada a falência ao tratamento apesar do uso de 5 ou mais classes de anti-hipertensivos, incluindo um diurético tiazídico de longa ação e um antagonista mineralocorticoide. A HAR é comum (10-20%) entre os hipertensos em geral, sendo decorrente de hiperatividade do Sistema Renina Angiotensina Aldosterona e retenção hidrossalina. Aqueles com HARf correspondem a 5-8% dos resistentes e parecem sofrer maior influência catecolaminérgica. Os resistentes tendem a ter maior idade, ao sobrepeso e à obesidade. Comorbidades incluem diabetes, apneia obstrutiva do sono e status de hiperaldosteronismo. Refratários são afro-americanos em maior proporção, mais jovens e, predominantemente, mulheres. Ambos são fortemente associados à elevada albuminúria, HVE, doenças cardio e cerebrovasculares, além da doença renal crônica. O fenômeno do jaleco branco parece ser mais evidente nos resistentes. Quanto ao tratamento, a intensificação da terapia diurética está indicada nos resistentes, enquanto na HARf, a falência à terapia impôs novas alternativas de tratamento (“simpaticolíticas”). Em conclusão, tanto a HAR quanto a HARf constituem-se desafios na prática clínica e devem ser abordadas como entidades clínicas distintas por profissionais especialistas que identifiquem comorbidades e venham a prover um tratamento específico, diversificado e individualizado.

Palavras-chave: Hipertensão Resistente; Hipertensão Refratária; Sistema Nervoso Simpático; Hiperaldosteronismo.

Submitted on: 05/20/2018.

Approved on: 09/05/2018.

Correspondence to:

Cibele Isaac Saad Rodrigues.
E-mail: cbele.sr@gmail.com

DOI: 10.1590/2175-8239-JBN-2018-0108

INTRODUCTION

Resistant hypertension (RH) has been studied in several different populations

since the end of the 20th century. Nevertheless, it was only in 2008 that the American Heart Association published



the first guidelines on RH, standardizing its definition and establishing the main risk factors, secondary causes, and the diagnostic and therapeutic approach to these patients¹. Thenceforth, many studies have demonstrated the high cardiovascular morbidity and mortality and begun to advocate for new therapeutic regimens (i.e. adding definitively spironolactone as the fourth-line drug choice), as well as new interventional therapies seeking for a better blood pressure (BP) control.

In an effort to define a subgroup of high-risk patients who should benefit the most from these new therapies, the refractory hypertension (RfH) definition was established in 2012² for individuals with worst BP control and, possibly, the worst cardiovascular outcomes.

Despite the final common pathway of an increased sympathetic tonus and hydrosaline retention, the current literature suggests the existence of different clinical phenotypes with different prognoses. These phenotypes would range from arterial hypertension that is responsive to initial treatment to RH and, more recently, to RfH.²

Although RfH seems to be an extreme phenotype of RH, recent studies have suggested different pathophysiological mechanisms. Whereas an increased sympathetic activity plays a more important role in the former, inappropriate hydrosaline retention due to a renin-angiotensin-aldosterone system (RAAS) hyperactivity is a major factor in RfH.³⁻⁶ Therefore, although hypertension is frequently understood as part of a *continuum*, a better comprehension about the prevalence of RfH in different populations, as well as its clinical and prognostic differences from RH is essential, especially in the post-mineralocorticoid-antagonist-receptor era⁷⁻⁹.

DISCUSSION

DEFINITION

RH is defined as an office BP that remains above the goal despite the use of 3 or more anti-hypertensive agents of different mechanisms of action at optimal doses, preferentially including a diuretic agent. Patients with controlled office BP on 4 or more drugs are also considered RH.^{1,10}

In parallel to this, the definition of RfH has been evolving since 2012,² being currently regarded as the failure of office BP control despite the use of 5 or more anti-hypertensive agents including a long-acting thiazide-like diuretic (ideally chlorthalidone) or a loop diuretics, according to estimated glomerular filtration

rate (eGFR), besides a mineralocorticoid receptor antagonist (e.g. spironolactone) as the fourth drug.⁷

EPIDEMIOLOGY

The prevalence of RH, as estimated by multiple multicenter cohorts, lies between 10-20% of all treated individuals.¹¹⁻¹⁴ The increased prevalence, despite the improvement of anti-hypertensive regimens in the last 30 years, is explained by the progressive ageing of the population and by the obesity pandemic.¹⁵ Analyses excluding pseudo-resistant hypertension are needed to estimate the true prevalence of RH.^{8,16} In Brazil, the Brazilian Longitudinal Study of Adult's Health (ELSA) found an 11% RH prevalence among a cohort of more than 15,000 individuals between 35 and 74 years old.¹⁷

The prevalence of RfH has been estimated by a limited number of studies. Of particular importance, is the prospective analysis conducted by Dudenbostel,¹⁸ which reported a 5% prevalence of RfH among the RH referred to a specialized hypertension clinic. Additionally, in the REGARDs study,¹⁹ similar rates (3.6%) have been described among patients with controlled or uncontrolled RH, highlighting the low prevalence of RfH (0.5%) among the entire population of hypertensive patients. Recently, the analysis of a Spanish ABPM Registry evidenced a prevalence of 8% of RfH among the RH patients (16.9%).²⁰

MECHANISMS

RH is mainly attributed to RAAS hyperactivity and consequently to excessive hydrosaline retention, as evidenced by BP reduction with diuretic therapy that is proportional to effective intravascular volume depletion.²¹ This mechanism appears to be multifactorial, being associated with increasing age, obesity, chronic kidney disease (CKD) and diabetes, Afro-American ethnicity, excessive sodium intake, and, remarkably, to the magnitude of the hyperaldosteronism status.²²

In contrast, RfH would be less volume-dependent, since, by definition, its treatment with the association of diuretic drugs fails to achieve the BP goals.^{4,5} Thus, refractory hypertensives seem to be under a greater sympathetic influence, having lower levels of plasmatic aldosterone and a reduced 24-hour sodium excretion. Recent studies comparing patients with refractory to resistant hypertension have shown increased markers of

sympathetic activity in the former group: higher heart rate, increased 24-hour norepinephrine excretion, and a higher peripheral resistance.^{4,18,23}

CLINICAL CHARACTERISTICS AND COMORBIDITIES

Resistant hypertensives tend to be older, overweight, or obese. Commonly associated comorbidities include CKD, diabetes, obstructive sleep apnea, left ventricle hypertrophy (LVH), cardiovascular and cerebrovascular diseases and, lastly, hyperaldosteronism status.^{1,10,11,24}

Refractory patients, compared to their controlled resistant counterparts, are more likely to be younger, Afro-american, and, predominately, females.²¹ Regarding associated comorbidities, the most common are heart failure¹⁸, stroke,² CKD with moderately increased albuminuria, diabetes, metabolic syndrome, cardiovascular diseases,¹⁹ and left ventricular hypertrophy.²

CLINICAL APPROACH

When assessing a patient with possible RH, we must consider many important factors to define the diagnostic approach (Table 1).

The first step is to exclude common reasons for pseudo-resistance: inaccurate measurement of BP (special attention should be paid to the adequate size of the cuff for obese patients), poor adherence to both pharmacological and nonpharmacological therapy (i.e. low-sodium diet, physical activity, and weight loss), and an inadequate therapeutic regimen, especially in relation to the use and dosage of the diuretic agents prescribed.^{1,27,28} Once the pseudo-resistance is excluded, the following steps are recommended:

TABLE 1 DIAGNOSTIC APPROACH IN RESISTANT HYPERTENSION²⁶

Diagnostic Approach

- 1) Check therapeutic adhesion
- 2) Rule out pseudo-resistance
- 3) Adjust anti-hypertensive scheme
- 4) Perform initial complementary exams (Table 2)
- 4) Investigate secondary hypertension:
 - Obstructive sleep apnea;
 - Primary aldosteronism;
 - Renovascular hypertension;
 - Renal parenchymal disease.
- 5) Control blood pressure - ABPM

ABPM, Ambulatory Blood Pressure Monitoring.

A) AMBULATORY BLOOD PRESSURE MONITORING (ABPM)

Even though the definitions of both RH and RfH rely on the office BP measurement higher than 140/90 mmHg, the ABPM is a crucial tool in the diagnosis and follow-up of these patients due to the high prevalence (37% in different series) and magnitude of the white-coat effect observed in these patients.^{11,28} (Table 2) Moreover, the ABPM allows patients to be classified into 4 distinct groups (Figure 1) that will determine the subsequent diagnostic evaluation and management: true RH (office BP \geq 140/90 mmHg and either daytime BP \geq 135/85 mmHg or night time BP \geq 120/70 mmHg), white-coat RH (office BP \geq 140/90 mmHg and either daytime BP $<$ 135/85 mmHg and night time BP $<$ 120/70 mmHg), masked RH (office BP $<$ 140/90 mmHg and either daytime BP \geq 135/85 mmHg or night time BP \geq 120/70 mmHg), and controlled RH (office BP $<$ 140/90 mmHg and either daytime BP $<$ 135/85 mmHg and night time BP $<$ 120/70 mmHg).^{11,28,29}

On the other hand, among RfH patients, the white-coat phenomenon has not yet been adequately studied. In an analysis of the Spanish ABPM Registry²⁰, the prevalence of the white-coat effect was lower among refractory when compared to resistant hypertensives (26.7% versus 37.1%, $p < 0.001$). In a recent small prospective study with patients with RfH, a prevalence of only 6.5% was found,³⁰ suggesting that this phenomenon is much less common among these patients.

In addition, ABPM is essential in the follow-up of these patients at high cardiovascular risk, since it is the only available tool to assess nocturnal blood pressure. In clinical practice, this information allows adjustments to therapeutic regimens based on chronotherapy.^{31,32} It is recommended that patients with RH use at least one of their anti-hypertensive drugs at bedtime.^{31,32} It has been demonstrated that chronotherapy was capable of reversing the non-dipper pattern in these patients.³³

It is known that the non-dipper status is the most common pattern among patients with resistant hypertension, affecting up to 65% of these patients.¹¹ Furthermore, it is considered an important prognostic marker, especially for coronary artery disease.³⁴ In addition, ambulatory blood pressure during the three periods, but especially at nighttime, are strong predictors of stroke.³⁵


The Spanish ABPM registry compared resistant with refractory hypertensives and identified higher ambulatory BP levels in the latter group, with a

TABLE 2 INITIAL COMPLEMENTARY EXAMS

Complementary exams	Indication
ABPM	White coat-effect and nocturnal BP pattern
Fasting plasma glucose/HbA1c	Screening of abnormal glucose tolerance or <i>diabetes mellitus</i>
Serum cholesterol, LDL - cholesterol, HDL - cholesterol	Screening of dyslipidemia
Serum uric acid	Monitoring of uric acid by diuretic use. Possible prognostic marker
Serum potassium	Monitoring potassium especially before the onset of spironolactone. Screening of primary aldosteronism
Renal evaluation:	
Serum creatinine	Calculation of estimated GFR (MDRD ou CKD-EPI) Available in: http://ckdepi.org/equations/gfr-calculator/
Urine analysis	Verification of urinary sediment
Urinary protein, creatinine and albuminuria	Calculation of protein/creatinine or albumin/creatinine ratio - asymptomatic target organ or established kidney diseases evaluation
Renal ultrasound	Verification of anatomical changes
12-lead ECG	Screening of left ventricular hypertrophy (voltage criteria and strain pattern)

Notes: ABPM, Ambulatory Blood Pressure Monitoring; HbA1c, Glycated haemoglobin; ECG, electrocardiogram; GFR, glomerular filtration rate.

Figure 1. Classification of resistant hypertension into four subgroups according to office and ambulatory blood pressure measurements: controlled, masked, white-coat, and true resistant hypertension (authorized reproduction from Muxfeldt et al²⁸)



smaller nocturnal BP reduction. The prevalence of the non-dipper and of the riser patterns was 42.7% and 19.3% among RH patients and of 45.2% and 26.0% among refractory hypertensives, respectively.²⁰

B) LABORATORY EXAMS

At first evaluation, it is necessary to assess the metabolic profile and the renal function (serum creatinine, calculation of the eGFR, and albuminuria dosage) (Table 2).

Patients with RfH have a higher prevalence of diabetes *mellitus* (48.1% *versus* 33.5%, $p < 0.001$) and dyslipidemia (61.9% *versus* 51.7%, $p < 0.001$) than patients with RH.²⁰ The association between CKD and RH is well established, as both a cause and a consequence of therapeutic failure. Besides, moderately increased albuminuria and a reduction in the GFR³⁶ identify patients with a high cardiovascular risk^{37,38} and albuminuria reduction may be used as a therapeutic goal in these patients.³⁹

A higher prevalence of a eGFR < 60 mL/min/1.73 m² (32.1% *versus* 23.6%, $p < 0.001$) and of moderately increased albuminuria (38.3% *versus* 24.5%, $p < 0.001$) was identified in RfH patients when compared to resistant hypertensives in an analysis of the Spanish ABPM registry.²⁰

c) ELECTROCARDIOGRAM (ECG)

The 12-lead ECG is a useful tool of low-cost and widely available even in primary health units (Table 2). Left ventricular hypertrophy identified on ECG is an important prognostic marker indicating that a subclinical lesion is under development, even in patients who seem to have a well-controlled office BP. These patients may be experiencing masked RH or isolated nocturnal RH.^{8,9} The diagnosis of LVH will guide the choice of the therapeutic regimen. Preferentially, an inhibitor of the RAAS should be chosen, aiming for the regression of the LVH.⁴⁰ The electrocardiographic diagnosis of LVH implies an increase in the cardiovascular risk and its prevention or regression aims to improve the prognosis.⁴⁰

On the Spanish ABPM registry, electrocardiographically-diagnosed LVH was more prevalent among patients with RfH than in those with RH (27.6% versus 14.9%, $p < 0.001$).²⁰

d) CAUSES OF SECONDARY HYPERTENSION

By definition, in all individuals with suspected RH or RfH, secondary hypertension should be excluded. The most prevalent causes are obstructive sleep apnea, primary aldosteronism, renovascular

hypertension, renal parenchymal disease, and pheochromocytoma (Table 3).⁴¹ Other causes as coarctation of the aorta, Cushing's syndrome, hypo or hyperthyroidism, and acromegaly should be investigated only in situations where there are stigmas of the disease.

THERAPEUTIC APPROACH

NONPHARMACOLOGICAL STRATEGIES

Obesity, as well as physical inactivity, high sodium intake, smoking, and alcoholism are strongly associated with anti-hypertensive treatment failure, all of them considered important risk factors for RH.^{1,11,34,42} In this way, it is imperative to reinforce the importance of lifestyle changes.⁸

- Reduction of dietary sodium intake: (below 2 g/day of sodium, corresponding to 5 g/day of salt);
- DASH diet: use of the Dietary Approaches to Stop Hypertension;⁴³
- Weight loss: preferentially a BMI < 25 kg/m²;
- Physical activity: practicing aerobic exercises, dynamic resistance training, and isometric resistance training weekly (at least 30 minutes on 5-7 days per week), after cardiology evaluation;

TABLE 3 SCREENING FOR SECONDARY CAUSES OF HYPERTENSION

Clinical findings	Suspected diagnosis	Additional investigation
Snore, diurnal somnolence, metabolic syndrome	Obstructive sleep apnea	STOP-BANG questionnaire, Epworth Somnolence Scale. Gold standard: Polysomnography (AIH > 5/hour; moderate apnea: AIH > 15/hour; severe apnea: > 30/hour)
Resistant hypertension with or without hypokalemia. Adrenal nodule	Primary aldosteronism or adrenal hyperplasia	Serum aldosterone > 15 ng/dL Aldosterone/renin ratio > 30 Confirmatory tests: fludrocortisone suppression or saline infusion. Helicoidal CT or MRI
Oedema, anorexia, fatigue, anemia, increased serum urea and creatinine, urinary sediment or anatomic changes	Renal parenchymal disease	Urinalysis, calculation of eGFR, renal ultrasound, urinary albumin/creatinine and protein/creatinine ratio
Abdominal bruit, flash pulmonary oedema, rapid deterioration in renal function after inhibitor of RAAS use.	Renovascular diseases	Renal Duplex Doppler Ultrasonography and/or Magnetic resonance angiography, spiral computed tomography, intra-arterial digital subtraction angiography.
Episodic or persistent high BP with headache, heavy sweating, and palpitations	Pheochromocytoma	Plasma and 24-hour catecholamines and/or metanephrines CT and MRI

AIH, apnea-hypopnea index; CT, computed tomography; eGFR, estimated glomerular filtration rate; MRI, magnetic resonance imaging; RAAS, renin-angiotensin-aldosterone system.

Adapted from Malachias MVB et al. 7^o Diretriz Brasileira de Hipertensão Arterial.⁹

- Smokers: quitting smoking, preferentially with assistance;
- Alcohol: reduce the consumption;
- Avoidance of drugs that increase blood pressure.

PHARMACOLOGICAL STRATEGIES

The initial cornerstone of resistant hypertension treatment is the association of at least three classes of different drugs: i) an appropriate diuretic, preferentially a long-action thiazide diuretic (ex. chlorthalidone) in patients with normal renal function, or loop diuretics should replace thiazides if eGFR is $< 30 \text{ mL/min/1.73m}^2$ or in other edematous state; ii) a RAAS inhibitor (angiotensin-converting enzyme inhibitors and angiotensin receptor blockers; iii) long-acting dihydropyridines calcium-channel blockers.^{1,8,9,27} Even though hydrochlorothiazide is the most widely prescribed diuretic, chlorthalidone is the diuretic of choice because of its long-acting effect with higher efficacy.^{8,9} For patients with CKD stage 4 or 5 (eGFR lower than $30 \text{ mL/min/1.73m}^2$), loop diuretics must be prescribed and administered at least twice a day.^{8,9}

Coronary artery diseases, heart failure, and arrhythmias are special situations when beta-blockers can substitute calcium antagonists at the initial therapeutic scheme with 3 drugs.^{8,9,27}

RH treatment should be based on diuretic therapy intensification, with special emphasis in the use of spironolactone as a fourth drug, because its association with thiazides provides additive effect in reducing BP.^{6,7,18,44} The ASPIRANT Trial⁴⁵⁻⁴⁷ showed that the addition of spironolactone (25 mg/day) *versus* placebo lowers systolic BP significantly, especially in older patients. Even in resistant hypertensives with CKD, the spironolactone may be used, except in cases of hyperkalemia.^{48,49}

Recently, the ReHOT study - a Brazilian multicenter study comparing spironolactone *versus* clonidine as a fourth-drug therapy in RH - found that both drugs achieved office and ambulatory BP control in similar rates, but spironolactone promoted greater decrease in systolic and diastolic 24-hour BP and diastolic daytime BP, without nighttime BP difference. Nevertheless, spironolactone was considered preferable as the fourth-drug therapy because of its easier posology, less adverse effects, and consequently better long-term adherence.⁵⁰

If after the four-drug scheme ambulatory BP remains uncontrolled, a fifth-line drug should be added. Possible fifth or sixth drugs are beta-blockers

(preferentially the ones with vasodilation effect, as carvedilol, bisoprolol⁵¹ or nebivolol), central alfa1-agonists (clonidine or doxazosin⁵¹), and direct vasodilators (hydralazine or minoxidil). The latter two are capable to lowering BP although they do not reduce cardiovascular morbidity and mortality.^{8,9}

Regarding RfH with failure in controlling BP despite the use of optimized therapeutic scheme with 5 or even 6 drugs, new interventions have emerged, as sympatholytic therapies.^{18,24} Among these new strategies, we highlight the following:

BAROREFLEX ACTIVATION THERAPY

The Rheos system is a programmable device that consists of a battery-powered implantable generator that works by electrically activating the carotid baroreflex. The Rheos Pivotal Trial did not identify long-term benefits.⁵²

RENAL SYMPATHETIC DENERVATION

The renal denervation procedure uses radiofrequency energy to ablate the nerves within the main renal arteries. This therapy was evaluated by three studies called SYMPPLICITY.⁵³ Different meta-analyses, including a Cochrane's revision, showed that the procedure was safe, but did not significantly decrease BP.⁵⁴⁻⁵⁶ The authors advised to await further trials with next-generation catheters, longer follow-up and bigger sample sizes, and especially with standardized procedures.⁵⁴

CONTINUOUS POSITIVE AIRWAY PRESSURE (CPAP)

Although the benefits in BP control with CPAP use in resistant hypertensives with moderate-severe sleep apnea are not well established with controversy results in different populations,⁵⁷ the CPAP should be indicated as an adjuvant treatment, in so far as the adherence is greater than 4 hours per night, improving the quality of life and probably reestablishing the dipper pattern.⁵⁸

CENTRAL ILIAC ARTERIOVENOUS ANASTOMOSIS

The ROX Medical arteriovenous coupler is a stent-like device that exhibits shape memory to self-expand, forming an AV anastomosis in central iliac. The ROX control HTN demonstrated significant BP decrease, possibly reducing cardiovascular morbidity in those patients.⁵⁹ Notwithstanding, this is an isolated study and more clinical evidence is necessary.

Table 4 summarizes the main differences between resistant and refractory hypertension observed in various populations.

TABLE 4 CHARACTERISTICS OF RESISTANT AND REFRACTORY HYPERTENSION

Characteristics	Resistant hypertension	Refractory hypertension
Prevalence	10-20%	5%
Mechanism	Volume-dependent	Increased sympathetic activity
Gender	Women	Women
Age	Older	Younger
Obesity	↑	↑↑
Diabetes	↑	↑↑
Dyslipidemia	↑	↑↑
Left ventricular hypertrophy	↑↑	↑↑↑
Moderately increased albuminuria	↑	↑↑
eGFR < 60 ml/min/1,73m ²	↑	↑↑
Coronary heart disease	↑	↑
Previous cardiovascular disease	↑↑	↑↑↑
Obstructive sleep apnea	↑	Undetermined
Aldosterone	↑	↔
Sodium	↑	↔
Cardiovascular risk	↑↑	Apparently increased

eGFR, estimated glomerular filtration rate.

CONCLUSION

Despite the common final pathway of hypertension encompassing hydrosaline retention and increased sympathetic tonus, the existence of many phenotypes with distinct clinical paths and prognosis, a broad spectrum ranging from easily controlled hypertension to RH, and more recently, RfH has been suggested.

Even though these two entities are frequently considered a continuum of the same process, it is interesting to observe that they seem to have different pathophysiological mechanisms, suggesting two distinct conditions.

RH patients compared with general hypertensives, are older and more obese. The principal associated comorbidities are established CKD, diabetes, sleep apnea, stroke, and cardiovascular diseases, all of them involving the hyperaldosteronism status. ABPM is mandatory in the diagnosis and follow-up of those patients because of a high magnitude of the white-coat effect.

Moreover, refractory hypertensives compared with controlled RH are younger, predominantly women, and Afro-american. They also have a high prevalence of heart failure, stroke, and CKD with moderately increased albuminuria and LVH. The white-coat effect seems to be less evident in those patients.

In RH, the therapeutic scheme should be based on the intensification of diuretic therapy, emphasizing the spironolactone as the fourth drug associated with a

long-acting thiazide, as chlorthalidone. On the other hand, as RfH usually fails all used classes of anti-hypertensives including association of different diuretics, RfH treatment is not well established and new therapies have been proposed such as sympatholytic intervention.

The unfavorable cardiovascular and renal prognosis of RH patients is well established, but future longitudinal studies are necessary to define the morbidity and mortality of RfH.

Resistant and refractory hypertension are challenges in clinical practice and should be addressed as different entities, ideally by specialized professionals capable of identifying comorbidities and to provide diversified and individualized treatment.

REFERENCES

- Carey RM, Calhoun DA, Bakris GLO, Brook RD, Daugherty SL, Dennison-Himmerlfarb CR et al. Resistant hypertension: detection, evaluation and management. A scientific statement from the American Heart Association. *Hypertension* 2018;72:e53-e90
- Acelajado MC, Pisoni R, Dudenbostel T, Dell'Italia LJ, Cartmill F, Zhang B, et al. Refractory hypertension: definition, prevalence, and patient characteristics. *J Clin Hypertens* 2012;14:7-12.
- Leibowitz A, Schiffrin EL. Immune mechanisms in hypertension. *Curr Hypertens Rep* 2011;13:465-72.
- Modolo R, de Faria AP, Almeida A, Moreno H. Resistant or refractory hypertension: are they different? *Curr Hypertens Rep* 2014;16:485.
- Dudenbostel T, Acelajado MC, Pisoni R, Li P, Oparil S, Calhoun D. Refractory Hypertension: Evidence of Heightened Sympathetic Activity as a Cause of Antihypertensive Treatment Failure. *Hypertension* 2015;66:126-33.

6. Yugar-Toledo JC, Modolo R, de Faria AP, Moreno H. Managing resistant hypertension: focus on mineralocorticoid-receptor antagonists. *Vasc Health Risk Manag* 2017;13:403-11.
7. de Souza F, Muxfeldt ES, Fiszman R, Salles GF. Efficacy of spironolactone therapy in patients with true resistant hypertension. *Hypertension* 2010;55:147-52.
8. Williams B, Mancia G, Spiering W, Rosei EA, Azizi M, Burnier M et al. 2018 European Society for Cardiology (ESC)/European Society of Hypertension (ESH) guidelines for the management of arterial hypertension. *J Hypertens* 2018;36(10):1953-2041.
9. Malachias MVB, Souza WKS, Plavnik FL, Rodrigues CIS, Brandão AA, Neves MFT, et al. 7ª Diretriz Brasileira de Hipertensão Arterial. *Arq Bras Cardiol* 2016;107:1-83.
10. Braam B, Taler SJ, Rahman M, Fillaus JA, Greco BA, Forman JP, et al. Recognition and Management of Resistant Hypertension. *Clin J Am Soc Nephrol* 2017;12:524-35.
11. de la Sierra A, Segura J, Banegas JR, Gorostidi M, de la Cruz JJ, Armario P, et al. Clinical features of 8295 patients with resistant hypertension classified on the basis of ambulatory blood pressure monitoring. *Hypertension* 2011;57:898-902.
12. Persell SD. Prevalence of resistant hypertension in the United States, 2003-2008. *Hypertension* 2011;57:1076-80.
13. Daugherty SL, Powers JD, Magid DJ, Tavel HM, Masoudi FA, Margolis KL, et al. Incidence and prognosis of resistant hypertension in hypertensive patients. *Circulation* 2012;125:1635-42.
14. Sim JJ, Bhandari SK, Shi J, Liu IL, Calhoun DA, McGlynn EA, et al. Characteristics of resistant hypertension in a large, ethnically diverse hypertension population of an integrated health system. *Mayo Clin Proc* 2013;88:1099-107.
15. Egan BM, Zhao Y, Axon RN, Brzezinski WA, Ferdinand KC. Uncontrolled and apparent treatment resistant hypertension in the United States, 1988 to 2008. *Circulation* 2011;124:1046-58.
16. Sarafidis PA, Georgianos P, Bakris GL. Resistant hypertension—its identification and epidemiology. *Nat Rev Nephrol* 2012;9:51-8.
17. Lotufo PA, Pereira AC, Vasconcellos PS, Santos IS, Mill JG, Bensenor IM. Resistant hypertension: risk factors, subclinical atherosclerosis, and comorbidities among adults—the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *J Clin Hypertens (Greenwich)* 2015;17:74-80.
18. Dudenbostel T, Siddiqui M, Gharpure N, Calhoun DA. Refractory versus resistant hypertension: Novel distinctive phenotypes. *J Nat Sci* 2017;3:pii: e430.
19. Calhoun DA, Booth JN 3rd, Oparil S, Irvin MR, Shimbo D, Lackland DT, et al. Refractory hypertension: determination of prevalence, risk factors, and comorbidities in a large, population-based cohort. *Hypertension* 2014;63:451-8.
20. Armario P, Calhoun DA, Oliveras A, Blanch P, Vinyoles E, Banegas JR, et al. Prevalence and Clinical Characteristics of Refractory Hypertension. *J Am Heart Assoc* 2017;6:pii: e007365.
21. Siddiqui M, Calhoun DA. Refractory versus resistant hypertension. *Curr Opin Nephrol Hypertens* 2017;26:14-9.
22. Hwang AY, Dietrich E, Pepine CJ, Smith SM. Resistant Hypertension: Mechanisms and Treatment. *Curr Hypertens Rep* 2017;19:56.
23. Calhoun DA. Refractory and Resistant Hypertension: Antihypertensive Treatment Failure versus Treatment Resistance. *Korean Circ J* 2016;46:593-600.
24. Muxfeldt ES, Salles GF. Pulse pressure or dipping pattern: which one is a better cardiovascular risk marker in resistant hypertension? *J Hypertens* 2008;26:878-84.
25. Departamento de Hipertensão Arterial da Sociedade Brasileira de Cardiologia. I Posicionamento Brasileiro sobre hipertensão arterial resistente. *Arq Bras Cardiol* 2012;99:576-85.
26. de Souza F, Muxfeldt ES, Salles GF. Prognostic factors in resistant hypertension: implications for cardiovascular risk stratification and therapeutic management. *Expert Rev Cardiovasc Ther* 2012;10:735-45.
27. Muxfeldt ES, de Souza F, Salles GF. Resistant hypertension: a practical clinical approach. *J Hum Hypertens* 2013;27:657-62.
28. Muxfeldt ES, Salles GF. How to use ambulatory blood pressure monitoring in resistant hypertension. *Hypertens Res* 2013;36:385-9.
29. Wei FF, Zhang ZY, Huang QF, Staessen JA. Diagnosis and management of resistant hypertension: state of the art. *Nat Rev Nephrol* 2018;14:428-41.
30. Siddiqui M, Judd EK, Oparil S, Calhoun DA. White-Coat Effect Is Uncommon in Patients With Refractory Hypertension. *Hypertension* 2017;70:645-51.
31. Hermida RC, Ayala DE, Mojón A, Fernández JR. Effects of time of antihypertensive treatment on ambulatory blood pressure and clinical characteristics of subjects with resistant hypertension. *Am J Hypertens* 2010;23:432-9.
32. Hermida RC, Ayala DE, Fernández JR, Mojón A, Crespo JJ, Ríos MT, et al. Bedtime Blood Pressure Chronotherapy Significantly Improves Hypertension Management. *Heart Fail Clin* 2017;13:759-73.
33. Hermida RC, Ayala DE, Fernández JR, Calvo C. Chronotherapy improves blood pressure control and reverts the nondipper pattern in patients with resistant hypertension. *Hypertension* 2008;51:69-76.
34. Muxfeldt ES, Cardoso CR, Salles GF. Prognostic value of nocturnal blood pressure reduction in resistant hypertension. *Arch Intern Med* 2009;169:874-80.
35. Salles GF, Cardoso CR, Muxfeldt ES. Prognostic influence of office and ambulatory blood pressures in resistant hypertension. *Arch Intern Med* 2008;168:2340-6.
36. Salles GF, Cardoso CR, Pereira VS, Fiszman R, Muxfeldt ES. Prognostic significance of a reduced glomerular filtration rate and interaction with microalbuminuria in resistant hypertension: a cohort study. *J Hypertens* 2011;29:2014-23.
37. Oliveras A, Armario P, Sierra C, Arroyo JA, Hernández-del-Rey R, Vazquez S, et al. Urinary albumin excretion at follow-up predicts cardiovascular outcomes in subjects with resistant hypertension. *Am J Hypertens* 2013;26:1148-54.
38. da Costa PM, Cortez AF, de Souza F, Mares GS, dos Santos BDM, Muxfeldt ES. Prognostic impact of baseline urinary albumin excretion rate in patients with resistant hypertension: a prospective cohort study. *J Hum Hypertens* 2018;32:139-49.
39. Salles GF, Cardoso CR, Fiszman R, Muxfeldt ES. Prognostic importance of baseline and serial changes in microalbuminuria in patients with resistant hypertension. *Atherosclerosis* 2011;216:199-204.
40. Salles GF, Cardoso CR, Fiszman R, Muxfeldt ES. Prognostic impact of baseline and serial changes in electrocardiographic left ventricular hypertrophy in resistant hypertension. *Am Heart J* 2010;159:833-40.
41. Pedrosa RP, Drager LF, Gonzaga CC, Sousa MG, de Paula LK, Amaro AC, et al. Obstructive sleep apnea: the most common secondary cause of hypertension associated with resistant hypertension. *Hypertension* 2011;58:811-7.
42. Brambilla G, Bombelli M, Seravalle G, Cifkova R, Laurent S, Narkiewicz K, et al. Prevalence and clinical characteristics of patients with true resistant hypertension in central and Eastern Europe: data from the BP-CARE study. *J Hypertens* 2013;31:2018-24.
43. Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, et al.; DASH-Sodium Collaborative Research Group. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med* 2001;344:3-10.
44. Dudenbostel T, Calhoun DA. Use of Aldosterone Antagonists for Treatment of Uncontrolled Resistant Hypertension. *Am J Hypertens* 2017;30:103-9.
45. Václavík J, Sedlák R, Plachy M, Navrátil K, Plásek J, Jarkovsky J, et al. Addition of spironolactone in patients with resistant arterial hypertension (ASPIRANT): a randomized, double-blind, placebo-controlled trial. *Hypertension* 2011;57:1069-75.

46. Vaclavik J, Sedlak R, Jarkovsky J, Kocianova E, Taborsky M. Effect of spironolactone in patients with resistant arterial hypertension in relation to age and sex: insights from the aspirant trial. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2014;158:384-90.
47. Vaclavik J, Sedlak R, Jarkovsky J, Kocianova E, Taborsky M. The effect of spironolactone in patients with resistant arterial hypertension in relation to baseline blood pressure and secondary causes of hypertension. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2013;157:50-5.
48. Abolghasmi R, Taziki O. Efficacy of low dose spironolactone in chronic kidney disease with resistant hypertension. *Saudi J Kidney Dis Transpl* 2011;22:75-8.
49. Pisoni R, Acelajado MC, Cartmill FR, Dudenbostel T, Dell'Italia LJ, Cofield SS, et al. Long-term effects of aldosterone blockade in resistant hypertension associated with chronic kidney disease. *J Hum Hypertens* 2012;26:502-6.
50. Krieger EM, Drager LE, Giorgi DMA, Pereira AC, Barreto-Filho JAS, Nogueira AR, et al.; ReHOT Investigators. Spironolactone Versus Clonidine as a Fourth-Drug Therapy for Resistant Hypertension: The ReHOT Randomized Study (Resistant Hypertension Optimal Treatment). *Hypertension* 2018;71:681-90.
51. Williams B, MacDonald TM, Morant S, Webb DJ, Sever P, McInnes G et al. British Hypertension Society s PATHWAY Study Group. Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial. *Lancet* 2015;386:2059-2068.
52. Bakris GL, Nadim MK, Haller H, Lovett EG, Schafer JE, Bisognano JD. Baroreflex activation therapy provides durable benefit in patients with resistant hypertension: results of long-term follow-up in the Rheos Pivotal Trial. *J Am Soc Hypertens* 2012;6:152-8.
53. Bakris GL, Townsend RR, Liu M, Cohen SA, D'Agostino R, Flack JM, et al.; SYMPLICITY HTN-3 Investigators. Impact of renal denervation on 24-hour ambulatory blood pressure: results from SYMPLICITY HTN-3. *J Am Coll Cardiol* 2014;64:1071-8.
54. Elmula FF, Jin Y, Larstorp AC, Persu A, Kjeldsen SE, Staessen JA. Meta-analysis of five prospective and randomized trials of renal sympathetic denervation on office and ambulatory systolic blood pressure in treatment resistant hypertension. *J Hypertens* 2015;33:e107.
55. Fadl Elmula F, Jin Y, Yang WY, Thijs L, Lu YC, Larstorp AC, et al.; European Network Coordinating Research On Renal Denervation (ENCOREd) Consortium. Meta-analysis of randomized controlled trials on renal denervation in treatment-resistant hypertension. *Blood Press* 2015;24:263-74.
56. Coppolino G, Pisano A, Rivoli L, Bolignano D. Renal denervation for resistant hypertension. *Cochrane Database Syst Rev* 2017;2:CD011499.
57. Iftikhar IH, Valentine CW, Bittencourt LR, Cohen DL, Fedson AC, Gislason T, et al. Effects of continuous positive airway pressure on blood pressure in patients with resistant hypertension and obstructive sleep apnea: a meta-analysis. *J Hypertens* 2014;32:2341-50.
58. Martínez-García MA, Capote F, Campos-Rodríguez F, Lloberes P, Díaz de Atauri MJ, Somoza M, et al.; Spanish Sleep Network. Effect of CPAP on blood pressure in patients with obstructive sleep apnea and resistant hypertension: the HIPARCO randomized clinical trial. *JAMA* 2013;310:2407-15.
59. Lobo MD, Ott C, Sobotka PA, Saxena M, Stanton A, Cockcroft JR, et al. Central Iliac Arteriovenous Anastomosis for Uncontrolled Hypertension: One-Year Results From the ROX CONTROL HTN Trial. *Hypertension* 2017;70:1099-105.