

Volume 36 Issue 4 -Oct/Nov/Dec 2014

EVALUATION OF FOOD INTAKE AND EXCRETION OF METABOLITES IN NEPHROLITHIASIS

AValiação da ingestão alimentar e excreção de metabólitos na nefrolitíase

Évellyn Alves Gordiano¹, Larissa Marques Tondin^{2,3}, Renata Costa de Miranda¹, Deise Regina Baptista¹, Maurício Carvalho¹

¹ Federal University of Paraná (UFPR).

² Pontifical Catholic University of Paraná (PUCPR).

³ Hospital de Clínicas.

Maurício Carvalho was not cited as one of the authors in the original article published on the Brazilian Journal of Nephrology 36 (4), and Table 1 contained an error. The corrected version of the Table is printed below:

	NL		CONTROL		<i>p</i> -value
	N	Mean (SD)	n	Mean (SD)	
Age	31	52.23 (10.56)	18	52.78 (14.05)	0.877 ^a
BMI	31	29.44 (4.92)	18	31.44 (5.30)	0.189 ^a
Excreted oxalate	21	23.19 (12.52)	8	25.56 (15.05)	0.659 ^a
Excreted citrate	21	812.45 (566.67)	9	458.00 (387.83)	0.039 ^{b*}

^a Student's *t*-test; ^b Mann-Whitney. *Statistically significant difference ($p < 0.05$).