





Patients' Description of Rectal Effluents Help Predict the Quality of Colonoscopy Preparation

Manuel Antonio Lescano Lescano¹ Rodrigo Strehl Machado² Maria Rachel Rohr¹
Sender Jankiel Miszputen¹

¹Department of Gastroenterology, Universidade Federal de São Paulo, São Paulo, SP, Brazil

²Department of Pediatrics, Universidade Federal de São Paulo, São Paulo, SP, Brazil

Address for correspondence Manuel Antonio Lescano Lescano, M.D., Department of Gastroenterology, Universidade Federal de São Paulo, Rua Pedro de Toledo, 861/869, Vila Clementino, São Paulo, SP, Brazil, 04039-032 (e-mail: manuellescanol@hotmail.com).

J Coloproctol 2024;44(2):e106–e110.

Abstract

Introduction Evaluation of patients' reports of characteristics of rectal effluents as a predictor of the quality of the colonoscopy preparation assessed by the endoscopist.

Methods A total of 270 patients, aged 18 or older, were consecutively included to perform an outpatient colonoscopy, for a period of 8 months. Demographic and clinical data were collected and evaluated, as well as the rectal effluents' characteristics and data concerning the colonoscopy. The quality of bowel preparation was evaluated by employing the Boston Bowel Preparation Scale. The association between rectal effluents and the quality of preparation was verified by binary logistic regression.

Results Of the 270 patients, 67.3% were female, with a mean age of 59.69 ± 12.48 years. Reports of dark and thick, dark orange, or brown and thick effluents produced a higher likelihood of inadequate preparation (OR 4.26, CI 95% 1.51; 11.14, $p = 0.004$).

Conclusions Reports of dark and thick, dark orange, or brown and thick rectal effluents are predictors of inadequate preparation in the endoscopist assessment.

Keywords

- ▶ colonoscopy
- ▶ rectal effluents
- ▶ bowel preparation

Introduction

Colonoscopy represents the most important method of evaluation of the lower gastrointestinal tract, allowing a complete assessment of the colonic mucosa.¹ Studies have that about 25% of the patients exhibited inadequate colonoscopy preparation,^{2,3} which denotes lower detection rates of preneoplastic lesions, an increased rate of complications during the procedure, as well as higher direct and indirect costs compared to procedures performed under ideal preparation.^{4,5}

In a meta-analysis that evaluated the impact of bowel preparation quality on adenoma detection rate, it was established that inadequate preparation is significantly associated with a lower adenoma detection rate⁶ and, consecutively, with an increased likelihood of interval cancer and mortality from colorectal cancer.⁷

In practice, prior to the procedure, the patient is questioned about the aspect of their rectal effluents to predict the quality of preparation. Three papers, two from the United States and one from South Korea approach this subject, though none of them demonstrated great consensus between patients' descriptions and the endoscopist's assessment.^{8–10}

Our main goal was to evaluate the patients' reports of the rectal effluent characteristics as a predictor of the quality of colonoscopy preparation as assessed by the endoscopist.

Methods

A total of 270 patients, aged 18 or older, were consecutively included to perform an outpatient colonoscopy, for a period of 8 months. They were referred to the Endoscopy Service of the General Hospital of Vitória da Conquista, Bahia, for the

received

July 31, 2023

accepted after revision

April 9, 2024

DOI <https://doi.org/>

10.1055/s-0044-1787289.

ISSN 2237-9363.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil



Fig. 1 Patient's description of the color of rectal effluents and the correlation with the quality of bowel preparation.

procedure. The study protocol followed the principles of the Declaration of Helsinki and was approved by the Ethics Committee of the Federal University of São Paulo (CAAE 23020319.3.00000.5505).

The following were used as exclusion criteria: formal restriction for the procedure (suspected colon perforation, toxic megacolon, severe colitis); patients with a medical history of inflammatory bowel disease; inability to properly comprehend the preparation instructions; patients who have undergone partial or total colectomy; patients scheduled for advanced therapeutic procedures (e.g., pre-scheduled polypectomy or mucosectomy); incomplete colonoscopy due to endoscopy restriction. The assigned nurse explained in verbal and written form the instructions for bowel preparation, followed by the patient's signature on the Informed Consent Form.

On the day of the procedure, prior to the examination, another nurse from the research team oversaw the clinical and epidemiological data questionnaires and collected data about the last rectal effluent. The description of the last rectal effluent was guided by **Figure 1**, and the patient should indicate which characteristic best matched the graph.

The two assigned endoscopists in charge of performing the procedures were unaware of the descriptions of rectal effluents provided by the patients. All participants underwent preparation by oral administration of two sachets of a laxative (PicoPrep®) consisting of 10.0 mg of sodium picosulfate, 3.5 g of magnesium oxide, and 12.0 g of citric acid anhydrous, as per verbal and written instructions.

All colonoscopies were performed in the afternoon. The preparation regimen used was the "split-dose" regimen. The Boston Bowel Preparation Scale was employed for comparison with the descriptions provided by the patients. A score of ≥ 2 in each segment was evaluated as adequate, and a score of ≤ 1 was inadequate. Prior to the start of the study, the endoscopists watched a video on the evaluation of colonoscopy preparation

using the Boston Bowel Preparation Scale as a method of enhancing the results' consensus.¹¹ Each of them had great experience in conducting the procedure, with more than 3,000 colonoscopies performed by each one.

Statistical Analysis

The following descriptions by patients before the procedure were considered inadequate preparation: dark, thick, and with particles; brown, thick, and with particles; and semi-clear dark orange. Mostly clear light orange and clear light yellow were considered adequate. Similarly, in the endoscopist's assessment, a Boston score of ≥ 6 indicated adequate preparation, while a Boston score of < 6 indicated inadequate preparation.

Qualitative variables were represented as absolute and relative frequencies, while quantitative variables were represented by mean \pm standard deviation (median). The latter ones were subjected to the Shapiro-Wilk test.

The connection between sociodemographic, clinical, and rectal effluent aspects as well as the quality of preparation, was assessed by the binary logistic regression model, with results presented as an odds ratio (OR) and their respective confidence interval (CI) of 95%. Additionally, there was also an examination of the sensitivity and specificity of rectal effluent reports regarding the quality of preparation as assessed by the endoscopist. The analyses were conducted using the R 4.0.5 software, and a level of $p < 0.05$ was considered statistically significant.

Results

Patients' Characteristics and Demographic Findings

Table 1 illustrates the demographic and clinical characterization of the entire sample, which consisted of 270 subjects: 67.4% were female; with a mean age of 56.69 ± 12.48 years; and 40% had finished primary education. More than half

Table 1 Sample's demographic and clinical characterization

Characteristics	Total (n = 270)	P-value
Sex	182 (67.4%)	0.796 ^Q
Female	88 (32.5%)	
Male		
Age (mean ± SD (median))	56.69 ± 12.48 (57.00)	0.216 ^T
< 60 years	161 (59.6%)	
60 years and older	109 (40.4%)	
Educational level	36 (13.3%)	0.899 ^Q
Illiterate	108 (39.7%)	
Primary education	40 (14.8%)	
Lower secondary education	76 (28.1%)	
High school	10 (3.7%)	
Higher education		
Colonoscopy indication	142 (52.5%)	0.712 ^Q
Prevention	111 (41.1%)	
Diagnosis	17 (6.2%)	
Post-polypectomy follow-up		
Associated diseases	33 (12.2%)	0.582 ^Q
Diabetes	6 (2.2%)	0.684 ^F
Depression	56 (20.7%)	1.000 ^Q
Constipation		
Medication use	29 (10.7%)	0.562 ^Q
Hypoglycemic agents	10 (3.7%)	0.540 ^Q
Antidepressants		
Surgical history	51 (18.8%)	0.090 ^Q
Abdominal	95 (52.1%)	0.948 ^Q
Gynecological* (n = 182)		

^Q Chi-squared test, ^F Fisher's exact test, ^T Student's t-test for independent samples.

* Gynecological surgeries were evaluated only in women.

(52.6%) of the colonoscopies were performed as a means of prevention, and, among associated diseases, 21.0% had constipation, 12.2% had diabetes, and 2.2% had depression. Only 11.0% made use of hypoglycemic agents, and 4.0% took antidepressants. Prior abdominal surgeries accounted for 18.8%, and 52.1% of gynecological surgeries were reported by women.

Colonoscopy Results

Cecal intubation could be accomplished in all colonoscopies.

Adenoma detection rate in the sample was 27.2% (CI 95% 21.9%; 32.5%), with 25.1% (CI 95% 18.8%; 31.4%) for women, and 31.5% (CI 95% 21.8%; 41.2%) for men. Diverticula were detected in 24.6% of the procedures (CI 95% 19.5%; 29.7%), as shown in ►Table 2.

Consensus between Rectal Effluents Reports and the Quality of Bowel Preparation.

►Table 3 shows that the reports of effluents that were dark and thick, brown and thick, or dark orange indicated increased rates of inadequate preparation (OR 4.26, CI 95% 1.51; 11.14, $p = 0.004$), as well as when adjusted for sex, educational level, presence of diverticula and colonoscopy indication (OR 3.58, CI 95% 1.22; 9.74, $p = 0.015$). Reports of dark and thick, brown and thick, or dark orange

Table 2 Evaluation of colonoscopy outcomes

Variables	
Adenoma detection rate	74 (27.2%)
Female	46 (25.1%)
Male	28 (31.5%)
Diverticula	67 (24.6%)

Chi-squared test.

effluents displayed a sensitivity of 30.4% and specificity of 90.7% for assessing inadequate preparation.

Accuracy of Patient Assessment

►Table 4 indicates that patients with more than eight years of education had a higher likelihood of accurately assessing rectal effluents (OR 3.70, CI 95% 1.61; 10.06, $p = 0.005$), as well as when adjusted for age group and associated diseases (OR 3.84, CI 95% 1.64; 10.60, $p = 0.005$).

Discussion

There is limited literature on patients' perceptions of the quality of bowel preparation. In 2004, Harewood, Wright, and Baron⁸ evaluated the descriptions of rectal effluents by 474 patients who underwent outpatient colonoscopy. The authors concluded that the patients either overestimated or underestimated the quality of rectal effluents when compared to the endoscopist's assessment.

In a subsequent study published by Fatima, Johnson, and Rex⁹ in 2010, 429 subjects underwent outpatient colonoscopy in three different hospitals. A slight consensus was verified between the patients' description of rectal effluents and the endoscopist's assessment. However, there was a higher likelihood of inadequate preparation when the effluent description was solid or liquid brown. It is also important to note that both studies did not employ a validated colonoscopy preparation assessment scale.

In 2015, So et al.¹⁰ assessed the description of 138 individuals about their last three rectal effluents, with photographic examples provided as scoring and employment of the Aronchick scale. The images with clear rectal effluents were assigned a score of 1, up to 5 for semi-solid stool effluents. A score of 3 in the sum of the last three rectal effluents represented the most adequate reported preparation. The authors found statistical significance, although with poor clinical relevance. The study had several limitations, including being conducted by a single endoscopist, the inappropriate use of the Aronchick scale as it is impractical and unreliable in practice, and relying on patients' accurate recall of their last three effluents.

In the present study, there was a four times higher chance of inadequate preparation when the description of rectal effluents was that it was dark and thick, brown, and thick, or dark orange (OR 4.26, CI 95% 1.51; 11.14, $p = 0.004$). This finding is similar to the one reported by Fatima, Johnson, and Rex,⁹ and as practical applicability, may optimize preparation before patient sedation and the start of the procedure. As suggested

Table 3 Predictor of quality of inadequate preparation

Variables	Gross		Adjusted	
	OR (CI 95%)	P-value	OR (CI 95%)	P-value
Sex				
Female	1.00	–	1.00	–
Male	2.44 (1.02; 5.85)	0.043	2.31 (0.94; 5.75)	0.068
Educational level				
Less than 8 years	1.00	–	1.00	–
8 years or higher	0.29 (0.07; 1.09)	0.054	0.37 (0.08; 1.17)	0.128
Diverticula				
No	1.00	–	1.00	–
Yes	0.62 (0.18; 1.73)	0.403	0.81 (0.22; 2.39)	0.722
Report of rectal effluents				
Dark and thick, brown and thick, or dark orange	4.26 (1.51; 11.14)	0.004	3.58 (1.22; 9.74)	0.015
Light yellow or light orange	1.00	–	1.00	–

Table 4 Factors associated with patient's assessment accuracy (whether the patient's assessment matched the endoscopist's)

Variables	Gross		Adjusted	
	OR (CI 95%)	P-value	OR (CI 95%)	P-value
Age group				
< 60 years	1.00	–	1.00	–
60 years and older	1.04 (0.55; 2.03)	0.895	1.27 (0.64; 2.57)	0.496
Educational level				
Less than 8 years	1.00	–	1.00	–
8 years or higher	3.70 (1.61; 10.06)	0.005	3.84 (1.64; 10.60)	0.005
Associated diseases				
Diabetes	0.95 (0.39; 2.68)	0.919	0.82 (0.32; 2.42)	0.705
Depression	0.40 (0.08; 2.95)	0.299	0.40 (0.07; 3.12)	0.324
Constipation	1.97 (0.84; 5.39)	0.147	1.72 (0.72; 4.80)	0.255

by a reference service in colonoscopy, following the reports of inadequate effluents, patients could be encouraged to keep consuming the laxative, as well as postpone the procedure and contact a nurse or the clinic's department if there are any doubts about adequate preparation.¹²

In this study, we assessed that the subjects with higher levels of education (> 8 years) had a greater chance of accurately evaluating their rectal effluents. This finding differs from what was detected in the works of Harewood, Wright and Baron⁸ and Fatima, Johnson, and Rex,⁹ where both showed that the variables that had higher accuracy were body mass index < 30, use of medications for constipation, use of sodium phosphate as a laxative for preparation, and age below 60 years, respectively.

Conclusion

Reports of dark and thick, dark orange or brown and thick rectal effluents are predictors of inadequate preparation in the endoscopist's assessment.

Conflicts of Interest

The authors declare no conflicts of interest.

References

- 1 Tang SJ, Sones JQ. Colonoscopy atlas of colon polyps and neoplasms. *J Miss State Med Assoc* 2016;57(03):68–71
- 2 Ness RM, Manam R, Hoen H, Chalasani N. Predictors of inadequate bowel preparation for colonoscopy. *Am J Gastroenterol* 2001;96(06):1797–1802

- 3 Harewood GC, Sharma VK, de Garmo P. Impact of colonoscopy preparation quality on detection of suspected colonic neoplasia. *Gastrointest Endosc* 2003;58(01):76–79
- 4 Sulz MC, Kröger A, Prakash M, Manser CN, Heinrich H, Misselwitz B. Meta-analysis of the effect of bowel preparation on adenoma detection: early adenomas affected stronger than advanced adenomas. *PLoS One* 2016;11(06):e0154149
- 5 Voigt J, Mosier M, Gralnek IM. Colonoscopy in poorly prepped colons: a cost effectiveness analysis comparing standard of care to a new cleansing technology. *Cost Eff Resour Alloc* 2021;19(01):25
- 6 Kaminski MF, Regula J, Kraszewska E, et al. Quality indicators for colonoscopy and the risk of interval cancer. *N Engl J Med* 2010;362(19):1795–1803
- 7 Clark BT, Rustagi T, Laine L. What level of bowel prep quality requires early repeat colonoscopy: systematic review and meta-analysis of the impact of preparation quality on adenoma detection rate. *Am J Gastroenterol* 2014;109(11):1714–1723, quiz 1724
- 8 Harewood GC, Wright CA, Baron TH. Assessment of patients' perceptions of bowel preparation quality at colonoscopy. *Am J Gastroenterol* 2004;99(05):839–843
- 9 Fatima H, Johnson CS, Rex DK. Patients' description of rectal effluent and quality of bowel preparation at colonoscopy. *Gastrointest Endosc* 2010;71(07):1244–1252.e2
- 10 So H, Boo SJ, Seo H, et al. Patient descriptions of rectal effluents may help to predict the quality of bowel preparation with photographic examples. *Intest Res* 2015;13(02):153–159
- 11 BBPS Educational Module. (available from: <http://www.cori.org/bbps/>)
- 12 Rex DK. Bowel preparation for colonoscopy: entering an era of increased expectations for efficacy. *Clin Gastroenterol Hepatol* 2014;12(03):458–462