



ORIGINAL ARTICLE

Association between general and abdominal obesity with high blood pressure: difference between genders^{☆,☆☆}



Alison O. Silva^a, Micaelly V. Silva^a, Lisley K.N. Pereira^a,
Wallacy M.N. Feitosa^{a,b}, Raphael M. Ritti-Dias^c, Paula R.B. Diniz^{d,e},
Luciano M.F.T. Oliveira^{a,b,d,e,*}

^a Faculdade Associação Caruaruense de Ensino Superior e Técnico (ASCES), Caruaru, PE, Brazil

^b Grupo de Pesquisa em Saúde Pública (GPESP), Caruaru, PE, Brazil

^c Hospital Israelita Albert Einstein, São Paulo, SP, Brazil

^d Núcleo de Telessaúde, Universidade de Pernambuco (UPE), Recife, PE, Brazil

^e Universidade Federal de Pernambuco (UFPE), Recife, PE, Brazil

Received 8 March 2015; accepted 27 May 2015

Available online 12 December 2015

KEYWORDS

Hypertension;
Adolescent;
Prevalence

Abstract

Objective: To assess the association between general and abdominal obesity with high blood pressure in adolescents of both genders from the public school system.

Methods: This was an epidemiological, descriptive, exploratory study, with a quantitative approach and local scope whose sample consisted of 481 high school students (aged 14–19), selected by using a random cluster sampling strategy. Blood pressure was measured through the use of automated monitor and was considered high when the pressure values were at or above the 95th percentile. The analyses were performed using the chi-squared test and binary logistic regression.

Results: The prevalence of high blood pressure was 6.4%, and it was higher among boys (9.0% vs. 4.7%, $p < 0.05$). There was no significant difference between general ($p = 0.903$) and abdominal obesity ($p = 0.157$) when genders were compared. After adjusting for age, high blood pressure was associated with general ($OR = 6.4$; $p < 0.001$) and abdominal obesity ($OR = 7.0$; $p < 0.001$)

[☆] Please cite this article as: Silva AO, Silva MV, Pereira LK, Feitosa WM, Ritti-Dias RM, Diniz PR, et al. Association between general and abdominal obesity with high blood pressure: difference between genders. J Pediatr (Rio J). 2016;92:174–80.

^{☆☆} Study carried out at Faculdade Associação Caruaruense de Ensino Superior e Técnico (ASCES), Caruaru, PE, Brazil.

* Corresponding author.

E-mail: luciano2308@hotmail.com (L.M.F.T. Oliveira).

PALAVRAS-CHAVE

Hipertensão;
Adolescente;
Prevalência

only among boys, when comparing the fourth quartile with the first quartile of body mass index ($\leq 18.6 \text{ kg/m}^2$ vs. $\geq 23.5 \text{ kg/m}^2$) and waist circumference ($\leq 69 \text{ cm}$ vs. $\geq 80.1 \text{ cm}$).

Conclusion: It was observed that general and abdominal obesity are associated with high blood pressure only in boys, regardless of age.

© 2015 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. All rights reserved.

Associação entre a obesidade geral e abdominal com a pressão arterial elevada: diferença entre gêneros**Resumo**

Objetivo: Analisar a associação entre obesidade geral e abdominal com a pressão arterial elevada em adolescentes de ambos os gêneros da rede de ensino público.

Métodos: Trata-se de um estudo epidemiológico, descritivo, exploratório, com abordagem quantitativa e abrangência municipal cuja amostra foi constituída de 481 estudantes (14-19 anos) do ensino médio, selecionado por meio de uma estratégia de amostragem aleatória de cluster. A pressão arterial foi medida através da utilização de equipamentos automáticos, sendo considerada elevada quando os valores pressóricos estivessem iguais ou acima do percentil 95. As análises foram realizadas através do teste de Qui-quadrado e da regressão logística binária.

Resultados: A prevalência de pressão arterial elevada foi de 6,4%, sendo maior entre os rapazes (9,0% vs. 4,7%, $p < 0,05$). Não foi observada diferença significante entre a obesidade geral ($p = 0,903$) e abdominal ($p = 0,157$) quando comparados os gêneros. Após o ajuste pela idade, a pressão arterial elevada foi associada com a obesidade geral ($OR = 6,4$; $p < 0,001$) e abdominal ($OR = 7,0$; $p < 0,001$) apenas entre os rapazes, quando comparado o quarto quartil com o primeiro quartil do índice de massa corporal ($\leq 18,6 \text{ Kg/m}^2$ Vs $\geq 23,5 \text{ Kg/m}^2$) e da circunferência da cintura ($\leq 69 \text{ cm}$ Vs $\geq 80,1 \text{ cm}$).

Conclusão: Foi observado que a obesidade geral e abdominal está associada com a pressão arterial elevada apenas entre os rapazes, independentemente da idade.

© 2015 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda. Todos os direitos reservados.

Introduction

High blood pressure (HBP) is considered a major risk factor for cardiovascular disease for adults, as well as children and adolescents.^{1,2} Its diagnosis and early treatment can prevent the occurrence of future adverse cardiovascular events, since HBP in childhood is a predictor of hypertension in adulthood.³ In Brazil, the prevalence of HBP in children ranges from 2.5%⁴ to 44.7%;⁵ as it is asymptomatic, the identification and treatment of HBP is often neglected.⁶

By consensus, general and abdominal obesity are considered as predisposing factors for HBP onset⁷; moreover, excess body fat accumulation in the early stages of life is associated with the onset of cardiovascular and metabolic diseases in adulthood.⁸ However, there is evidence demonstrating that body fat distribution is more important than obesity alone.^{9,10} In this sense, it has been observed that abdominal fat accumulation is closely related to HBP, and it is considered the biggest contributor to metabolic complications of the obese population.¹¹

Another factor that can influence blood pressure (BP) values is the difference between genders.^{12,13} Studies have confirmed that boys have higher BP than girls.¹⁴⁻¹⁶ However, the association between obesity and HBP is usually assessed, but the interaction in relation to gender is not tested,

and this factor can distort the found results.^{1,16} Therefore, the objective of this study was to analyze the association between general and abdominal obesity with HBP in adolescents from public schools, while considering the difference between genders.

Methods

This was a descriptive study with a quantitative approach that integrated a school-based cross-sectional epidemiological survey of municipal scope. The sample consisted of students aged 14–19 years, of both genders, enrolled in the state public high schools in the city of Caruaru, state of Pernambuco, Brazil. The total population was estimated at 8833 young individuals distributed in 15 schools, according to the State Secretariat of Education and Culture.

The following parameters were used to calculate sample size: 95% of confidence interval; maximum tolerable error of 2 percentage points; design effect ($deff = 2$); and, because this study comprehended the analysis of multiple risk behaviors and different frequencies of occurrence, the estimated prevalence was defined as 50%. Additionally, to minimize the limitations caused by eventual losses in the application and/or inadequate completion of the questionnaires, it was decided to add 20% to the sample size.

To select the required sample, a two-stage random cluster sampling strategy was used, and "school" and "class" represented, respectively, the sample units in the first and second stage. All public schools in the city of Caruaru were considered eligible for inclusion in the study. In the first stage, school density was used as the stratification criterion in each microregion of the city (Gerência Regional de Educação – Gere), according to their size. In the second stage, the density of classes in the selected schools was considered by school shift (day and night) as a criterion to choose, by drawing lots, those in which the questionnaires would be applied. All students in the selected classes were asked to participate in the study, regardless of their age. After their application, the questionnaires answered by students older than the established age (19 years) were excluded.

Data collection was carried out from June to November of 2014. The questionnaires were collectively applied in the classroom, without the presence of their teachers, and the students were continuously assisted by five researchers (two professors and three undergraduate students), to clarify doubts when filling out the questionnaires. The study was approved by the Ethics Committee in Research with human beings of Faculdade Asces (CAAE-22210913.8.0000.5203/CEP-ASCES: 072403/2013). Subject participation was voluntary and anonymous, and a passive parental consent form was obtained.

Personal information, as well as socioeconomic and sociodemographic variables, were acquired through the translated and adapted version of the Global School-based Student Health Survey (GSHS), proposed by the World Health Organization (WHO). At the anthropometric measurements, all adolescents wore light clothing and were barefoot. Body mass was measured in an electronic Beurer scale (Beurer GmbH, Ulm, Germany) with a maximum capacity of 150 kg and accuracy of 100 g. Height was measured using a portable stadiometer (Wiso, SC, Brazil) with an accuracy of 0.5 cm. Body mass index (BMI) was calculated by dividing body weight in kilograms by the square of height in meters. Waist circumference (WC) was determined as the minimum circumference between the iliac crest and the last rib, using a non-extendable metric tape (Sanny, SP, Brazil) graduated in millimeters. After the calculations of BMI and waist circumference, the data were categorized into quartiles as used in a previous study with adolescents.¹⁷

Blood pressure was measured using an automated oscillometric device (Omron HEM-742, Omron Healthcare, INC, IL, USA) was used, which was previously calibrated and validated for adolescents.¹⁸ Cuff adjustment followed the recommendations of the American Heart Association.¹⁹ Three blood pressure measurements were taken, after a one-minute interval between them, carefully following the protocols recommended by the VI Brazilian Guidelines on Hypertension,²⁰ as well as in relation to the positioning of the arm and body, prior rest, instrument calibration, size of the sphygmomanometer cuff, number of measurements taken, and the choice of the value used to determine prevalence.

Regarding the BP value used, the first measurement was discarded and the mean of the last two measurements was used, both for the systolic and diastolic pressure. The criteria used to classify the subjects were those established by

the National High Blood Pressure Education Program²¹; thus, for the age group 14–17 years, HBP was considered when the systolic blood pressure (SBP) or diastolic blood pressure (DBP) was equal to or greater than the 95th percentile of the Task Force Report High Blood Pressure in Children and Adolescents, 1996 reference tables for age and gender, adjusted for the height percentile of the assessed adolescent. For students aged 18–19, the values used for young adults were considered as parameters.

The final data tabulation was performed using the EpiData program, version 3.1 (Epidata Association, Odense, Denmark), a public domain system, which was also used to carry out the electronic procedures for data entry control through the check function (controls). In order to detect errors, data entry was repeated and, using the duplicate

Table 1 Socioeconomic and demographic characteristics and prevalence of high blood pressure and general and abdominal obesity in high school students from state public schools, Pernambuco, Brazil, 2014.

Variables	Total = 481	
	n	%
<i>Gender</i>		
Boys	221	45.9
Girls	260	54.1
<i>Age (years)</i>		
14–15	183	38.1
16–17	219	45.6
18–19	78	16.3
<i>Place of residence</i>		
Urban	407	84.6
Rural	74	15.4
<i>Skin color</i>		
White	128	26.7
Non-white	351	73.3
<i>Maternal education</i>		
>8 years of schooling	47	11.6
<8 years of schooling	359	88.4
<i>Occupation</i>		
Works	122	25.5
Does not work	357	74.5
<i>High blood pressure</i>		
No	450	93.6
Yes	31	6.4
<i>Body mass index</i>		
Very low risk (≤ 18.6)	121	25.2
Low risk (18.7–20.6)	120	24.9
Moderate risk (20.7–23.4)	120	24.9
High risk (>23.5)	120	24.9
<i>Waist circumference</i>		
Very low risk (≤ 69)	136	28.3
Low risk (69.1–73)	108	22.5
Moderate risk (73.1–80)	123	25.6
High risk (>80.1)	114	23.7




Figure 1 High risk prevalence rates of general ($BMI > 23.5$) and abdominal ($WC > 80.1$) obesity and high blood pressure (per centile $> 95^{\text{th}}$) in high school students of both genders from state public schools, Pernambuco, Brazil, 2014.
* $p < 0.05$.

file comparison function, typing errors were detected and corrected.

Data analysis was performed using SPSS (SPSS Inc. Statistics for Windows, Version 10.0, IL, USA). Frequency distribution was observed in the descriptive analysis. Pearson's chi-squared test was used in the inferential analysis to analyze the isolated association between HBP and general and abdominal obesity, the difference between the BP of boys and girls, and general and abdominal obesity between boys and girls, as well as to analyze the variables in the model, explore the potential confounders, and identify the need for statistical adjustment of the analyses.

In the multivariate analysis, binary logistic regression was used, by estimating the odds ratio (OR) and 95% confidence intervals to express the degree of association between the independent variables (general and abdominal obesity) and the dependent variable (HBP), using adjustment for age, as performed in other studies.^{22,23} In addition to the biological plausibility, the age variable showed, in relation to BP, a statistical significance level < 0.20 ($p < 0.20$), a criterion used for entry in the statistical model using the backward method. After obtaining the predictive variables of the final model, the interaction occurrence was tested. The significance level was set at $p < 0.05$ for all tests.

Skin color, maternal level of schooling, occupation, and place of residence did not enter the statistical model, as they had a statistical significance level < 0.20 in relation to BHP: $p = 0.764$, $p = 0.884$, $p = 0.259$, and $p = 0.526$, respectively.

Results

Nine schools of the public state educational system located in the city of Caruaru were assessed. Of the 569 students who were in class on the day of data collection, 31 adolescents refused to participate and 26 were not allowed to participate in the study by their parents or guardians, totaling 57 refusals. Data were collected from 512 adolescents; however, the final sample, corresponding to students aged

between 14 and 19 years, comprised 481 students, of whom 54.1% were females. The characteristics of the adolescents are shown in Table 1.

The prevalence of HBP was 6.4%, being higher among boys (9.0% vs. 4.7%, $p < 0.05$). There was no significant difference between general ($p = 0.903$) and abdominal ($p = 0.157$) obesity when genders were compared, as shown in Fig. 1.

After adjusting for age, as shown in Table 2, HBP was associated with general ($OR = 6.44$; $p = 0.001$) and abdominal ($OR = 6.97$; $p = 0.001$) obesity only in boys, when comparing the fourth quartile with the first quartile of the BMI ($\leq 18.6 \text{ kg/m}^2$) vs. ($\geq 23.5 \text{ kg/m}^2$) and waist circumference ($\leq 69 \text{ cm}$ vs. ($\geq 80.1 \text{ cm}$)).

Discussion

This study aimed to analyze the association between general and abdominal obesity with HBP in adolescents of both genders. The main findings were: (i) the prevalence of HBP was higher in boys; (ii) there was no significant difference between general and abdominal obesity when comparing genders; and (iii) HBP was associated with general and abdominal obesity only in boys.

The prevalence of HBP observed in this study was 6.4%, lower than that found in other studies with adolescents using the $HBP > 95^{\text{th}}$ percentile as a criterion for HBP diagnosis.^{14,15,24–26} However, methodological differences, number of measurements, and the different reference criteria are the main causes of the great variability found in HBP prevalence among the investigations. In this study, the authors chose to use greater accuracy when diagnosing HBP, considering that a higher methodological stringency based on the hypertension guideline may result in a lower prevalence of HBP.²⁷

As in other studies,^{14–16} boys had higher BP than girls. In this sense, studies using ambulatory BP monitoring techniques in children showed that with increasing age, there is an increase in blood pressure in both boys and girls; however, after puberty onset, BP in boys is higher than that of girls of

Table 2 Raw and adjusted odds ratio between high blood pressure and general and abdominal obesity (quartiles) in high school students of both genders, from state public schools, Pernambuco, Brazil, 2014.

Anthropometric indices	High blood pressure							
	Odds ratio (Raw)	95% CI	p-Value	Overall p-Value	Odds ratio (adjusted ^a)	95% CI	p-Value	Overall p-Value
Boys								
<i>Waist circumference (cm)</i>								
Very low risk (≤ 69)	1				1			
Low risk (69.1–73)	0.46	0.04–5.26	0.535	<0.001	0.38	0.03–4.40	0.439	<0.001
Moderate risk (73.1–80)	1.75	0.31–10.00	0.526		1.31	0.22–7.76	0.765	
High risk (> 80.1)	7.93	1.69–37.18	0.009		6.97	1.47–33.15	0.015	
<i>Body mass index (kg/m²)</i>								
Very low risk (≤ 18.6)	1				1			
Low risk (18.7–20.6)	0.46	0.04–5.19	0.527	<0.001	0.38	0.03–4.43	0.442	<0.001
Moderate risk (20.7–23.4)	2.70	0.50–14.59	0.247		2.31	0.42–12.73	0.337	
High risk (> 23.5)	7.57	1.60–35.70	0.011		6.44	1.33–31.04	0.020	
Girls								
<i>Waist circumference (cm)</i>								
Very low risk (≤ 69)	1				1			
Low risk (69.1–73)	0.38	0.04–3.54	0.399	0.260	0.34	0.04–3.15	0.343	0.245
Moderate risk (73.1–80)	0.33	0.03–3.06	0.331		0.33	0.04–3.10	0.335	
High risk (> 80.1)	2.22	0.60–8.25	0.233		2.23	0.59–8.41	0.235	
<i>Body mass index (kg/m²)</i>								
Very low risk (≤ 18.6)	1				1			
Low risk (18.7–20.6)	0.25	0.03–2.34	0.227	0.462	0.24	0.03–2.24	0.210	0.463
Moderate risk (20.7–23.4)	0.23	0.03–2.16	0.200		0.21	0.02–2.00	0.174	
High risk (> 23.5)	1.52	0.41–5.68	0.529		1.54	0.41–5.82	0.524	

CI, confidence interval.

^a Adjusted for age.

the same age.²⁸ Corroborating these findings, a longitudinal study of 1267 adolescents found that male adolescents were more likely to develop HBP in their lifetime.¹²

One of the major risk factors for hypertension is general obesity.^{2,16} A systematic review with meta-analysis showed that obese children had blood pressure levels approximately 40% higher than children with normal weight.² Regarding body fat location, an association between general and abdominal obesity with HBP, through BMI and WC, was observed in a study with adolescents; however, sample stratification by gender was not performed at the analysis and the interaction between variables was not tested.²⁴

Usually, there is a correlation between a possible increase in blood pressure in boys and a higher prevalence of fat accumulation in the abdominal region, but curiously, even with no difference between general and abdominal obesity when comparing genders, it was observed that the HBP was associated with general and abdominal obesity only in boys. These results are similar to those observed by Payeb et al.²³ In a study carried out in Iran with 13,486 adolescents, those authors found a higher percentage of obesity in girls, although the HBP values were higher in boys.

A possible explanation for the higher HBP in boys is that sex hormones play an important role in BP regulation,²⁹ as increased levels of gonadal hormones may occur during childhood and adolescence and these can modify the

synthesis, release, and bioactivity of these factors, inducing vascular relaxation and consequently influencing BP.¹³ Moreover, Landazuri et al.³⁰ verified that the levels of angiotensin-converting enzyme (ACE) were increased in boys aged 11–17 years, while they were decreased in girls of the same age group. ACE was the main enzyme of the renin-angiotensin system, which in turn plays an important role in heart function and BP regulation. The significant reduction in ACE activity observed in girls, but not in boys, indicates that estradiol and testosterone may have a diverse influence on angiotensin converting enzyme (ACE) activity in humans, particularly in relation to HBP.

The study has strengths and limitations that deserve to be mentioned. The cross-sectional design and the correlative nature of the data prevent us from establishing a causal association between HBP and obesity. Moreover, the adolescents' maturational age, a point which may be related to hormonal release, was not assessed. Among the strengths, this study had a representative sample; the sampling procedures were established to ensure that the sample consisted of adolescent students that attended schools at different school shifts, and methodological stringency was employed when collecting BP data.

According to the data obtained during the study, it was observed that even with no difference in fat distribution between genders, general and abdominal obesity are

associated with HBP only in boys, regardless of age. The results show the need to stratify the sample by gender when the aim is assessing BP in adolescents. Additionally, it was observed that HBP in boys may not be related to the prevalence of general and abdominal fat accumulation. In this sense, it is recommended to carry out studies using a longitudinal design to assess the influence of sex hormones on BP in adolescents.

Funding

The research was funded by Faculdade ASCES Scientific Initiation Program.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Singhal V, Agal P, Kamath N. The prevalence of elevated blood pressure and the association of obesity in asymptomatic female adolescent offsprings of hypertensive and normotensive parents. *J Clin Diagn Res.* 2012;6:1158–61.
2. Friedemann C, Heneghan C, Mahtani K, Thompson M, Perera R, Ward AM. Cardiovascular disease risk in healthy children and its association with body mass index: systematic review and meta-analysis. *BMJ.* 2012;345:e4759.
3. Li S, Chen W, Srinivasan SR, Bond MG, Tang R, Urbina EM, et al. Childhood cardiovascular risk factors and carotid vascular changes in adulthood: the Bogalusa Heart Study. *JAMA.* 2003;290:2271–6.
4. de Rezende DF, Scarpelli RA, de Souza GF, da Costa JO, Scarpelli AM, Scarpelli PA, et al. Prevalence of systemic hypertension in students aged 7 to 14 years in the municipality of Barbacena, in the State of Minas Gerais, in 1999. *Arq Bras Cardiol.* 2003;81:381–6.
5. de Araújo TL, de Lopes MV, Cavalcante TF, Guedes NG, Moreira RP, Chaves ES, et al. Análise de indicadores de risco para hipertensão arterial em crianças e adolescentes. *Rev Esc Enferm USP.* 2008;42:120–6.
6. Brasil, Ministério da Saúde. Hipertensão arterial sistêmica para o Sistema Único de Saúde. (Série A. Normas e Manuais Técnicos). In: Secretaria de Atenção à Saúde Departamento de Atenção Básica, editor. Brasília: Ministério da Saúde; 2006. pp. 7–84.
7. Vasques C, Mota M, Correia T, Lopes V. Prevalence of overweight/obesity and its association with sedentary behavior in children. *Rev Port Cardiol.* 2012;31:783–8.
8. Williams SM. Weight and height growth rate and the timing of adiposity rebound. *Obes Res.* 2005;13:1123–30.
9. Hasselmann MH, Faerstein E, Werneck GL, Chor D, Lopes CS. Associação entre circunferência abdominal e hipertensão arterial em mulheres: Estudo Pró-Saúde. *Cad Saude Publica.* 2008;24:1187–91.
10. Jardim PC, Gondim Mdo R, Monego ET, Moreira HG, Vitorino PV, Souza WK, et al. Hipertensão arterial e alguns fatores de risco em uma capital brasileira. *Arq Bras Cardiol.* 2007;88: 452–7.
11. de Menezes TN, Rocha FL, de Oliveira Belém PL, Pedraza DF. Obesidade abdominal: revisão crítica das técnicas de aferição e dos pontos de corte de indicadores antropométricos adotados no Brasil. *Cien Saude Colet.* 2014;19:1741–54.
12. Dasgupta K, O'Loughlin J, Chen S, Karp I, Paradis G, Tremblay J, et al. Emergence of sex differences in prevalence of high systolic blood pressure. *Circulation.* 2006;114:2663–70.
13. Khalil RA. Sex hormones as potential modulators of vascular function in hypertension. *Hypertension.* 2005;46:249–54.
14. Barros MV, Ritti-Dias RM, Honda Barros SS, Mota J, Andersen LB. Does self-reported physical activity associate with high blood pressure in adolescents when adiposity is adjusted for? *J Sports Sci.* 2013;31:387–95.
15. Christofaro DG, Ritti-Dias RM, Chiolero A, Fernandes RA, Casonatto J, de Oliveira AR. Physical activity is inversely associated with high blood pressure independently of overweight in Brazilian adolescents. *Scand J Med Sci Sports.* 2013;23: 317–22.
16. Hoffmann M, Pio da Silva AC, Siviero J. Prevalência de hipertensão arterial sistêmica e interrelações com sobrepeso, obesidade, consumo alimentar e atividade física, em estudantes de escolas municipais de Caxias do Sul. *Pediatria (São Paulo).* 2010;32:163–72.
17. Kuschnir MC, Mendonça GA. Risk factors associated with arterial hypertension in adolescents. *J Pediatr (Rio J).* 2007;83: 335–42.
18. Christofaro DG, Fernandes RA, Gerage AM, Alves MJ, Polito MD, Oliveira AR. Validação do monitor de medida de pressão arterial O mron HEM 742 em adolescentes. *Arq Bras Cardiol.* 2009;92:9–14.
19. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al. Recommendations for blood pressure measurement in humans and experimental animals. *Circulation.* 2005;111:697–716.
20. Sociedade Brasileira de Cardiologia, Sociedade Brasileira de Hipertensão, Sociedade Brasileira de Nefrologia. VI Diretrizes Brasileiras de Hipertensão Arterial. *Arq Bras Cardiol.* 2010;95:1–51.
21. National High Blood Pressure Education Program Working group on High Blood Pressure in Children and Adolescents. The Fourth Report on the diagnosis, evaluation and treatment of high blood pressure in children and adolescents. *Pediatrics.* 2004;114:555–76.
22. Flechtnner-Mors M, Neuhauser H, Reinehr T, Roost H-P, Wiegand S, Siegfried W, et al. Blood pressure in 57,915 pediatric patients who are overweight or obese based on five reference systems. *Am J Cardiol.* 2015;115:1587–94.
23. Payab M, Kelishadi R, Qorbani M, Motlagh ME, Ranjbar SH, Ardalan G, et al. Association of junk food consumption with high blood pressure and obesity in Iranian children and adolescents: the CASPIAN-IV study. *J Pediatr (Rio J).* 2015;91: 196–205.
24. Christofaro DG, Ritti-Dias RM, Fernandes RA, Polito MD, Andrade SM, Cardoso JR, et al. Detecção de hipertensão arterial em adolescentes através de marcadores de adiposidade geral e abdominal. *Arq Bras Cardiol.* 2011;96:465–70.
25. Gomes JC. O atual ensino da ética para os profissionais de saúde e seus reflexos no cotidiano do povo brasileiro. *Revista Bioética.* 2009;4:53–64.
26. Griz LH, Viégas M, Barros M, Griz AL, Freese E, Bandeira F. Prevalence of central obesity in a large sample of adolescents from public schools in Recife, Brazil. *Arq Bras Endocrinol Metabol.* 2010;54:607–11.
27. Magalhaes MG, Oliveira LM, Christofaro DG, Ritti-Dias RM. Prevalência de pressão arterial elevada em adolescentes brasileiros e qualidade dos procedimentos metodológicos empregados: revisão sistemática. *Rev Bras Epidemiol.* 2013;16:849–59.

28. Jackson LV, Thalange NK, Cole TJ. Blood pressure centiles for Great Britain. *Arch Dis Child.* 2007;92:298–303.
29. Reckelhoff JF, Fortepiani LA. Novel mechanisms responsible for postmenopausal hypertension. *Hypertension.* 2004;43:918–23.
30. Landazuri P, Granobles C, Loango N. Gender differences in serum angiotensin-converting enzyme activity and blood pressure in children: an observational study. *Arq Bras Cardiol.* 2008;91:382–8.