

Carlos Alberto Velasco-Benítez \*,  
Herney Andrés García-Perdomo 

Universidad del Valle, Cali, Colombia

\* Corresponding author.

E-mail: [carlos.velasco@correounivalle.edu.co](mailto:carlos.velasco@correounivalle.edu.co)  
(C.A. Velasco-Benítez).

<https://doi.org/10.1016/j.jpeds.2018.11.011>  
0021-7557/

© 2019 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Answer to Letter to the Editor “Some flaws that could change the effect sizes and limit the extrapolation of these results”<sup>☆</sup>



## Resposta à Carta ao Editor “Alguns pontos fracos que podem alterar os tamanhos dos efeitos e limitar a extrapolação desses resultados”

Dear Editor,

We appreciate the attention of those who read and analyzed our work; we found the comments and considerations made about the recently published article very interesting. Please find below the answers to the main points raised in the Letter to the Editor:

Constipation is an increasingly present condition found in the pediatric population. It can be classified as organic, when it is associated with intestinal and extra intestinal disorders, and as functional, when there are no warning symptoms or organic causes, which occurs more frequently.<sup>1</sup> The pathophysiology of functional constipation is not well understood, but it is known to be multifactorial. Recently, the diagnostic criteria for functional constipation were updated and redefined in the Rome IV consensus, aiding in the differentiation between functional constipation and constipation with an organic cause.<sup>2</sup>

Intestinal constipation is a frequent disease in the pediatric population and the initial management often includes advice on increasing the fiber intake. However, there have been few studies that have proven the efficacy of fiber use in the treatment of childhood intestinal constipation. The European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN), in its Consensus on the management of childhood intestinal constipation in 2014, for the first time emphasized that there was no evidence to justify prescribing an increase in fiber consumption.<sup>3</sup> Aiming to gather more current evidence on the use of fiber in the treatment of constipation in pediatric patients, a systematic review with meta-analysis was proposed.

The search strategy is one of the most relevant aspects in a systematic review. We decided to implement a comprehensive strategy, including multiple terms to define the study population and study design. We did not include endpoint criteria in our search terms, in order to increase sensitivity. Moreover, we searched for articles published in five different languages (Portuguese, English, Spanish, French, and German) and in four databases (PubMed, Embase, LILACS, and the Cochrane Library), above what is recommended in the Cochrane Handbook.

Additionally, our search in the gray literature occurred through the evaluation of references in the 42 articles that were fully read. We did not search smaller databases or non-indexed poster publications in the main databases because this would have required resources beyond those available to us, with low probability of finding large studies with good methodological quality.

The lack of publication in PROSPERO (<https://www.crd.york.ac.uk/prospero>) is a real limitation of our study; however, its absence does not prevent the use of the results in clinical practice, since our search was broad.

The assessment of the quality of evidence selected for this systematic review with meta-analysis was performed using the Cochrane Collaboration tool and the Jadad Scale. This evaluation determines the strength of the recommendation, but also represents the degree of confidence on which to base a decision or recommendation.<sup>4</sup> We understand that the use of these two tools for bias evaluation is not entirely necessary, but there is no loss for the reader in having this information available.

Unfortunately, because this is a field with few studies and low methodological consistency, we chose not to exclude studies based on diagnostic criteria for functional constipation. In fact, the current Rome IV criteria, published in 2016, were not used by any of the studies. Moreover, the type of intervention varied in the studies, which further exacerbates the outcome heterogeneity. Therefore, we decided to include all the studies in the final analysis because they showed a similar patient profile.

The analysis of sensitivity through one-by-one exclusion of the studies, published in the article, found consistency of the main results in the outcomes of bowel movement frequency, fecal consistency, therapeutic success, abdominal pain, and fecal incontinence outcomes. We then performed a sensitivity analysis excluding articles that did not use the Rome diagnostic criteria in their research, as suggested by Kokke et al.<sup>5</sup> and Loening-Baucke et al.<sup>6</sup>; once again, no significant alteration in the analyzed outcomes was found.

We still think that large studies with clear inclusion criteria, continuous follow-up, and well-defined treatment are

<sup>☆</sup> Please cite this article as: Mello PP. Answer to Letter to the Editor “Some flaws that could change the effect sizes and limit the extrapolation of these results”. *J Pediatr (Rio J)*. 2019;95:375–6.

necessary to define the real therapeutic power of fiber in constipation. Although there is a risk of a beta-error, given that we do not detect an actual fiber efficacy, currently we do not believe there is sufficient evidence to suggest the use of fiber. The results of our meta-analysis are important to define this fact and stimulate the production of further evidence in this field.

### Conflicts of interest

The author declares no conflicts of interest.

### References

1. Morais MB, Maffei HV. Constipation. *J Pediatr (Rio J)*. 2000;76:5147–56.
2. Drossman DA. Functional gastrointestinal disorders: history, pathophysiology, clinical features and Rome IV. *Gastroenterology*. 2016;150:1262–79.
3. Tabbers MM, DiLorenzo C, Berger MY, Faure C, Langendam MW, Nurko S, et al. Evaluation and treatment of functional constipation in infants and children: evidence-based recommendations from ESPGHAN and NASPGHA. *J Pediatr Gastroenterol Nutr*. 2014;58:258–74.
4. Brasil. Ministério da Saúde. Secretaria de Ciência, Tecnologia e Insumos Estratégicos. Departamento de Ciência e Tecnologia. Diretrizes metodológicas: elaboração de revisão sistemática e metanálise de ensaios clínicos randomizados. Normas e Manuais Técnicos. Brasília: Ministério da Saúde; 2012.
5. Kokke FT, Scholtens PA, Alles MS, Decates TS, Fiselier TJ, Tolboom JJ, et al. A dietary fiber mixture versus lactulose in the treatment of childhood constipation: a double-blind randomized controlled trial. *J Pediatr Gastroenterol Nutr*. 2008;47:592–7.
6. Loening-Baucke V, Miele E, Staiano A. Fiber (glucomannan) is beneficial in the treatment of childhood constipation. *Pediatrics*. 2004;113:259–64.

Patricia Piccoli de Mello  a,b,c

<sup>a</sup> Hospital Moinhos de Vento, Porto Alegre, RS, Brazil

<sup>b</sup> Grupo Hospitalar Conceição, Porto Alegre, RS, Brazil

<sup>c</sup> Universidade Federal do Rio Grande do Sul (UFRGS), Saúde da Criança e do Adolescente, Porto Alegre, RS, Brazil

E-mail: [patriciamellomed@gmail.com](mailto:patriciamellomed@gmail.com)

<https://doi.org/10.1016/j.jped.2019.02.002>

0021-7557/

© 2019 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).