



ORIGINAL ARTICLE

# Birth weight and its association with blood pressure and nutritional status in adolescents<sup>☆,☆☆</sup>



Vanessa Roriz Ferreira<sup>a</sup>, Thiago Veiga Jardim<sup>a,b,\*</sup>, Thais Rolim Póvoa<sup>c</sup>,  
Karla Lorena Mendonça<sup>a</sup>, Flávia Nogueira Nascente<sup>a</sup>, Carolina Sousa Carneiro<sup>a</sup>,  
Weimar Sebba Barroso<sup>a</sup>, Polyana Morais<sup>a</sup>, Maria Gondim Peixoto<sup>a</sup>,  
Ana Luiza Lima Sousa<sup>a</sup>, Paulo Cesar Brandão Veiga Jardim<sup>a</sup>

<sup>a</sup> Universidade Federal de Goiás (UFG), Liga de Hipertensão Arterial, Goiânia, GO, Brazil

<sup>b</sup> Brigham & Women's Hospital, Division of Cardiovascular Medicine, Boston, United States

<sup>c</sup> Universidade Estadual de Goiás (UEG), Escola Superior de Educação Física e Fisioterapia do Estado de Goiás (ESEFFEGO), Goiânia, GO, Brazil

Received 9 January 2017; accepted 22 March 2017

Available online 24 August 2017

## KEYWORDS

Birth weight;  
Body mass index;  
Waist circumference;  
Height;  
Blood pressure

## Abstract

**Objective:** The management of children with low birth weight is not the same in countries with different resources. The authors assessed the association of birth weight with blood pressure and nutritional status in a representative sample of adolescents from a Brazilian state, aiming to identify possible consequences of these differences.

**Methods:** A cross-sectional school-based study was conducted with adolescents (12–18 years) enrolled in public and private schools. Birth weight, office blood pressure, home blood pressure measurements, and nutritional status (body mass index, height z-score for the age, and waist circumference) were assessed. The association of birth weight with the outcomes (blood pressure, height, body mass index, and waist circumference) was studied through univariate and multivariable linear regression models.

**Results:** A total of 829 adolescents with a mean age of  $14.6 \pm 1.62$  years were included; 43.3% were male, and 37.0% from private schools. The prevalence of low birth weight was 8.7%. Mild low height prevalence was higher among those adolescents with low/insufficient birth weight when compared to those with normal/high birth weight (11.7 vs. 4.2%;  $p < 0.001$ ). In the multiple linear regression analysis, for each increase of 100 g in birth weight, height increased by 0.28 cm (95% CI: 0.18–0.37;  $p < 0.01$ ). Birth weight did not influence office blood pressure and home blood pressure, body mass index, or waist circumference of adolescents.

<sup>☆</sup> Please cite this article as: Ferreira VR, Jardim TV, Póvoa TR, Mendonça KL, Nascente FN, Carneiro CS, et al. Birth weight and its association with blood pressure and nutritional status in adolescents. J Pediatr (Rio J). 2018;94:184–191.

<sup>☆☆</sup> Study conducted at Universidade Federal de Goiás (UFG), Liga de Hipertensão Arterial, Goiânia, GO, Brazil.

\* Corresponding author.

E-mail: thiagolairin@hotmail.com (T.V. Jardim).

**PALAVRAS-CHAVE**

Peso ao nascer;  
Índice de massa corporal;  
Circunferência da cintura;  
Estatura;  
Pressão arterial

**Conclusions:** Birth weight was directly associated to height, but not associated to blood pressure, body mass index, and waist circumference in adolescents from an urban area of a developing country.

© 2017 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Peso ao nascer e sua associação com pressão arterial e estado nutricional em adolescentes

#### Resumo

**Objetivo:** O manejo de crianças com baixo peso ao nascer não é o mesmo em países com diferentes recursos. Investigamos a associação do peso ao nascer com a pressão arterial e o estado nutricional em uma amostra representativa de adolescentes de uma capital brasileira com o objetivo de identificar possíveis consequências destas diferenças.

**Métodos:** Estudo transversal de base escolar conduzido com adolescentes (12-18 anos) matriculados em escolas públicas e privadas. Investigou-se o peso ao nascer, a pressão arterial, e o estado nutricional, por meio do índice de massa corporal, do escore z de estatura para idade e da circunferência da cintura.

**Resultados:** Um total de 829 adolescentes com uma idade média  $14.6 \pm 1.62$  anos foram incluídos, 43.3% do sexo feminino e 37.0% de escolas privadas. A prevalência de baixo peso ao nascer foi 8.7%. Baixa estatura leve foi mais prevalente nos adolescentes com peso ao nascer baixo/insuficiente ( $11.7 \times 4.2\%$  -  $p < 0.001$ ). Na análise de regressão linear múltipla, para cada aumento de 100g no peso ao nascer, a estatura aumentou em 0.28 cm (IC 95% = 0.18-0.37;  $p < 0.01$ ). O peso ao nascer não influenciou a pressão arterial (casual e residencial), o índice de massa corporal e a circunferência da cintura dos adolescentes.

**Conclusões:** O peso ao nascer esteve diretamente associado à altura, mas não associado à pressão arterial, índice de massa corporal e circunferência da cintura em adolescentes de uma área urbana de um país em desenvolvimento.

© 2017 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The hypothesis of fetal origin of diseases or “programming” was first proposed by Barker, in 1990.<sup>1</sup> The main concept of this hypothesis is that the stress caused by fetal malnutrition, whose main biomarker is low birth weight (LBW), would inalterably modify the physiology or metabolism of an organ, increasing disease susceptibility in adult life.

Undoubtedly, uterine life is a critical period of development, due to rapid growth, cellular differentiation, and functional maturation of the organs. These rather sensitive processes would be particularly susceptible to alterations in the nutrient medium.<sup>1</sup>

Birth weight is an important health indicator that reflects the social and economic conditions of women during pregnancy. Together with prematurity, birth weight is the main factor associated with neonatal death in Brazil. Babies with LBW present greater morbidity and mortality in the first year of life.<sup>2</sup>

LBW children (BW <2500 g) can be born at term or before term, and have varying degrees of social and medical risk. Because this is not a homogeneous group, patients have a

broad spectrum of growth, health, and developmental outcomes. While the vast majority of LBW children present normal outcomes, as a group they generally have higher rates of subnormal growth, illnesses, and neurodevelopmental problems.<sup>3</sup>

Recent studies showed that individuals with LBW due to inadequate intrauterine development are prone to developing hypertension, obesity, and low height, among other conditions, in late stages of life.<sup>4-6</sup>

Even though the association of LBW with a number of conditions has been demonstrated in different contexts,<sup>7-9</sup> the mechanisms behind these associations are still not completely understood and need to be more investigated.<sup>10</sup> Furthermore, few publications in middle-income countries focused on this matter. Considering that the management of LBW children is not the same in countries with different resources, research conducted in these countries can help to identify the outcomes related to LBW in adolescence and adult life.

Due to the scarcity of studies on fetal programming of diseases in middle-income countries such as Brazil, particularly considering the youth population, this article aimed

to assess the association of birth weight with blood pressure (BP) and nutritional status in a representative sample of adolescents from a Brazilian state capital.

## Methods

A cross-sectional school-based study was conducted with a representative sample of adolescents enrolled in public and private schools of Goiania, a state capital in the Midwest region of Brazil. The study population was composed by adolescents, aged between 12 and 18 years. Schools were selected by probabilistic conglomerate sampling; students were randomly selected and stratified by age and gender. Exclusion criteria were chronic disease, use of medication that could influence on BP, physical disability that would not allow anthropometric evaluation, pregnancy, and absence of information on birth weight.

The study was approved by the institution's ethics committee (protocol no. 017/2010). The eligible adolescents who agreed to participate in the study signed an informed consent form and so did their parents/guardians.

A total of 1.221 adolescents who met the inclusion criteria were invited to participate in the study. Forty-eight refused (3.8%), four (0.5%) did not have the anthropometric measurements performed, and 340 (41.0%) did not provide information about birth weight; they were excluded, resulting in 829 adolescents (67.9% of the original group) in the final sample.

The sample size allowed to estimate the association of birth weight with anthropometric variables and BP in adolescents, detecting a difference of 15% for boys and girls separately, with a two-tailed  $\alpha$  of 0.05 and a test power of 80%.

The instrument for data collection was a standardized questionnaire, previously tested in a pilot study,<sup>11</sup> with questions regarding the individual's identification, including gender, age, skin color (white or non-white), and family history of hypertension and obesity, as well as socioeconomic classification of the adolescent.

## Anthropometric measures

Nutritional status was evaluated through weight, height and waist circumference (WC), using standardized procedures.<sup>12</sup> Weight was measured on a Kratos<sup>®</sup> electronic scale (Kratos<sup>®</sup>, SP, Brazil), with capacity of 150 kg and variation of 50 g and height was assessed with Secca<sup>®</sup> stadiometer (Secca<sup>®</sup>, SP, Brazil), graded to an accuracy of 0.1 cm. Adolescents were classified based on the height z-score for the age: mild low height ( $-1 < z\text{-score} \leq -2$ ) and normal height ( $z\text{-score} \geq -1$ ).<sup>13</sup> Body mass index (BMI) was calculated and classified according to the specific standard reference for age and gender proposed by the World Health Organization (WHO).<sup>14</sup> WC measurements were performed with a Sanny<sup>®</sup> (Sanny<sup>®</sup>, SP, Brazil) 200 cm inextensible tape with 0.1 cm variation, and the cut off points were those proposed by Taylor et al.<sup>15</sup>; students were categorized as normal and increased WC.

## Birth weight

In the study presentation letter, parents or guardians were required to answer whether the birth was premature or not and to write the birth weight, according to data from the children's birth card. Birth weight was classified as low ( $BW < 2500$  g), insufficient ( $2500 \text{ g} \leq BW < 3000$  g), normal ( $3000 \text{ g} \leq BW < 4000$  g), or high ( $BW \geq 4000$  g).<sup>12</sup> Preterm birth was used to adjust birth weight in the regression model.

## Office blood pressure (OBP) measurement

Measurements were performed in school, with the OMRON<sup>®</sup> HEM-705CP semiautomatic device (HEM-705CP OMRON<sup>®</sup>, CA, USA), and cuffs in three different sizes were used, according to the right arm circumference. The equipment was previously validated for adolescents.<sup>16</sup> Two BP measurements following the proper techniques<sup>17</sup> were performed in two different encounters within one-week interval. For the analysis, the mean values of the second measurements at each time point were considered. High OBP was defined as values  $\geq 95$ th percentile for sex, age, and height.<sup>17</sup>

## Home blood pressure (HBP) measurement

The same equipment and cuffs sizes for OBP were used. The device was given to the adolescents, who were also properly trained on how to use it. Two right arm measurements should be performed, in sitting position, after 5 min of rest, in the morning (between 6 and 10 am) and in the evening (between 6 and 10 pm), for six consecutive days (total of 24 readings).<sup>18</sup>

Home measurements saved in the memory of the equipment were printed and compared with those reported in the HBP form filled by the adolescent. When there was a discrepancy between measures (more measures from the equipment memory than from the HBP form) or when the adolescent reported not being the only person to have used the equipment, the exam was excluded from the analysis. Exams were considered valid when at least 12 measurements (50%) were performed according to standardized protocols.<sup>18</sup> HBP was classified as high when the mean systolic and/or diastolic pressure of the six days was  $\geq 95$ th percentile for age, gender, and height.<sup>19</sup>

## Statistical analysis

Data were tabulated in double entry with the Epi-Info (Epi-Info, version 6.04, Division of Health Informatics & Surveillance, Centers of Disease Control and Prevention). Variable categorization and statistical analysis were performed with the STATA software (SPSS for Windows, version 13, Chicago, USA). Kolmogorov-Smirnov's test was used to assess whether continuous variables had normal distribution. Pearson's chi-squared test was used to compare BP and nutritional status categories between low/insufficient and normal/high birth weight adolescents. The birth weight relation with the outcomes (BP, height, BMI, and WC) was studied through univariate and multivariable linear regression

**Table 1** Overall characteristics of a representative sample of adolescents from a Brazilian state capital. Goiânia, Brazil.

Variable	n	Value
Sex (male), n (%)	829	359 (43.3%)
School type (private), n (%)	829	307 (37.0%)
Skin color (white), n (%)	829	397 (47.9%)
Age (years), mean (SD)	829	14.68 (1.62)
Birth weight (g), mean (SD)	829	3277.93 (597.33)
Weight (kg), mean (SD)	829	57.36 (13.52)
Height (m), mean (SD)	829	1.66 (0.09)
Body mass index (kg/m <sup>2</sup> ), mean (SD)	829	20.79 (3.92)
Waist circumference (cm), mean (SD)	829	70.81 (9.09)
Office SBP <sup>a</sup> (mmHg), mean (SD)	777	112.15 (12.50)
Office DBP <sup>b</sup> (mmHg), mean (SD)	777	66.61 (8.03)
Home SBP <sup>c</sup> (mmHg), mean (SD)	701	112.82 (10.21)
Home DBP <sup>c</sup> (mmHg), mean (SD)	701	66.59 (6.36)
Preterm birth, n (%)	829	55 (6.7%)
Family history of hypertension, n (%)	829	216 (26.1%)
Family history of obesity, n (%)	829	80 (9.7%)

<sup>a</sup> Mean systolic blood pressure (SBP) values of second measurements in different moments (one week apart).

<sup>b</sup> Mean diastolic blood pressure values (DBP) of second measurements in different moments (one week apart).

<sup>c</sup> Mean systolic blood pressure values of all home blood pressure measurements.  
SD, standard deviation.


analysis. Variables that presented *p*-value <0.20 in the univariate analysis and absence of collinearity between independent variables (Pearson’s correlation coefficient <0.80) were included in the multivariable model. The variables included in the model for adjustment were selected based on scientific plausibility.<sup>20</sup> A significance level of 5% was considered.

## Results

A total of 829 adolescents with a mean age of 14.68 (±1.62) years were included in the analysis, 43.3% were male and 37.0% from private schools. OBP measurements were not performed in 53 (6.4%) subjects and 128 (15.4%) did not perform a valid HBP measurement. The mean birth weight was 3277.93 g (±597.33) and 47.9% referred to themselves as being white. The overall characteristics of the group are shown in Table 1.

The prevalence of LBW was 8.7%. The distribution among the birth weight categories was homogenous between males and females, and the only difference was found in the high birth weight category, in which the proportion of males was higher than females, as shown in Fig. 1.

Table 2 presents the nutritional status, characterized by BMI, height, and WC distribution, and stratified by low/insufficient and normal/high birth weight. The overall overweight/obesity prevalence was 22.9%. The prevalence of mild low height was higher among those adolescents with low/insufficient birth weight when compared to those



**Figure 1** Birth weight category distribution in the overall sample and stratified by sex in adolescents. Goiânia, Brazil.

<sup>a</sup>Difference between male and female – statistically significant at  $\alpha = 0.05$ .

with normal/high birth weight (11.7 vs. 4.2%; *p* < 0.001). The prevalence of adolescents with abdominal obesity was 15.7%.

The prevalence of high BP, defined by OBP, was 8.5%; in turn, when this prevalence was defined by HBP, the rate was 3.8%. Table 2 shows the prevalence of hypertension stratified by the measurement technique and the comparison between birth weight categories. No difference in hypertension prevalence was found among the birth weight categories.

In the studied sample, birth weight did not influence OBP and HBP. A positive influence was detected initially on BMI and WC; however, it was not sustained after adjusted analysis (*p* = 0.274 and 0.835, respectively). In turn, for each 100 g increase in birth weight, height increased by 0.33 cm (95% CI: 0.23–0.43; *p* < 0.01) and, after multivariable analysis, the increase went to 0.28 cm (95% CI: 0.18–0.37; *p* < 0.01). The birth weight results were adjusted to prematurity. The independent variables used in the model explained 37.6% of height variation (Table 3).

## Discussion

This study assessed a representative sample of more than 800 secondary students from a Brazilian capital in the Midwest region of the country focusing on LBW and its association with BP and nutritional status. Birth weight was a predictor of height, but not of BMI and BP in adolescents from an urban area of Brazil.

The sample’s LBW rate (8.7%) was higher than the 6.1% from another Brazilian sample<sup>21</sup> and the 6.9% found in a previously published study in Goiânia.<sup>22</sup> This finding has a negative impact on newborn’s health and survival. However, higher LBW prevalence can be observed in more developed regions of Brazil; this phenomenon can be explained by the improvement in health assistance for pregnant women and infants, allowing babies with low weight to survive.<sup>23</sup>

High birth weight prevalence (9.0%) was also superior to the 6.5% previously reported for newborns in Goiânia.<sup>24</sup> Since fetal macrosomia is related to a higher risk of death and prematurity, as well as complications such as obstetrical trauma, dystocia, and neonatal hypoglycemia,<sup>25</sup> further

**Table 2** Distribution of nutritional status and blood pressure levels by birth weight categories in adolescents. Goiânia, Brazil.

Variable	<i>n</i>	Total sample <i>n</i> (%)	<i>n</i>	Low/insufficient birth weight <i>n</i> (%)	<i>n</i>	Normal/high birth weight <i>n</i> (%)	<i>p</i> -value <sup>a</sup>
<b>BMI</b>	829		230		599		0.285
Low		30 (3.6)		13 (5.6)		17 (2.8)	
Normal weight		609 (73.5)		166 (72.2)		443 (74.0)	
Overweight		127 (15.3)		34 (14.8)		93 (15.5)	
Obesity		63 (7.6)		17 (7.4)		46 (7.7)	
<b>Height</b>	829		230		599		<0.001
Normal		777 (93.7)		203 (88.3)		574 (95.8)	
Mild low		52 (6.3)		27 (11.7)		25 (4.2)	
<b>WC</b>	829		230		599		0.386
Normal		699 (84.3)		198 (86.1)		501 (83.6)	
Increased		130 (15.7)		32 (13.9)		98 (16.4)	
<b>OBP</b>	776		214		562		0.134
<95th percentile		710 (91.5)		201 (93.9)		509 (90.6)	
≥95th percentile		66 (8.5)		13 (6.1)		53 (9.4)	
<b>HBP</b>	701		188		513		0.436
<95th percentile		674 (96.1)		179 (95.2)		495 (96.5)	
≥95th percentile		27 (3.8)		9 (4.8)		18 (3.5)	
<b>OBP and HBP</b>	816		224		592		0.505
<95th percentile		805 (98.6)		220 (98.2)		585 (98.8)	
≥95th percentile		11 (1.4)		4 (1.8)		7 (1.2)	
<b>OBP or HBP</b>	816		224		592		0.239
<95th percentile		734 (89.9)		206 (92.0)		528 (89.2)	
≥95th percentile		82 (10.1)		18 (8.0)		64 (10.8)	

BMI, body mass index; WC, waist circumference; OBP, office blood pressure; HBP, home blood pressure.

<sup>a</sup> Pearson's chi-square test.

**Table 3** Association of birth weight with blood pressure and nutritional status in adolescents. Goiânia, Brazil, 2010–2011 (*n* = 829).

Variables <sup>a</sup>	$\beta$ simple (95% CI)	<i>p</i> -value <sup>b</sup>	$\beta$ adjusted (95% CI)	<i>R</i> <sup>2</sup>	<i>p</i> -value <sup>c</sup>
<b>OBP (mmHg)<sup>d</sup></b>					
Systolic	−0.01 (−0.16–0.14)	0.872	–	–	–
Diastolic	0.03 (−0.06–0.13)	0.466	–	–	–
<b>HBP (mmHg)<sup>e</sup></b>					
Systolic	0.08 (−0.05–0.20)	0.240	–	–	–
Diastolic	−0.01 (−0.08–0.07)	0.881	–	–	–
<b>BMI (kg/m<sup>2</sup>)</b>	0.05 (0.01–0.09)	0.023	0.03 (−0.02–0.07) <sup>f</sup>	0.095	0.274
<b>WC (cm)</b>	0.13 (0.03–0.23)	0.013	−0.01 (−0.09–0.11) <sup>f</sup>	0.174	0.835
<b>Height (cm)</b>	0.33 (0.23–0.43)	<0.001	0.28 (0.18–0.37) <sup>g</sup>	0.376	<0.001

CI, confidence interval;  $\beta$ , not standardized regression coefficient; *R*<sup>2</sup>, determination coefficient; OBP, office blood pressure; HBP, blood pressure measurement; BMI, body mass index; WC, waist circumference.

<sup>a</sup> Expressed by 100 g.

<sup>b</sup> Simple linear regression analysis.

<sup>c</sup> Multiple linear regression analysis.

<sup>d</sup> *n* = 776.

<sup>e</sup> *n* = 701.

<sup>f</sup> Model adjusted for prematurity, socioeconomic classification, sex, public or private schools, family history of hypertension, height, and age.

<sup>g</sup> Model adjusted for prematurity, socioeconomic classification, sex, public or private schools, family history of hypertension, BMI, WC, birth-height, and age.

investigations are needed to clarify if this finding is a tendency and the possible factors behind this growth.

A notable aspect of this study was that children with LBW became shorter in adolescence. This result is in accordance with a 5.9 cm reduction in height, at the end of adolescence, observed in LBW children in the county of Hordaland (Norway)<sup>4</sup> as well as with a higher occurrence of low height (10.3%) in a group of adults who were born small for gestational age (SGA) when compared with the control group.<sup>26</sup>

Most SGA children recover their height in the two initial years of life; however, it is estimated that between 10% and 15% will maintain a height deficit ( $-2$  z-scores) throughout life. For those children, therapy with recombinant human growth hormone was recently approved in the United States and in Europe.<sup>20</sup>

Nonetheless, it is not clear whether the relation of birth weight with height results from genetic mechanisms, uterine influence, lifestyle, social condition, or a combination of these factors. A low height programming mechanism is probably related to changes on thyroid hormone synthesis, through phenylalanine and tyrosine reduction. It would cause oxygen consumption reduction and growth retardation.<sup>7</sup> Another explanation would be a predisposition to early sexual and bone maturation, which would result in a linear growth deficit.<sup>1</sup> Despite not being the focus of this study, the finding of loci (HMGA2 and LCORL) that genetically link intrauterine growth with postnatal height is noteworthy.<sup>27</sup>

The most recent WHO<sup>28</sup> recommendation has modified the previous classification of mild low height to normal, probably due to the world decline in height deficit prevalence among young individuals in the population. However, the height increase tendency throughout generations demands reassessing the reference curves and revising the very definition of low height, in order not to underestimate nutritional disorders in the population, mainly in LBW newborns.<sup>26</sup>

Birth weight was directly associated to BMI and WC increase in adolescents; however, this relation was not sustained after an adjusted analysis. The variations of these anthropometric variables may be totally or partially explained by postnatal period factors, more than by fetal biology.<sup>8</sup> These results corroborated studies<sup>4,29,30</sup> that found no significant association of birth weight with low weight or obesity during and at the end of adolescence.

Controversies about this relation remain. A Chilean study concluded that approximately one-third of obesity cases in high school students could have been averted with early interventions in newborns with macrosomia.<sup>6</sup> A Brazilian cohort study with newborns from the most economically developed region of the country observed a 1.2 kg/m<sup>2</sup> (95% CI: 0.0–2.4) increase in BMI at the age of 18 years in subjects with birth weight  $\geq 4$  kg.<sup>31</sup>

Overweight was observed in more than one fifth of this population, exceeding by six times the frequency of underweight. These results confirm the nutritional transition observed in middle-income countries, characterized by progressive reduction of undernutrition and continuous increase in obesity prevalence. Among Brazilians aged 10–19, the

prevalence of underweight was reported in 3.4% while that of overweight/obesity was 20.5%.<sup>32</sup>

No association was observed between birth weight and OBP/HBP in adolescence, contradicting authors, who propose that LBW due to inadequate uterine development would be associated with higher risk of developing hypertension in youth.<sup>5,8</sup>

Nevertheless, the investigation on the impact of birth weight on BP is controversial. Another Brazilian study, which used ambulatory blood pressure (ABP) monitoring, revealed higher SBP at nighttime and lower dipper patterns of SBP in students with LBW; however, it did not find correlation with daytime SBP.<sup>9</sup>

A limitation of the present study was the lack of information about gestational age. Without that, it was not possible to establish whether the SGA condition was due to intrauterine growth restriction (IUGR). However, the term SGA presents limitations from an epidemiological perspective since it disregards factors such as biological variability, gender, multiplicity, ethnicity, and parity, inferring that all small babies result from a pathological IUGR. In this sense, there is no golden standard method to determine IUGR with accuracy, which hampers the accomplishment of studies on fetal programming. Moreover, birth weight is the only perinatal health indicator widely collected in Brazil, due to its practicality and strong association with infant mortality.<sup>33</sup>

Other limitations refer to the socioeconomic condition, which does not necessarily correspond to the context in which the child was born; the collection of birth variables, which was performed exclusively by parents or guardians of the adolescents; and the absence of data on sexual maturation, parents' height and smoking during gestation. Considering that 67.9% of the enrolled adolescents were included in this cross-sectional analysis, it is also important to consider the potential selection bias as a limitation.

Another factor which may have influenced the present results was the use of oscillometric method to measure BP rather than auscultatory, as well as the use of only two BP measurements. This methodology was chosen since the available recommendations for BP measurements in adolescents are based on experts' opinions, rather than on well-conducted experimental studies.<sup>34</sup>

Birth weight is a complex variable that is influenced by many factors of fetal and maternal genotype. Contemporary life course perspectives also acknowledge the complex interactions of biological factors with the environment. These are the challenges in furthering understanding the fetal origin of diseases given the complexity of environmental factors.<sup>33</sup>

In conclusion, LBW was a predictor of growth deficit in adolescence. There was no association between birth weight and OBP, HBP, WC, and BMI in this age group. The high prevalence of newborns with low and high birth weight in the studied population demonstrates the need for a better prenatal assistance and nutritional attention to the pregnant woman, in order to promote maternal health, prevent neonatal mortality, and ensure normal growth/development of the child.

## Funding

This study was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação de Amparo à Pesquisa do Estado de Goiás (FAPEG).

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgements

The authors would like to thank the adolescents, relatives, and guardians who participated in the study, as well as to thank the support offered by the Hypertension League from the Federal University of Goiás.

## References

- Barker DJ. The fetal and infant origins of adult disease. *BMJ*. 1990;301:1111.
- Lansky S, Friche AA, Silva AA, Campos D, Bittencourt SD, Carvalho ML, et al. Birth in Brazil survey: neonatal mortality, pregnancy and childbirth quality of care. *Cad Saúde Pública*. 2014;30:S192–207.
- Hack M, Klein NK, Taylor HG. Long-term developmental outcomes of low birth weight infants. *Future Child*. 1995; 5:176–96.
- Odberg MD, Sommerfelt K, Markestad T, Elgen IB. Growth and somatic health until adulthood of low birthweight children. *Arch Dis Child Fetal Neonatal Ed*. 2010;95:F201–5.
- Chen W, Srinivasan SR, Berenson GS. Amplification of the association between birthweight and blood pressure with age: the Bogalusa Heart Study. *J Hypertens*. 2010;28: 2046–52.
- Loaiza S, Atalah E. Birth weight and obesity risk at first grade of high school in a non-concurrent cohort of Chilean children. *Public Health Nutr*. 2012;16:228–32.
- Bagnoli F, Laura F, Sara N, Salvatore G. Thyroid function in small for gestational age newborns: a review. *J Clin Res Pediatr Endocrinol*. 2013;5:2–7.
- Sayers S, Singh G, Mott S, McDonnell J, Hoy W. Relationships between birthweight and biomarkers of chronic disease in childhood: Aboriginal Birth Cohort Study 1987–2001. *Paediatr Perinat Epidemiol*. 2009;23:548–56.
- Salgado CM, Jardim PC, Teles FB, Nunes MC. Low birth weight as a marker of changes in ambulatory blood pressure monitoring. *Arq Bras Cardiol*. 2009;92:113–21.
- Bassareo PP, Marras AR, Cugusi L, Zedda AM, Mercurio G. The reasons why cardiologists should consider prematurity at birth and intrauterine growth retardation among risk factors. *J Cardiovasc Med (Hagerstown)*. 2016;17:323–9.
- Nascente FM, Jardim TV, Peixoto MD, Carneiro CS, Mendonça KL, Póvoa TI, et al. Sedentary lifestyle and its associated factors among adolescents from public and private schools of a Brazilian state capital. *BMC Public Health*. 2016;16:1177.
- World Health Organization (WHO). Physical status: the use and interpretation of anthropometry. Technical report series. Geneva: WHO; 1995.
- World Health Organization (WHO). Measuring change in nutritional status. Guidelines for assessing the nutritional impact of supplementary feeding programme. Geneva: WHO; 1983.
- World Health Organization (WHO). Growth reference data for 5–19 years. Geneva: WHO; 2007.
- Taylor RW, Jones IE, Williams SM, Goulding A. Evaluation of waist circumference, waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual-energy X-ray absorptiometry, in children aged 3–19 y. *Am J Clin Nutr*. 2000;72:490–5.
- Stergiou GS, Yiannes NG, Rarra VC. Validation of the Omron 705 IT oscillometric device for home blood pressure measurement in children and adolescents: the Arsakion School Study. *Blood Press Monit*. 2006;11:229–34.
- National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics*. 2011;114:555–76.
- Sociedade Brasileira de Cardiologia (SBC), Sociedade Brasileira de Hipertensão (SBH), Sociedade Brasileira de Nefrologia (SBN). [V Guidelines for ambulatory blood pressure monitoring (ABPM) and III Guidelines for home blood pressure monitoring (HBPM)]. *Arq Bras Cardiol*. 2011;97:1–24.
- Stergiou GS, Yiannes NG, Rarra VC, Panagiotakos DB. Home blood pressure normalcy in children and adolescents: the Arsakeion School study. *J Hypertens*. 2007; 25:1375–9.
- Clayton PE, Cianfarani S, Czernichow P, Johannsson G, Rapaport R, Rogol A. Management of the child born small for gestational age through to adulthood: a consensus statement of the International Societies of Pediatric Endocrinology and the Growth Hormone Research Society. *J Clin Endocrinol Metab*. 2007;92:804–10.
- Viana K de J, Taddei JA, Cocetti M, Warkentin S. Birth weight in Brazilian children under two years of age. *Cad Saude Publica*. 2013;29:349–56.
- Giglio MR, Lamounier JA, Morais Neto OL, César CC. Low birth weight in a cohort of newborns in Goiânia-Brazil in 2000. *Rev Bras Ginecol Obstet*. 2005;27:130–6.
- Silva AA, Silva LM, Barbieri MA, Bettiol H, Carvalho LM, Ribeiro VS, et al. The epidemiologic paradox of low birth weight in Brazil. *Rev Saúde Pública*. 2010;44:767–75.
- Costa BM, Paulinelli RR, Barbosa MA. Association between maternal and fetal weight gain: cohort study. *Sao Paulo Med J*. 2012;130:242–7.
- Bamberg C, Hinkson L, Henrich W. Prenatal detection and consequences of fetal macrosomia. *Fetal Diagn Ther*. 2013;33: 143–8.
- Jaquet D, Collin D, Lévy-Marchal C, Czernichow P. Adult height distribution in subjects born small for gestational age. *Horm Res*. 2004;62:92–6.
- Horikoshi M, Yaghoobkar H, Mook-Kanamori DO, Sovio U, Taal HR, Hennig BJ, et al. New loci associated with birth weight identify genetic links between intrauterine growth and adult height and metabolism. *Nat Genet*. 2013;45:76–82.
- World Health Organization (WHO). Growth reference data for 5–19 years; 2007. Available from: <http://www.who.int/growthref/en/> [cited June 2016].
- Fåhræus C, Wendt L-K, Nilsson M, Isaksson H, Alm A, Andersson-Gäre B. Overweight and obesity in twenty-year-old Swedes in relation to birthweight and weight development during childhood. *Acta Paediatr*. 2012;101:637–42.
- Wang Y, Gao E, Wu J, Zhou J, Yang Q, Walker MC, et al. Fetal macrosomia and adolescence obesity: results from a longitudinal cohort study. *Int J Obes (Lond)*. 2009; 33:923–8.
- Goldani MZ, Haeffner LS, Agranonik M, Barbieri MA, Bettiol H, Silva AA. Do early life factors influence body mass index in adolescents? *Braz J Med Biol Res*. 2007;40: 1231–6.
- Brazilian Institute of Geography and Statistics (IBGE). National Survey of Households 2008. Rio de Janeiro: IBGE;

- 2008, v. 29. Available from: <http://biblioteca.ibge.gov.br/visualizacao/livros/liv42672.pdf> [cited June 2016].
33. Urquia ML, Ray JG. Seven caveats on the use of low birth-weight and related indicators in health research. *J Epidemiol Community Health*. 2012;66:971–5.
  34. Lurbe E, Agabiti-Rosei E, Cruickshank JK, Dominiczak A, Erdine S, Hirth A, et al. 2016 European Society of Hypertension guidelines for the management of high blood pressure in children and adolescents. *J Hypertens*. 2016;34:1887–920.