

## RESEARCH NOTE

## Enterotoxigenic Capacity of *Campylobacter coli* Strains Isolated from Humans and Bovines

A Tresierra-Ayala/<sup>+</sup>, H Fernández\*, J Zamora\*\*

Departamento de Microbiología, Universidad Nacional de la Amazonía Peruana, Casilla Postal 751, Iquitos, Perú \*Instituto de Microbiología Clínica, Facultad de Medicina \*\*Instituto de Microbiología, Facultad de Ciencias, Universidad Austral de Chile, Casilla Postal 567, Valdivia, Chile

Key words: *Campylobacter coli* - ileal loop test - enterotoxigenic capacity - pathogenicity - reservoirs

In developing countries, *Campylobacter coli* and *C. jejuni* are important agents associated with diarrhetic syndrome. In Latin America, their isolation rates vary from 0.6% in Cuba to 23.0% in Iquitos, in the Peruvian jungle region (H Fernández 1992 *J Braz Ass Develop Sci* 44: 39-43). However, most of the studies have been done only with *C. jejuni*, in spite of that *C. coli* has similar importance (T Daikoku et al. 1989 *FEMS Microbiol Lett* 58: 33-36).

Enterotoxigenic capacity has been identified as a pathogenic mechanism in *Campylobacter* (G Ruiz-Palacios et al. 1983 *Lancet* 2: 250-251, H Fernández & L Trabulsi 1995 *Biol Res* 28: 205-210). Some researchers have reported that this property is similar in both species (W Johnson & H Lior 1986 *J Clin Microbiol* 24: 275-281), however others have determined that the enterotoxigenic capacity of *C. coli* is lower than *C. jejuni* or this bacterium has not enterotoxic property (T Daikoku et al. 1989 *FEMS Microbiol Lett* 58: 33-36, C Fricker & R Park 1989 *J Appl Bacteriol* 66: 477-490).

This work received financial support from Grants Fondecyt 59-89; Fondecyt 217-90 and General Office of Investigation - UNAP.

\*Corresponding author. Fax: 51-94-23.4723

Received 5 February 1996

Accepted 29 March 1996

The aim of this study was to evaluate the presence of enterotoxigenic capacity in 15 strains of *C. coli* isolated from humans (7 strains) and from bovines (8 strains), previously maintained under freezing temperature (-35°C). Using the Lior biotyping scheme (H Lior 1984 *J Clin Microbiol* 20: 636-640), 3 belonged to biovar I and 12 to biovar II (Table).

TABLE

Distribution of biovars of *Campylobacter coli* strains studied according to the isolation source and determination of their enterotoxigenic capacity

Source	Biovars		Total TOX/TOT
	I	II	
	TOX/TOT	TOX/TOT	
Human	1/2	3/4	4/7 (57.1)
Bovine	1/1	5/7	6/8 (75.0)
Total	2/3 (66.7)	8/12 (66.7)	10/15 (66.7)

TOX/TOT: no. of enterotoxigenic strains/no. of strains studied; ( ) : % of strains with enterotoxigenic capacity. The difference between the enterotoxigenic capacity of human and bovine strains was not significant ( $p > 0.05$ ).

The enterotoxigenic capacity was determined by the ligated ileal loop test in rats (S Saha et al. 1988 *J Med Microbiol* 26: 87-91). A cell-free supernatant of each strain was prepared from a 72 hr culture in *Brucella* broth incubated at 37°C in microaerophilic conditions and 500 µl were inoculated individually into single 5-cm-long ligated ileal loops of adult Holtzman rats (170 to 250 g). Sterile *Brucella* broth and the cell-free supernatant of a known enterotoxigenic *C. coli* strain were used as negative and positive control, respectively. After 18 hr, the animals were killed by ether overdose and the ileum was examined. Distention of the loops with fluid accumulation was considered as a positive result (98 µl/cm of intestine or more, as determined by Student's t test). The volume/length ratio in negative control was consistently <20 µl/cm. The tests were done in two rats simultaneously.

The enterotoxigenic capacity was determined in 66.7% of the strains, belonging two strains to the biovar I and eight to the biovar II. Some investigators have reported rates higher than our values. H Bok et al. (1991 *J Clin Microbiol* 29: 1262-1264) and W Johnson and H Lior (1986 *J Clin Microbiol* 24: 275-281) determined rates of 100% and 83.2%, respectively.

Moreover, our results showed that 57.1% of the culture supernatants of human strains and 75% of the supernatants of bovine strains produced fluid

accumulation. They are in agreement with the results reported in England by C Fricker and R Park (1989 *J Appl Bacteriol* 66: 477-490) because in both studies, the difference between the human and bovine strains, in relation to their enterotoxigenic capacity, was not significant ( $p > 0.05$ ). Consequently, this characteristic was not associated with the biovar neither with the strains source.

The presence of enterotoxigenic strains in bovines increases the evidences that these animals could be an important infection source for humans. Some authors have expressed that in developing countries, the close association between man and

animals may lead to a cycle of infection that would increase the toxigenicity of *Campylobacter* strains by consecutive passages through the intestinal tract (S Saha et al. 1988 *J Med Microbiol* 26: 87-91). A similar enhancement of other pathogenic factors of *Campylobacter* has been suggested by D Newell et al. (1985 *J Hyg* 94: 45-54).

The present study suggests that the enterotoxigenicity could be an important pathogenic mechanism by which *C. coli* may produce disease. For this reason we consider that this bacterium must have the same importance as *C. jejuni* in relation to their enterotoxigenicity.