

Nutrient omission effect on growth and nutritional status of assai palm seedlings¹

Fábio Reis Ribeiro Araújo², Ismael de Jesus Matos Viégas³,
Raimundo Lázaro Moraes da Cunha⁴, Werica Larissa Farias de Vasconcelos²

ABSTRACT

The Amazonian assai palm has a great socioeconomic importance, but most of its commercial plantations take place in uplands and low natural fertility soils, what may hinder its development. This study aimed at evaluating the effect of nutrient omission on growth and nutritional status of assai palm seedlings (Ver-o-Peso cultivar). The experimental design was completely randomized, with 14 treatments and 5 replicates. The treatments consisted of complete fertilization with liming; no fertilization and no liming (control); complete fertilization with individual omission of N, P, K, Ca, Ca with no liming, Mg, Mg with no liming, S, B, Cu and Zn. Plant height, stem diameter and dry mass, leaf and total shoot dry mass, and leaf nutrients content and accumulation were evaluated. The initial growth of the assai palm plants was limited by the omission of P, N, K, Ca, Mg and Cu. The production of leaf dry mass was decreasingly affected by the omission of P > Cu > N > K > Mg, while leaf area was limited by the individual omissions of Ca > N > P > K > Mg > Zn. Plant development, measured by relative growth of shoots, was affected by lack of Ca > P > N > Mg > Cu > K, with an average reduction of 31 %. The nutrients most needed by the assai palm plants, as evidenced by nutrients contents and accumulation in the leaf dry mass, are: N > K > S > Ca > Mg > P > Mn > Zn > B > Cu.

KEYWORDS: *Euterpe oleracea* Mart.; palm tree; plant nutrition.

RESUMO

Efeito da omissão de nutrientes no crescimento e estado nutricional de mudas de açaizeiro

O açaizeiro é uma palmeira amazônica de grande importância socioeconômica, mas a maioria de seus plantios comerciais ocorre em terras firmes e de baixa fertilidade natural, o que pode causar diminuição em seu desenvolvimento. Objetivou-se avaliar o efeito da omissão de nutrientes no crescimento e estado nutricional de mudas de açaizeiro (cultivar Ver-o-Peso). O delineamento experimental foi inteiramente casualizado, com 14 tratamentos e 5 repetições. Os tratamentos consistiram de adubação completa com calagem; sem adubação e sem calagem (testemunha); adubação completa com omissões individuais de N, P, K, Ca, Ca sem calagem, Mg, Mg sem calagem, S, B, Cu e Zn. Foram avaliados a altura das plantas, diâmetro e massa seca do caule, massa seca das folhas e aérea total, teores e acúmulo de nutrientes nas folhas. O crescimento inicial dos açaizeiros foi limitado pela omissão de P, N, K, Ca, Mg e Cu. A produção de massa seca das folhas foi afetada em ordem decrescente pela omissão de P > Cu > N > K > Mg, enquanto a área foliar foi restringida por omissões individuais de Ca > N > P > K > Mg > Zn. O desenvolvimento das plantas, medido pelo crescimento relativo da parte aérea, foi afetado pela carência de Ca > P > N > Mg > Cu > K, com redução média de 31 %. Os nutrientes mais exigidos pelos açaizeiros, evidenciados pelos teores e acúmulos de nutrientes na massa seca das folhas, são: N > K > S > Ca > Mg > P > Mn > Zn > B > Cu.

PALAVRAS-CHAVE: *Euterpe oleracea* Mart.; palmeira; nutrição vegetal.

INTRODUCTION

Assai palm (*Euterpe oleracea* Mart.) is native from Amazonia, being abundant in the estuary of the Amazon River, in lowland and flooded forest areas. The main product of the assai palm is its fruit, which is basically consumed *in natura* in juices. It is also an economic alternative for the production of heart of palm, because of its multi-stemmed clump growth

habit, making its continued exploitation viable, when managed rationally (Menezes et al. 2008).

In 2015, the national production of assai fruit, in Brazil, totaled more than one million tons, with the main producer being the Pará State (91.6 %), which reached 154,000 ha (IBGE 2015). In 2016, Embrapa Amazônia Oriental will launch the assai palm BRS Ver-o-Peso cultivar, which is destined for planting in irrigated upland, and will be able to produce off

1. Manuscript received in Apr./2016 and accepted for publication in Oct./2016 (<http://dx.doi.org/10.1590/1983-40632016v46a40770>).

2. Universidade Federal do Sul e Sudeste do Pará, Instituto de Estudos em Desenvolvimento Agrário e Regional, Faculdade de Ciências Agrárias, Marabá, PA, Brazil. *E-mails*: fabioaraujo@unifesspa.edu.br, werica_lee@hotmail.com.

3. Universidade Federal Rural da Amazônia, Campus Capanema, Capanema, PA, Brazil. *E-mail*: ismael.viegas@ufra.edu.br.

4. Universidade Federal Rural da Amazônia, Campus Belém, Belém, PA, Brazil. *E-mail*: cunhalazaro@yahoo.com.br.

season fruits with a higher yield than the BRS Pará cultivar.

Assai palm naturally occurs in lowlands, which are ecosystems with eutrophic, acidic, clay-silty soils and a good natural fertility, due to depositions of sediments brought in by tides (Oliveira et al. 2007). These soils receive a regular contribution of various sediments rich in N, Ca, P, K and Mg that are transported by the Amazon River and its tributaries from the Andean mountains (Almeida et al. 2004). However, the majority of assai palm commercial plantations are located in upland soils, where Ferralsols of low natural fertility dominate and the provision of nutrients through fertilization is necessary to achieve high yields (Viégas et al. 2004).

Palms require high levels of nutrients for both vegetative growth and reproduction. In addition, an adequate supply of fertilizer is necessary to promote a greater initial growth (Bovi et al. 2002). Experiments with fertility may help to determine which nutrients are most limiting to plant development (Lopes 2001, Viégas et al. 2004). The first research on assai palm nutrition emphasized that macronutrients interfere with the production of total dry matter by young plants in the following order: $K > Mg > P > N > Ca > S$ (Haag et al. 1992, Oliveira et al. 2002).

Nutritional disorders cause a decrease in the development and production of any crop, and commonly manifested visual symptoms tend to be more pronounced in the leaves. However, yield may often be compromised even before symptoms occur (Malavolta 1980). In this context, the missing element technique evaluates the nutritional requirements of crops and is an effective tool for acquiring qualitative information on the nutrients that most limit plant growth (Laviola & Dias 2008, Miranda et al. 2010).

Thus, the present study aimed at evaluating the growth, dry mass production and content and accumulation of macro and micronutrients in assai palm seedlings (BRS Ver-o-Peso cultivar), using a sandy-textured Ferralsol substrate under the effect of individual nutrients omission.

MATERIAL AND METHODS

The experiment was conducted under greenhouse conditions at the Universidade Federal Rural da Amazônia, in Belém, Pará State, Brazil, from September 2013 to April 2014. The substrate used was a sandy-textured Ferralsol (FAO 2015),

“Latossolo Amarelo” in the Brazilian system (Embrapa 2013), collected in Belém, at a depth of 0-30 cm, which has low natural fertility and is deep and well drained.

Prior to the experiment, a physico-chemical analysis of the soil was performed (Embrapa 1997), resulting in the following profile: clay = 11 %; silt = 6 %; sand = 83 %; pH (H₂O) = 4.8; P = 25.28 mg dm⁻³; K = 20.3 mg dm⁻³; Ca = 0.70 cmol_c dm⁻³; Mg = 0.10 cmol_c dm⁻³; S = 6.8 mg dm⁻³; Al = 0.70 cmol_c dm⁻³; H + Al = 6.30 cmol_c dm⁻³; B = 0.33 mg dm⁻³; Cu = 0.89 mg dm⁻³; Mn = 3.6 mg dm⁻³; Zn = 1.41 mg dm⁻³; V (%) = 11.9; OM = 23.6 g kg⁻¹; SB = 0.85 cmol_c dm⁻³. The high levels of P indicate that the soil had been previously fertilized.

The pH (H₂O) was determined using a glass electrode placed in a 1:2.5 soil to liquid suspension. The amounts of Ca and Mg were measured by atomic absorption, after their extraction with KCl at 1N; P was extracted with a solution of H₂SO₄ + HCl 0.025N and measured with a spectrophotometer (wavelength of 660 nm); K was determined by flame photometry and Al by titration with NaOH 0.025N.

The experimental design was completely randomized, with 14 treatments and 5 replicates, with each container corresponding to one experimental unit. The treatments were: complete (complete fertilization with all the nutrients plus liming); control (without fertilization and without liming); complete fertilization with liming and omission of nitrogen (-N), phosphorus (-P), potassium (-K), calcium (-Ca), calcium without liming (-Ca/-lime), magnesium (-Mg), magnesium without liming (-Mg/-lime), sulfur (-S), boron (-B), copper (-Cu), manganese (-Mn) and zinc (-Zn).

Liming was performed using the base saturation method, aiming to raise the base saturation up to 60 % (Cravo et al. 2010), applying dolomitic limestone with 32 % of CaO, 14 % of MgO, 67 % of NP and relative power of total neutralization of 95 %. The lime was homogeneously mixed with the substrate and incubated for 30 days.

Seedlings were produced using seeds of the new assai palm BRS Ver-o-Peso cultivar, developed by the Embrapa Amazônia Oriental. The seeds were sown in seedbeds containing a mixture of black earth, sawdust and aged cattle manure in the proportion of 3:1:1. When seedlings reached an average height of 10 cm and showed one pair of leaves, they were transplanted in pairs into plastic containers with 5 kg

of air-dried fine earth that had been previously sieved (4 mm mesh).

At 45 days after planting, the seedlings were thinned for homogenization and all containers were provided with mineral fertilization, according to their designated treatments. Fertilization was performed according to Viégas et al. (2009), with the following doses, taking into account the missing element technique: N = 100 mg kg⁻¹ [-CO(NH₂)₂]; P = 50 mg kg⁻¹ (-NaH₂PO₄); K = 90 mg kg⁻¹ (-KCl); Ca = 30 mg kg⁻¹ (-CaCl₂); Mg = 30 mg kg⁻¹ (-MgCl₂); S = 7.5 mg kg⁻¹ (-NaSO₄); B = 1.2 mg kg⁻¹ (-H₃BO₃); Cu = 1.0 mg kg⁻¹ (-CuSO₄); Mn = 4 mg kg⁻¹ (-MnSO₄); and Zn = 5 mg kg⁻¹ (-ZnSO₄) of soil.

A fertilizer containing N and K was applied at 45, 90 and 150 days after planting. The soil was irrigated with distilled water, while weighing the pots to maintain substrates at 80 % of field capacity. Plant height, stem diameter at 2.5 cm from the substrate and leaf number and size (length and width of leaflets) were measured.

Plants were collected at eight months after planting the seedlings, separating the leaflets, petioles, stems and roots. The separate organs were packed in paper bags, identified and placed in a forced-circulation oven at 70 °C, until a constant mass was reached, at which point the dried material was weighed and ground in a Wiley mill, in order to determine the nutrient content. The shoot relative growth (SRG) was calculated as it follows: SRG (%) = (DMNO/DMCT) x 100, where DMNO = total shoot dry mass for each nutrient omission and DMCT = total shoot dry mass for the complete treatment. Leaf area (LA) was calculated following an adaptation of the method by Clement & Bovi (2000), as it follows: LA (m²) = (L x W) x 0.535, where L = leaf length, W = leaf width and 0.535 = constant adjusting the shape of a rectangle to the shape of a leaf. The determination of leaf macro and micronutrients was performed according to Embrapa (2010).

An analysis of variance (Anova) was performed on the data for the following variables: plant height, stem diameter, leaf and stem dry mass, and total content and accumulation of macro and micronutrients. To determine the accumulation of nutrients, the content (g kg⁻¹) of each nutrient was multiplied by the leaf dry mass.

For variables significantly affected by the treatments, the Dunnett's test was used to compare the treatments means with that of the complete

treatment, at 5 %. Analyses were performed using the Action[®] software.

RESULTS AND DISCUSSION

The results for plant height, stem height and diameter, shoot dry biomass, leaf area and relative shoot growth, according to the treatments, are shown in Table 1.

Nutrient omission treatments exhibited values for plant height and stem height and diameter less than, or equal to, that of the complete treatment. The absence of liming and fertilization in the control plants restricted plant height by 41.5 %, stem height by 45.2 % and stem diameter by 36.5 %, in relation to the complete treatment. These results justify the importance of nutrients fertilization for the proper development of assai palm plants cultivated in sandy-textured Ferralsol. Stem height and diameter for the control plants were the only estimates that differed significantly from the complete treatment.

The omissions of the macronutrients N, P, K, Ca/-lime, Mg and Mg/-lime were the most restricting to the initial growth of assai palm plants. The omission of these nutrients was responsible for an average decrease of 22.1 % in plant height, relatively to the complete treatment, with N, P and K decreasing the plant height by 20.3 %, 23.3 % and 20.3 %, respectively. These results corroborate Viégas et al. (2004), who found that the omission of N, P and K in medium-textured Ferralsol limited the plant height of assai palm (BRS Pará cultivar) by 35.4 %, 38.5 % and 26 %, respectively. Viégas et al. (2009) also showed that the individual omission of N, P and K in Ferralsol limits the plant height of assai palm.

The omission of Ca/-lime restricted plant height by 24.4 %, showing the importance of the joint practices of fertilization and liming. However, the effect of the absence of liming was not found with the individual omission of Mg and Mg/-lime, which restricted plant height by 23.1 % and 21.1 %, respectively. Viégas et al. (2008) indicated that the omission of Mg restricted the growth of assai palm seedlings by 50.8 %.

Regarding the micronutrients omission, only the omission of Cu showed a significant difference from the complete treatment, with a reduction of 20.3 % in plant height. Viégas et al. (2004) recorded a reduction of 7.7 % in plant height of assai palm with the omission of Cu.

For shoot dry biomass, all the treatments were less than, or equal to, the complete treatment (Table 1). The individual nutrients whose omission had the greatest restriction to leaf dry mass production, relatively to the complete treatment, were P with a reduction of 34.3 %; N with 32.5 %; K with 30.7 %; and Mg with 28.1 %. Regarding the omission of micronutrients, the only treatment with a significant difference was Cu, with a reduction of 33.7 % in leaf dry mass, if compared to the complete treatment. The control treatment restricted the leaf dry mass gain by 66.1 %.

For the stem dry mass production, the individual nutrients that were most restrictive when omitted were N, P, Ca/-lime and Mg/-lime, with reductions relatively to the complete treatment of 35.2 %, 36.3 %, 39.8 % and 34.1 %, respectively. The other individual omissions did not exhibit significant differences, and the control experienced a decrease of 77.3 % in stem dry mass.

For the shoot dry mass production, the omitted elements that limited mass gain, relatively to the complete treatment, were the same as those for stem dry mass: N, P, Ca/-lime and Mg/-lime, with reductions of 33.3 %, 34.5 %, 37 % and 24.8 %, respectively. The absence of fertilization and liming in the control treatment decreased the gain of shoot dry mass by 71.7 %.

Viégas et al. (2004) observed a decrease in leaf dry mass and stem and shoot dry mass, in assai palm plants, as an effect of the individual omission of N, P, K, Ca and Mg, relatively to the complete treatment. For leaves, the mass of the complete treatment was 51.4 g, while the omissions of N, P, K and Mg reduced growth by 49.3 %, 49 %, 26.8 % and 21.7 %, respectively. For stems, the omission of N reduced growth by 70.7 %, P by 74 %, K by 40.9 %, Ca by 39.6 % and Mg by 24.6 %. For shoot dry mass, the omissions were responsible for a decrease of 58.98 % in the absence of N, 60.21 % for P, 33.15 % for K, 27.21 % for Ca and 37.14 % for Mg.

Leaf area was restricted by the individual omission of the nutrients N, P, K and Mg by 40.6 %, 39.2 %, 35.9 % and 37.6 %, respectively. The omission of Zn decreased leaf area by 35.8 %, relatively to the complete treatment. The control treatment had its leaf area reduced by 69 %. Viégas et al. (2008) showed that the number of leaves on assai palm plants, in treatments with individual omission of N, P and K, were reduced significantly (around 50 %, 33 % and 30 %, respectively), in relation to the complete treatment.

The shoot relative growth followed the descending order of complete > Ca > S > B > Zn > Mg > Mn > K > Cu > Mg/-lime > N > P > Ca/-lime > control. Therefore, the development of the plant,

Table 1. Height (HT), stem height (SH), stem diameter (SD), leaf dry mass (LDM), stem dry mass (SDM), shoot dry mass (SHDM), leaf area (LA) and relative growth (RG) of assai palm, according to the treatment. Treatments: complete (complete fertilization with all nutrients and liming); control (without fertilization and without liming); complete fertilization with liming and omission of a nutrient represented by its chemical symbol (-X); complete fertilization without liming and with nutrient omission represented by its chemical symbol (-X/-lime).

Treatment	HT	SH	SD	LDM	SDM	SHDM	LA	RG
	cm	cm	mm	g	g	g	cm ²	%
Complete	72.00	22.00	16.85	9.53	33.62	43.15	1,776.72	100.0
-N	57.40*	18.40	15.38	6.43*	21.79*	28.76*	1,055.64*	65.4
-P	55.20*	18.00	14.84	6.26*	21.40*	28.25*	1,079.61*	64.1
-K	57.40*	20.20	16.36	6.60*	25.94	34.34	1,139.37*	75.4
-Ca	60.20	21.80	16.34	9.48	33.36	42.84	1,543.03	99.3
-Ca/-lime	54.40*	18.60	14.20	6.91	20.25*	27.17*	948.31*	63.0
-Mg	55.40*	18.80	15.27	6.85*	29.27	39.12	1,108.63*	83.7
-Mg/-lime	56.80*	19.00	16.19	8.37	22.15*	32.46*	1,322.22	70.7
-S	67.00	22.00	16.39	8.91	33.20	42.11	1,204.72	97.6
-Cu	57.40*	20.40	14.42	6.32*	26.11	41.79	1,254.77	75.2
-Zn	58.60	20.00	16.22	8.58	30.94	32.74	1,140.61*	91.6
-Mn	60.40	20.80	16.18	7.64	27.48	35.12	1,196.98	81.4
-B	62.80	20.60	16.20	9.00	31.99	39.52	1,239.26	95.0
Control	42.15*	12.05*	10.70*	3.23*	8.97*	12.20*	551.26*	28.3
CV (%)	11.50	13.50	9.30	19.63	22.97	21.50	28.82	25.1

* Average significantly different from the complete treatment, in the same column, according to the Dunnett's test ($p < 0.05$).

during the experimental period, was most affected by the total omission of nutrients in the control, with a reduction of 71.7 %. Phosphorus affected shoot relative growth the most, decreasing it by 35.9 %, followed by N with a 34.6 % reduction, Mg/-lime with 29.3 %, Cu with 24.8 % and K with 24.6 %.

The limitations to the development of assai palm seedlings, in the present study, may be explained by the low level of nutrients present in the substrate. Although the soil used displayed a satisfactory amount of P, it had a low base saturation index ($V = 11.92\%$) and unsatisfactory amounts of K (20.28 mg dm^{-3}), Ca ($0.7 \text{ cmol}_c \text{ dm}^{-3}$) and Mg ($0.1 \text{ cmol}_c \text{ dm}^{-3}$). As observed by Cravo et al. (2010), the average concentrations of these elements for this crop in the Pará State are $41\text{-}60 \text{ mg dm}^{-3}$ for K, $0.5\text{-}1.5 \text{ cmol}_c \text{ dm}^{-3}$ for Ca and $0.5\text{-}1.5 \text{ cmol}_c \text{ dm}^{-3}$ for Mg. The low levels of these elements may be considered limiting factors, in accordance with the Liebig's law (Malavolta 2006), as these low levels may restrict the absorption of other nutrients.

Thus, restrictions to growth and dry biomass gain caused by the omission of N, P, K, Ca and Mg, in this study, may be explained by the metabolic functions that result in plant biomass growth. Nitrogen is a constituent of the chlorophyll molecule, as well as various cell components, including proteins, amino acids and nucleic acids, and its deficiency

alters the activity of enzymes from the biochemical phase of photosynthesis and from carbohydrate and nitrogen metabolism, thereby decreasing the rate of photosynthesis and thus the plant growth (Taiz & Zeiger 2013). Phosphorus is a structural element of nucleotides and phospholipids, and it acts on the transfer of energy through phosphorus compounds (Cunha et al. 2009). Therefore, P deficiency leads to lower growth by reducing the production and transport of energy.

The absence of Mg affects the size, structure and function of chloroplasts, since this element is a constituent of the chlorophyll molecule, in addition to activating enzymes involved in the processes of respiration, photosynthesis and nucleic acid synthesis (Taiz & Zeiger 2013). Potassium acts in maintaining a favorable hydric system, among other functions, by osmotic stomatal regulation, influencing the transport and storage of carbohydrates, synthesis of proteins, enzyme activity and leaf starch synthesis (Armenguard et al. 2009).

The nutrient contents on the leaf dry mass of assai palms, for each treatment, are shown in Table 2.

In general, the treatments with omission of an individual nutrient caused a reduction in the content of that nutrient in the plant. In the leaf dry mass of the complete treatment, the descending sequence of contents for the different macronutrients was $N > K >$

Table 2. Levels of macro and micronutrients in the leaf dry matter of assai palm cultivated in sandy-textured Ferralsol, according to the treatment. Treatments: complete (fertilization with all nutrients and liming); control (without fertilization and without liming); complete fertilization with liming and omission of a nutrient represented by its chemical symbol (-X); complete fertilization without liming and with nutrient omission represented by its chemical symbol (-X/-lime).

Treatment	Content									
	N	P	K	Ca	Mg	S	Cu	Zn	Mn	B
	g kg ⁻¹					mg kg ⁻¹				
Complete	24.14	1.25	14.52	2.61	2.19	7.39	7.22	93.17	99.93	54.45
-N	12.88*	1.02*	9.50*	7.20*	3.04*	5.47*	4.08*	66.22*	76.66*	58.77
-P	22.69	0.91*	9.62*	5.50	2.45	6.52	6.00*	60.15*	84.78*	60.21
-K	24.20	1.24	4.77*	6.44*	3.10*	6.47	5.14*	64.41*	83.75*	66.98*
-Ca	24.27	1.19	11.67*	6.52*	2.98	4.50	4.43*	66.64*	71.33*	55.30
-Ca/-lime	22.40	1.10	13.17	3.09	2.00	6.34	8.93*	100.34*	116.48*	50.54
-Mg	21.98	1.25	9.86	5.49*	1.80	5.79*	6.21	71.51*	90.46	52.85
-Mg/-lime	23.03	1.23	18.11	2.58	0.81*	5.13*	6.65	80.23*	109.62	55.71
-S	21.21*	1.19	9.50	7.34*	3.04*	6.48	5.23*	62.71*	84.13*	54.95
-Cu	22.78	1.18	13.05	7.05*	3.16*	5.65*	5.09*	59.17*	82.80*	54.09
-Zn	18.74*	1.11	8.98*	6.54*	2.65	3.02*	5.17*	34.11*	89.31	54.88
-Mn	23.31	1.21	11.55	7.18*	2.88	5.68*	6.23	64.40*	34.66*	46.51
-B	22.84	1.24	13.29	6.68*	2.86	6.85	5.99*	53.65*	95.28	17.47*
Control	15.72*	0.80*	4.48*	7.06*	2.29*	4.77*	4.25*	31.03*	34.26*	32.21*
CV (%)	10.21	11.60	25.74	19.40	15.14	10.22	7.73	9.09	10.47	23.30

* Average significantly different from the complete treatment, in the same column, by the Dunnett's test ($p < 0.05$).

S > Ca > Mg > P, while for the micronutrients it was Mn > Zn > B > Cu (Table 3). Fernandes et al. (2013) concluded that, for the peach palm (*Bactris gasipaes* Kunth) grown in a complete solution, the descending sequence of macronutrient concentration in the leaves and stems was N > K > Ca > P > Mg > S.

The content of N in the leaf dry mass of the complete treatment was 24.14 g kg⁻¹, a value higher than those found in similar experiments with assai palm in a Ferralsol: 15.67 g kg⁻¹ (Viégas et al. 2009); 19.39 g kg⁻¹ (Gonçalves 2004); and 16.6 g kg⁻¹ (Haag et al. 1992).

The absorption of N was limited by the omission of S and Zn (16.3 % and 22.9 % lower than the full treatments, respectively), whereas lower values of absorption were obtained for the treatments omitting N, with a reduction of 47.6 % in N levels, and the control with 34.9 % less N. The decrease in the level of N, as a result of the omission of S, is related to the decrease in the sugar content, accumulation of carbohydrates and protein synthesis (Hawkesford 2000), because of the low conversion of N to an organic form (Prado et al. 2007).

The content of P obtained in the present study (1.25 g kg⁻¹), in the complete treatment, is consistent with those observed by Haag et al. (1992) (1.3 g kg⁻¹) and Gonçalves (2004) (1.69 g kg⁻¹). Phosphorus was limited by the omission of N, P and Ca/-lime

(18.08 %, 27.2 % and 11.36 % lower than the complete treatment, respectively). The omission of all nutrients reduced the content of P by 35.68 %, since the absorption of P in low proportion was favored by the omission of Mg.

The content of K (14.52 g kg⁻¹) obtained in the complete treatment is four times higher than that obtained by Viégas et al. (2009) (3.52 g kg⁻¹), but comparable with the results by Haag et al. (1992) (19.6 g kg⁻¹) and Gonçalves (2004) (9.42 g kg⁻¹), with both using a nutrient solution with a silica substrate.

The K content was limited by the omissions of control > K > N > Zn > P > Mg. The level of K in the control was 69.2 % lower, if compared to the complete treatment. The omission of K from the soil favored the absorption of Ca by the assai palms, what may be related to the competitive inhibition between both nutrients in the absorption process (Malavolta et al. 1997).

The content of Ca obtained in the complete treatment was statistically greater than all the omissions by an average of 160 %. It is believed that the high absorption of Ca may have been enhanced by the supplementation of this element via soil liming with dolomitic limestone, in the majority of treatments.

The leaf content of Mg obtained in the complete treatment (2.19 g kg⁻¹) is comparable to

Table 3. Accumulation of macro and micronutrients in leaf dry material of assai palm cultivated in sandy-textured Ferralsol, according to the treatments. Treatments: complete (fertilization with all nutrients and liming); control (without fertilization and without liming); complete fertilization with liming and the omission of a nutrient represented by its chemical symbol (-X); complete fertilization without liming and with nutrient omission represented by its chemical symbol (-X/-lime).

Treatment	Accumulation (mg plant ⁻¹)									
	N	P	K	Ca	Mg	S	Cu	Zn	Mn	B
Complete	229.51	11.85	138.71	24.79	20.76	70.74	0.068	0.884	0.973	0.517
-N	92.09*	7.20*	65.65*	49.77	21.19	37.91*	0.029*	0.460*	0.533*	0.409
-P	156.46*	6.08*	65.74*	37.17	16.63	44.53*	0.042*	0.411*	0.579*	0.413
-K	202.21	10.39	41.09*	54.81*	26.24	54.33	0.043*	0.540*	0.704	0.564
-Ca	234.70	11.59	108.33	62.75*	28.86	43.49*	0.043*	0.638*	0.689	0.537
-Ca/-lime	155.81*	7.65*	87.75*	20.35	13.55	43.88*	0.060	0.691	0.780	0.351
-Mg	215.14	12.23	96.60	56.02*	17.39	57.29	0.061	0.709	0.895	0.522
-Mg/-lime	192.62	10.29	152.85	21.67	6.81*	42.90*	0.056	0.675	0.919	0.466
-S	190.10	10.69	84.12*	65.51*	27.38	57.74	0.046*	0.560*	0.746	0.489
-Cu	150.14*	7.81*	89.33	46.03	20.87	37.44*	0.034*	0.392*	0.548*	0.360
-Zn	159.73	9.52	75.86*	55.55*	22.44	25.87*	0.044*	0.291*	0.764	0.471
-Mn	176.69	9.15	90.21	54.65*	21.91	43.10*	0.048*	0.496*	0.265*	0.355
-B	223.44	12.17	130.16	65.38*	27.99	67.01	0.058	0.527*	0.933	0.172*
Control	51.36*	2.52*	15.24*	23.57*	7.55*	17.38*	0.014*	0.100*	0.109*	0.111*
CV (%)	22.79	23.70	32.41	30.17	27.05	23.90	23.41	23.57	23.70	24.72

* Averages significantly different from the complete treatment, in the same column, by the Dunnett's test ($p < 0.05$).

that by Viégas et al. (2009), who obtained a Mg leaf content of 2.2 g kg⁻¹ in the complete treatment, and close to that obtained by Haag et al. (1992) (3.5 g kg⁻¹). Also, the liming process may have caused the levels of Mg to average 40 % greater than the complete treatment, in the treatments with individual omission of N, K, Ca, S and Cu. Only the treatment with the simultaneous omission of Mg and liming (Mg/-lime) had a lower content than the complete treatment with statistical significance (-63 %).

Sulphur was less absorbed with the omission of N, Mg, Mg/-lime, Cu, Mn and Zn, especially the latter, which limited the absorption of S by leaves by 59.2 %, even exceeding the absorbance restriction of the control.

The contents of Cu, Zn and Mn were lower than the complete treatment for the majority of individual omission treatments. This may also be attributed to liming, since the increase of soil pH decreases the availability of micronutrients, thereby affecting their absorption by plants (Malavolta 2006). In this regard, the omission of Ca/-lime increased the absorption of Cu by 23.7 % and Zn by 26.15 %, and the omission of Mg/-lime increased the content of Mn by 9.7 %, relatively to the control. Viégas et al. (2009) considered Mn the most limiting micronutrient for assai palms growth on medium-textured Ferralsol, corroborating the present study and the importance of Mn as the most absorbed micronutrient.

The content of B in the complete treatment was 54.24 mg kg⁻¹, which is similar to the 51 mg kg⁻¹ obtained by Haag et al. (1992) and higher than the 31.34 mg kg⁻¹ observed by Viégas et al. (2009), with their complete treatments.

The content of B decreased by 68.17 % with the omission of the nutrient itself, while in the control it decreased by 41 %, if compared to the complete treatment. The absorption of B was favored by 23.09 %, with the omission of K. Viégas et al. (2004), studying the effect of doses of B on the growth of assai palm plants, concluded that the estimated optimum dose was 2.3 mg L⁻¹, and that at doses higher than 2.5 mg L⁻¹ of B, there was a negative effect on growth. Other studies have identified B as limiting to the development of Amazonian palms (Viégas 1993, Lins 2000, Viégas et al. 2009), but the present study does not corroborate this, as also observed by Haag et al. (1992).

The decreasing order of total nutrient accumulation in leaves, relatively to the complete treatment, was N > K > S > Ca > Mg > P > Mn > Zn > B > Cu (Table 3), showing that the nutrients most required by assai palms of the studied variety are N, K and S.

Viégas et al. (2009), who cultivated the assai palm BRS Pará cultivar on Ferralsol, showed the following decreasing order of macronutrients: N > Ca > K > Mg > S > P. For the peach palm, an Amazonian plant of the same botanical family, Fernandes et al. (2013) found the decreasing order of macronutrient accumulation to be N > K > Ca > P > Mg > S > Na. These data reveal that N and K are the nutrients most accumulated in the initial phase of development not only for assai palm, but also for other plants from the Arecaceae family. But a discordance, regarding the accumulation of S, remains, since, in the present study, it was the third most required macronutrient, whereas Viégas et al. (2009) and Fernandes et al. (2013) found it to be the fifth and sixth, respectively.

CONCLUSIONS

1. The initial growth of assai palm plants in sandy-textured Ferralsol is limited, on average, by 22 %, by the omission of the nutrients N, P, K, Ca, Mg and Cu;
2. The leaf dry mass production by assai palm is affected in a descending order by the omission of P > Cu > N > K > Mg, at an average of 31.9 %. Leaf area is restricted by the omission of the individual nutrients Ca > N > P > K > Mg > Zn;
3. The plant development measured by the shoot relative growth is affected by the lack of Ca > P > N > Mg > Cu > K, with an average reduction of 31 %;
4. The nutrients most required by the assai palm BRS Ver-o-Peso cultivar, as evidenced by the levels of nutrients accumulation in leaf dry mass, follow the order N > K > S > Ca > Mg > P > Mn > Zn > B > Cu.

ACKNOWLEDGMENTS

The authors thank the Postgraduate Program in Agrarian Sciences of the Universidade Federal Rural da Amazônia; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), for the scholarship granted to the first author; and Nelson Cruz de Oliveira, for helping in the experiment.

REFERENCES

- ALMEIDA, S. S.; AMARAL, D. D.; SILVA, A. S. L. Análise florística e estrutura de florestas de Várzea no estuário amazônico. *Acta Amazonica*, v. 34, n. 4, p. 513-524, 2004.
- ARMENGUAD, P. et al. Multilevel analysis of primary metabolism provides new insights into the role of potassium nutrition for glycolysis and nitrogen assimilation in *Arabidopsis* roots. *Plant Physiology*, v. 150, n. 2, p. 772-785, 2009.
- BOVI, M. L. A.; GODOY, J. R. G.; SPIERING, S. H. Respostas de crescimento de pupunheira à adubação NPK. *Scientia Agrícola*, v. 59, n. 1, p. 161-166, 2002.
- CLEMENT, C. R.; BOVI, M. L. A. Padronização de medidas de crescimento e produção em experimentos com pupunheira para palmito. *Acta Amazonica*, v. 30, n. 3, p. 349-362, 2000.
- CRAVO, M. S.; VIÉGAS, I. J. M.; BRASIL, E. C. *Recomendações de adubação e calagem para o Estado do Pará*. Belém: Embrapa Amazônia Oriental, 2010.
- CUNHA, A. C. M. et al. Papel da nutrição mineral na formação de raízes adventícias em plantas lenhosas. *Pesquisa Florestal Brasileira*, v. 58, n. 1, p. 35-47, 2009.
- EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA (Embrapa). *Manual de métodos de análise de solo*. 2. ed. Rio de Janeiro: Embrapa-CNPQ, 1997.
- EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA (Embrapa). *Métodos de análises bromatológicas de alimentos: métodos físicos, químicos e bromatológicos*. Pelotas: Embrapa Clima Temperado, 2010.
- EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA (Embrapa). *Sistema brasileiro de classificação de solos*. Brasília, DF: Embrapa Solos, 2013.
- FERNANDES, A. R.; MATOS, G. S. B.; CARVALHO, J. G. Deficiências nutricionais de macronutrientes e sódio em mudas de pupunheira. *Revista Brasileira de Fruticultura*, v. 35, n. 4, p. 1178-1189, 2013.
- FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS (FAO). *International soil classification system for naming soils and creating legends for soil maps*. Rome: FAO, 2015.
- GONÇALVES, A. S. *Crescimento, composição mineral e sintomas visuais de deficiências de macronutrientes e boro, em plantas de açaízeiro (Euterpe oleracea Mart.)*. 2004. 88 f. Dissertação (Mestrado em Agronomia) - Universidade Federal Rural da Amazônia, Belém, 2004.
- HAAG, H. P.; SILVA FILHO, N. L.; CARMELLO, Q. A. C. Carência de macronutrientes e de boro em plantas de açaí (*Euterpe oleracea Mart.*). In: CONGRESSO NACIONAL SOBRE ESSÊNCIAS NATIVAS, 2., 1992, São Paulo. *Anais...* São Paulo: Unipress, 1992. v. 1, p. 477-479.
- HAWKESFORD, M. J. Plant response to sulphur deficiency and the genetic manipulation of sulphate transporters to improve S-utilization efficiency. *Journal of Experimental Botany*, v. 51, n. 1, p. 131-138, 2000.
- INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA (IBGE). *Levantamento sistemático da produção agrícola paraense (LSPA/IBGE)*. Belém: IBGE, 2015.
- LAVIOLA, B. G.; DIAS, L. A. S. Teor e acúmulo de nutrientes em folhas e frutos de pinhão-manso. *Revista Brasileira de Ciência do Solo*, v. 32, n. 5, p. 1969-1975, 2008.
- LINS, P. *Resposta do coqueiro (Cocos nucifera L.) à aplicação de N, P, K e Mg nas condições edafoclimáticas de Moju - PA*. 2000. 89 f. Dissertação (Mestrado em Agronomia) - Faculdade de Ciências Agrárias do Pará, Belém, 2000.
- LOPES, M. L. B. *Mercado e distribuição dos retornos sociais do manejo do açaí para produção de frutos*. 2001. 73 f. Dissertação (Mestrado em Economia) - Universidade da Amazônia, Belém, 2001.
- MALAVOLTA, E. *Elementos de nutrição mineral de plantas*. São Paulo: Agronômica Ceres, 1980.
- MALAVOLTA, E. *Manual de nutrição de plantas*. São Paulo: Agronômica Ceres, 2006.
- MALAVOLTA, E.; VITTI, G. C.; OLIVEIRA, S. A. *Avaliação do estado nutricional das plantas: princípios e aplicações*. Piracicaba: Potafos, 1997.
- MENEZES, E. M. S.; TORRES, A. T.; SRUR, A. U. S. Valor nutricional da polpa de açaí (*Euterpe oleracea Mart*) liofilizada. *Acta Amazonica*, v. 38, n. 2, p. 311-316. 2008.
- MIRANDA, R. S. et al. Deficiência nutricional em plântulas de feijão-de-corda decorrente da omissão de macro e micronutrientes. *Revista Ciência Agronômica*, v. 41, n. 3, p. 326-333, 2010.
- OLIVEIRA, M. S. P. et al. *Cultivo do açaízeiro para produção de frutos*. Belém: Embrapa Amazônia Oriental, 2002. (Circular técnica, 26).
- OLIVEIRA, M. S. P.; FARIAS NETO, J. T.; PENA, R. S. *Açaí: técnicas de cultivo e processamento*. Fortaleza: Instituto Frutal, 2007.
- PRADO, R. M.; ROMUALDO, L. M.; ROZANE, D. E. Omissão de macronutrientes no desenvolvimento e no

estado nutricional de plantas de sorgo (cv. BRS 3010) cultivadas em solução nutritiva. *Científica*, v. 35, n. 2, p. 122-128, 2007.

TAIZ, L.; ZEIGER, E. *Plant physiology*. 4. ed. Porto Alegre: Artmed, 2013.

VIÉGAS, I. J. M. *Crescimento do dendezeiro (Elaeis guineensis, Jacq.), concentração, conteúdo e exportação de nutrientes nas diferentes partes de plantas com 2 a 8 anos de idade, cultivadas em Latossolo Amarelo distrófico; Tailândia, Pará*. 1993. 217 f. Tese (Doutorado em Agronomia) - Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo, Piracicaba, 1993.

VIÉGAS, I. J. M. et al. Avaliação da fertilidade de um Latossolo Amarelo textura média para o cultivo do açaizeiro no Estado do Pará. *Revista de Ciências Agrárias*, v. 52, n. 1, p. 23-35, 2009.

VIÉGAS, I. J. M. et al. Efeitos das omissões de macronutrientes e boro na sintomatologia e crescimento em plantas de açaizeiro (*Euterpe oleraceae* Mart). *Revista de Ciências Agrárias*, v. 50, n. 1, p. 129-141, 2008.

VIÉGAS, I. J. M. et al. Limitações nutricionais para o cultivo do açaizeiro em Latossolo Amarelo textura média, Estado do Pará. *Revista Brasileira de Fruticultura*, v. 26, n. 2, p. 382-384, 2004.