

Papéis Avulsos de Zoologia

Museu de Zoologia da Universidade de São Paulo

Volume 57(25):321-326, 2017

www.mz.usp.br/publicacoes
www.revistas.usp.br/paz

ISSN impresso: 0031-1049
ISSN on-line: 1807-0205

POST-EMBRYONIC DEVELOPMENT OF INTRAMANDIBULAR GLANDS OF *FRISELLA SCHROTTKYI* (FRIESE, 1900) (HYMENOPTERA: APIDAE) WORKERS

ANDRÉ HENRIQUE DE OLIVEIRA¹

LUCIANE CRISTINA DE OLIVEIRA LISBOA¹

JOSÉ EDUARDO SERRÃO²

JOSÉ COLA ZANUNCIO³

EDMILSON AMARAL DE SOUZA^{1,4}

ABSTRACT

Exocrine glands play important role in social organization of insects, such as caste and inter-caste differentiation. Due their functional and structural plasticity, morphological studies on such glands contribute to better understanding the biology of social bees. Therefore, the aim of the study was to characterize the sequence of the post-embryonic development of intramandibular glands of Friesella schrottkyi (Fries, 1900) (Hymenoptera: Apidae) workers using histological and histochemical analyses. The mandibles of pupae at different developmental stages and newly emerged adults were analyzed. The intramandibular glands of F. schrottkyi presented two types: class I glands, in the mandible epidermis and class III glands, inside the mandible cavity that open onto external surface. The intramandibular glands of F. schrottkyi developed during the transition from the prepupae to the white-eyed pupae, as shown by the morphological changes. Black-eyed pupae of F. schrottkyi presented fully developed intramandibular glands.

KEY-WORDS: Exocrine glands; Morphology; Development; Stingless bees.

INTRODUCTION

Stingless bee workers show plasticity of tasks they perform during their life, which can change according to the demands of the colony (Van Bethem *et al.*, 1995). The ability to perform different tasks

in the colony is related to the worker bee age and morpho-physiology of the exocrine glands. The morphology of these glands and the chemical composition of their secretions are parameters to analyze the functional plasticity of these glands (Cruz-Landim & Abdalla, 2002).

¹ Universidade Federal de Viçosa (UFV), Instituto de Ciências Biológicas e da Saúde, Campus Rio Paranaíba, Rodovia MG-230, km 7, CEP 38810-000, Rio Paranaíba, MG, Brasil.

² Universidade Federal de Viçosa (UFV), Departamento de Biologia Geral, Campus Viçosa, Avenida Peter Henry Rolfs, s/nº, Campus Universitário, CEP 36570-900, Viçosa, MG, Brasil.

³ Universidade Federal de Viçosa (UFV), Departamento de Entomologia, Campus Viçosa, Avenida Peter Henry Rolfs, s/nº, Campus Universitário, CEP 36570-900, Viçosa, MG, Brasil.

⁴ E-mail: edmilson.souza@ufv.br

The main exocrine secretory cells of the insects are class I and III. In class I, the epidermal cells are in close association with the endocuticle and release their secretions to the tegument surface by diffusion through the cuticle, whereas in class III glands, the secretory cells are detached from the epidermis, releasing the secretion through a canaliculus connected to a pore in the body surface (Noirot & Quennedey, 1991).

Mandibular glands can be of two types: ecto-mandibular gland (mandibular gland) and meso-mandibular gland (intramandibular gland). The ectomandibular glands are better studied than the mesomandibular glands. The term “mandibular glands” is used to distinguish them from the intramandibular glands (Cruz-Landim & Abdalla, 2002). According Nedel (1960), the mandibular glands of bees present secretory activity, spherical shape, and isolated cells with independent canaliculi. Moreover, epidermal hypertrophied cells inside the mandibles of some stingless bees were also described and classified as exocrine glands class I (Costa-Leonardo, 1978).

The function of the intramandibular gland is unknown, but its secretion may contribute to the lubrication of mandibles (Schoeters & Billen, 1994; Amaral & Caetano, 2006) or to release of scent trail pheromones (Martins *et al.*, 2013, 2015) in ants. In the stingless bee *Plebeia emerina* it is postulate that the intramandibular secretion helps in manipulation of propolis (Santos *et al.*, 2009). The post-embryonic development of the intramandibular glands has been reported in ants (Martins *et al.*, 2013), but is lacking in stingless bees.

Friesella is a monospecific genus of stingless bees with small workers (3-5 mm). *Friesella schrottkyi* (Friese, 1900) (Hymenoptera: Apidae) is a highly eusocial bee with labor division, overlapping generations, and food storage (Michener, 1974).

The aim of this study was to characterize the sequence of the post-embryonic development of intramandibular glands in the stingless bee *F. schrottkyi* using histological and histochemical analyses.

MATERIALS AND METHODS

Bee collection

Colonies of *F. schrottkyi* bees were maintained in Rio Paranaíba (19°11'37"S, 46°14'50"W), state of Minas Gerais, Brazil. Brood combs were removed from colonies and transferred to the Laboratory of Morphology of Arthropods at the Universidade Fed-

eral de Viçosa – Campus de Rio Paranaíba. Worker pupae at different developmental stages characterized by eye pigmentation (white-, pink-, brown-, and black-eyed pupae) were obtained from brood and newly emerged adults classified by tergite pigmentation collected in the nest (Bueno, 1981; Fagundes *et al.*, 2006).

Histology and histochemistry

The mandibles of *F. schrottkyi* pupae and adult workers were dissected and transferred to Zamboni fixative solution (Stefanini *et al.*, 1967), dehydrated in a graded ethanol series, embedded in historesin Leica, and sectioned at 3 µm thick. The sections were stained with toluidine blue-borax and analyzed using a light microscope.

Some mandible sections were submitted to Periodic acid-Schiff (PAS) staining (Bancroft & Gamble, 2008) for identification of neutral polysaccharides and glycoconjugates and to mercury-bromophenol (Bancroft & Gamble, 2008) for localization of total protein.

RESULTS

Histological analysis showed variations in the mandible according to the developmental phase of *F. schrottkyi* pupae. White- and pink-eyed pupae had only larval cuticle (Figs. 1A-B). The brown-eyed pupae showed both larval (external) and adult (onto the epidermis) cuticles (Figs. 1C-D). Black-eyed pupae had thicker adult cuticle (Fig. 1E).

The epidermis of mandibles in white- and pink-eyed pupae was formed by multiple cell layers with different morphology in the mandible regions. The mandible apex had irregular epidermal cells (Fig. 1A) and the basal region had columnar cells with basophil apex and central spherical nucleus with decondensed chromatin (Fig. 1B). Brown-eyed pupae showed columnar epidermal cells at the base of the mandible (Fig. 1C) and flattened ones in the middle and apical mandible regions (Fig. 1D). Black-eyed pupae and newly emerged workers showed flattened cells in the mandible (Figs. 1E-F).

Cells on the cavity of the mandible are involved in differentiation processes with granules and peripheral cytoplasm, and were found at higher quantity in the base mandible in white-, pink-, and brown-eyed pupae (Fig. 1C). Black-eyed pupae showed completely differentiated spherical cells with a large

nucleus with decondensed chromatin and cytoplasm with many granules (Fig. 1E). These cells had a canaliculus that opens on to the mandible surface in black-eyed pupae and newly emerged workers (Fig. 1F).

From white to brown-eyed pupae, the epidermal and spherical cells showed a strong positive reaction for PAS (Fig. 2A) and proteins (Fig. 2B). Epidermal cells of black-eyed pupae and newly emerged workers were slightly positive for PAS (Fig. 2C) and the granules of spherical cells were positive (Fig. 2D) and both cell types were positive for protein in black-eyed pupae and newly emerged workers (Figs. 2E-F). Sensilla were present on the mandible (Fig. 1D).

DISCUSSION

The morphology of *F. schrottkyi* intramandibular glands is similar to other Hymenoptera species (Cruz-Landim & Abdalla, 2002; Cruz-Landim *et al.*, 2011; Martins *et al.*, 2013). Epidermal cells of exocrine glands are class I with secretory features in newly emerged *F. schrottkyi* workers. However, large nuclei with decondensed chromatin in the spherical class III secretory cells suggest high activity (Silva-Zacarin *et al.*, 2008).

The new cuticle of brown-eyed *F. schrottkyi* pupae is synthesized in the previous developmental phas-




FIGURE 1: Mandible sections of *Friesella schrottkyi* (Hymenoptera, Apidae) worker pupae with toluidine blue. (A): Apex of the mandible in white-eyed pupae showing the larval cuticle (lc) and epidermal cells (ep). (B): Mandible of pink-eyed pupae showing epidermal cells (ep) with basophilic apex (*). (C-D): Mandible in brown-eyed pupae showing the new cuticle (ct), epidermal cells (ep) and secretory cells (sc). (E): Mandible in black-eyed pupae showing secretory cells (sc) and epidermal cells (ep). ct: cuticle; n: nucleus. (F): Mandible in newly emerged worker showing epidermal cells (ep), secretory cells (sc) and canaliculus (ca). n: nucleus.

es because the apex of the epidermal cells was positive for proteins, which may be used in the production of the new cuticle. The epidermal cells of immature insects synthesize the cuticle partly constituted by proteins, and the cuticular layers are still deposited to a lesser extent in adult insects as reported for *Melipona quadrifasciata* (Cruz-Landim *et al.*, 2011).

The multiple layers of epidermal cells in the mandible of white-eyed pupae changes from a single layer of squamous cells to the cubic epidermal cells in

black-eyed pupae, as found for *Plebeia emerina* (Santos *et al.*, 2009), *Melipona rufiventris* and *M. quadrifasciata anthioides* (Costa-Leonardo, 1978). The presence such epithelial cells, only in workers, could suggest some exclusive function for workers. Some authors attribute to the intramandibular secretion role in handling adhesive resins and propolis in *P. emerina* (Santos *et al.*, 2009; Cruz-Landim *et al.*, 2011).

Precursor cells of the class III glands found in the mandible of white-eyed pupae of *F. schrottkyi* are




FIGURE 2: Mandible sections of *Friesella schrottkyi* (Hymenoptera: Apidae) worker pupae submitted to histochemical tests. (A): White-eyed pupae showing the mandible cavity and epidermal cells (ep) Periodic acid-Schiff (PAS) positive. lc; larval cuticle. (B): Brown-eyed pupae showing the secretory cells (sc) and epidermal cells (ep) positive to protein. (C-D): Black-eyed pupae showing the epidermal cells (ep) and secretory cells (sc) Periodic acid-Schiff positive. ct: cuticle. (E-F): Black-eyed pupae showing the epidermal cells (ep) and secretory cells (sc) positive to protein. n: nucleus; ct: cuticle; s: sensilla; ca: canaliculus.

characterized by cytoplasm granules, but with complete differentiation in black-eyed pupae. According to Martins *et al.* (2013), the cells of the class III of the ant *Pachycondyla verenae* are also completely differentiated in black-eyed pupae.

The large amounts of glycoconjugates and proteins in the mandibular cells from white to brown-eyed pupae of *F. schrottkyi* suggest a high metabolic rate, perhaps due to the cellular reorganization. The positive staining for polysaccharides and glycoconjugates in the mandibles of the pupae is probably due to glycogen, which is an energetic fuel for high metabolic activity as shown in the pupae of the ant, *Solenopsis invicta*, which use about 75% of their carbohydrates during the pupal stage (Wheeler & Buck, 1992). Mandibles of white-eyed pupae of *Pachycondyla verenae* showed intense cell reorganization with positive reactions for glycoconjugates and proteins (Martins *et al.*, 2013).

The decrease of polysaccharides and glycoconjugates in the mandibles of *F. schrottkyi* during development of class III cells along the pupae suggest that this gland may be associated with secretion, likely found in the ants *Atta sexdens rubropilosa* (Amaral & Caetano, 2006) and *Pachycondyla verenae* (Martins *et al.*, 2013).

The development of intramandibular glands during the transition from prepupae to white-eyed pupae of *F. schrottkyi* differs from the lack of differentiation of the intramandibular glands in *P. verenae* prepupae (Martins *et al.*, 2013). The process of differentiation of intramandibular glands beginning in white-eyed pupae and completed in black-eyed pupae is still little known in stingless bees, but epidermal and non epidermal organs have complete differentiation in black-eyed pupae (Neves *et al.*, 2002, 2003; Azevedo *et al.*, 2008).

The post-embryonic development of intramandibular glands in *F. schrottkyi* occurs during the pupal period and completes the development during the black-eyed pupal stage.

RESUMO

As glândulas exócrinas desempenham importantes funções na organização social dos insetos, como a diferenciação de castas e inter-castas. Devido à sua plasticidade estrutural e funcional, estudos morfológicos destas glândulas contribuem para o melhor entendimento da biologia das abelhas sociais. Por isso, o objetivo deste estudo foi acompanhar o desenvolvimento pós-embriônico das glândulas intramandibulares de operárias da abelha sem ferrão Friesella schrottkyi (Fries, 1900) (Hymenoptera:

Apidae) com análises histológicas e histoquímicas. As mandíbulas das pupas em diferentes estágios do desenvolvimento e operárias recém-emergidas foram analisadas. As glândulas intramandibulares de F. schrottkyi são divididas em dois tipos: glândulas da classe I na epiderme da mandíbula e da classe III, inseridas na cavidade da mandíbula que se abrem na superfície externa. As glândulas intramandibulares de F. schrottkyi se desenvolvem durante a transição de pré-pupa para pupa de olho branco como observado pelas alterações morfológicas nas células. As pupas de olhos pretos de F. schrottkyi apresentaram glândulas intramandibulares completamente desenvolvidas.

PALAVRAS-CHAVE: Glândulas exócrinas; Morfologia; Desenvolvimento; Abelhas sem ferrão.

ACKNOWLEDGMENTS

We would like to thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Fundação de Amparo a Pesquisa do Estado de Minas Gerais (FAPEMIG) and Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA) for their financial support.

REFERENCES

- AMARAL, J.B. & CAETANO, F.H. 2006. The intramandibular gland of leaf-cutting ants (*Atta sexdens rubropilosa* 1908). *Micron*, 37:154-160.
- AZEVEDO, D.O.; GUS-MATIELLO, C.P.; RÖNNAU, M.; ZANUNCIO, J.C. & SERRÃO, J.E. 2008. Post-embryonic development of the antennal sensilla in *Melipona quadrifasciata anthidioides* (Hymenoptera: Meliponini). *Microscopy Research and Technique*, 71:196-200.
- BANCROFT, J.D. & GAMBLE, M. 2008. *Theory and practice of histological techniques*. London, Churchill Livingstone.
- BUENO, O.C. 1981. *Diferenciação dos ovários e determinação do número de ovários em Apis mellifera* (Doctoral Thesis). Universidade de São Paulo, São Paulo.
- COSTA-LEONARDO, A.M. 1978. Glândulas intramandibulares em abelhas sociais. *Ciência & Cultura*, 30:835-838.
- CRUZ-LANDIM, C. & ABDALLA, F.C. 2002. *Glândulas exócrinas das abelhas*. Ribeirão Preto, FUNPEC-RP.
- CRUZ-LANDIM, C.; GRACIOLI-VITTI, L.F. & ABDALLA, F.C. 2011. Ultrastructure of the intramandibular gland of workers and queens of the stingless bee, *Melipona quadrifasciata*. *Journal of Insect Science*, 11:1-9.
- FAGUNDES, I.B.; CAMPOS, L.A.O. & SERRÃO, J.E. 2006. Tergite pigmentation indicates hypopharyngeal gland developmental degree in *Melipona quadrifasciata* (Hymenoptera: Apidae: Meliponini). *Sociobiology*, 48:51-62.
- MARTINS, L.C.B.; DELABIE, J.H.C.; ZANUNCIO, J.C. & SERRÃO, J.E. 2013. Post-embryonic development of intramandibular glands

- in *Pachycondyla verenae* (Forel) (Hymenoptera: Formicidae) workers. *Sociobiology*, 60:154-161.
- MARTINS, L.C.B.; NASCIMENTO, F.S.; CAMPOS, M.C.G.; LIMA, E.R.; ZANUNCIO, J.C. & SERRÃO, J.E. 2015. Chemical composition of the intramandibular glands of the ant *Neoponera villosa* (Fabricius, 1804) (Hymenoptera: Ponerinae). *Chemoecology*, 25:25-31.
- MICHENER, C.D. 1974. *The social behavior of the bees*. Cambridge, Mass., The Belkman Press of Harvard University Press.
- NEDEL, O.J. 1960. Morphologie und Physiologie der Mandibeldrüse einiger Bienen-Arten (Apidae). *Zoomorphology*, 49:139-183.
- NEVES, C.A.; BHERING, L.L.; SERRÃO, J.E. & GITIRANA, L.B. 2002. FMRFamide-like midgut endocrine cells during the metamorphosis in *Melipona quadrifasciata anthidioides* (Hymenoptera, Apidae). *Micron*, 33:453-460.
- NEVES, C.A.; GITIRANA, L.B. & SERRÃO, J.E. 2003. Ultrastructural study of the metamorphosis in the midgut of *Melipona quadrifasciata anthidioides* (Apidae, Meliponini) worker. *Sociobiology*, 41:443-459.
- NOIROT, C. & QUENNEDEY, A. 1991. Gland, gland cell, glandular units: some comments on terminology and classification. *Annals of Society Entomologic of France*, 27:123-128.
- SANTOS, C.G.; MEGIOLARO, F.; SERRÃO, J.E. & BLOCHTEIN, B. 2009. Morphology of the head salivary and intramandibular glands of the stingless bee *Plebeia emerina* (Fries) (Hymenoptera, Meliponini) workers associated with propolis. *Annals of the Entomological Society of America*, 102:137-143.
- SCHOETERS, E. & BILLEN, J. 1994. The intramandibular gland, a novel exocrine structure in ants (Insecta, Hymenoptera). *Zoomorphology*, 114:125-131.
- SILVA-ZACARIN, E.C.; TABOGA, S.R. & SILVA DE MORAES, R.L. 2008. Nuclear alterations associated to programmed cell death in larval salivary glands of *Apis mellifera* (Hymenoptera: Apidae). *Micron*, 39:117-127.
- STEFANINI, M.; MARTINO, C.D.E. & ZAMBONI, L. 1967. Fixation of ejaculated spermatozoa for electron microscopy. *Nature*, 216:173-174.
- VAN BETHEM, F.D.J.; IMPERATRIZ-FONSECA, V.L. & VELTHUIS, H.W. 1995. Biology of the stingless bee *Plebeia remota* (Holmberg): observation and evolutionary implications. *Insectes Sociaux*, 42:71-87.
- WHEELER, D.A. & BUCK, N.A. 1992. Protein, lipid and carbohydrate use during metamorphosis in the fire ant, *Solenopsis xyloni*. *Physiological Entomology*, 17:397-403.

Aceito em: 22/05/2017

Publicado em: 13/06/2017

Editor Responsável: Kelli dos Santos Ramos

