

Bone changes caused by experimental *Solanum malacoxylon* poisoning in rabbits¹

José Ignacio Aguirre^{2,3}, María Soledad Gomar², Silvio Igal³, María Alejandra Quiroga², Enrique Leo Portiansky² and Eduardo Juan Gimeno^{2*}

ABSTRACT.- Aguirre J.I., Gomar M.S., Igal, S., Quiroga M.A., Portiansky E.L & Gimeno E.J. 2005. **Bone changes caused by experimental *Solanum malacoxylon* poisoning in rabbits.** [Alterações ósseas causadas na intoxicação experimental por *Solanum malacoxylon* em coelhos.] *Pesquisa Veterinária Brasileira* 25(1):34-38. Institute of Pathology "Prof. Dr. Bernardo Epstein", School of Veterinary Sciences, National University of La Plata, P.O.Box 296, (1900) La Plata, Argentina. E-mail: ejgimeno@fvc.unlp.edu.ar

The aim of this study was to describe the bone changes observed after a daily oral administration of the calcinogenic plant *Solanum malacoxylon* (syn. *S. glaucophyllum*) (*Sm*) during 9 days. The *Sm*-poisoned rabbits had an increase of bone resorption in the endosteal surface of the cortical zone and also in the surface covered by osteoblasts of the primary and secondary spongiosa of the trabecular bone compartment. Moreover, the epiphyseal growth plates in long bones appeared narrower than in the control rabbits, with reduction of the proliferative and hyperthrophic chondrocyte zones. The electron microscopic study revealed a significant decrease of proteoglycans in the hyperthrophic chondrocyte zone evidenced by a significant reduction of ruthenium red positive granules in the poisoned rabbit. Altogether, these data suggest that cell differentiation may play a pivotal role in the pathogenesis of *Sm*-induced bone lesions.

INDEX TERMS: *Solanum malacoxylon*, *S. glaucophyllum*, calcinogenic plant, rabbit, bone tissue, growth plate.

INTRODUCTION

Solanum malacoxylon (*Sm*) (syn. *S. glaucophyllum*) is a calcinogenic plant responsible for producing the Enzootic Calcinosis of cattle and sheep in Argentina, Brazil, Paraguay and Uruguay. The disease is characterized by the calcification of soft tissues, especially aorta, heart, lungs, and kidneys (Worker & Carrillo 1967, Puche & Bingley 1995, Tokarnia et al. 2000). The plant is highly toxic for cattle and causes considerable economic losses in one of the most important meat production areas of Argentina (Okada et al. 1977). Grazing animals in other parts of the world develop a similar disease but induced by other calcinogenic plants (Morris 1978, Jubb et al. 1993).

Sm contains high levels of 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) as glycoside derivatives (Wasserman et al. 1976). In fact, *Sm* contains not only 1,25(OH)₂D₃ but also other metabolites, like vitamin D₃ and 25(OH)D₃ (Esparza et al. 1982).

Bone changes have been studied in natural occurring and experimentally induced calcinosis. Increased bone formation (Döbereiner & Dämmrich 1974) and osteosclerosis (Krook et al. 1975) have been reported in spontaneous disease. Fibrosis of the bone marrow, endosteal and periosteal hyperostosis, and abnormal formation of ground substance by osteoblasts and fibroblasts were also recorded (Dämmrich et al. 1975).

It has been shown that bone tissue responds differently when a calcinogenic plant is administered. It has been suggested that the diversity of responses may depend upon exposure time to the plant, species sensitivity and bone type specificity (Barros et al. 1996).

Many researchers found that the increase in bone mass observed after *Sm* ingestion may be due to an increase in osteoid production by osteoblasts (Carrillo 1973, Dämmrich et al. 1975, Norrdin et al. 1975, Gimeno 1980). However, inhibition of bone resorption has also been reported in different animal species

¹Received on May 25, 2004.

Accepted for publication on September 10, 2004.

²Institute of Pathology "Prof. Dr. Bernardo Epstein", School of Veterinary Sciences, National University of La Plata. P.O.Box 296, (1900) La Plata.

*Author for correspondence. E-mail: ejgimeno@fvc.unlp.edu.ar

³Department of Electron Microscopy, School of Veterinary Sciences, National University of La Plata, P.O.Box 296, (1900) La Plata, Argentina.

(Krook et al. 1975, Santos et al. 1976, Gimeno 1980, Riet-Correa et al. 1987).

Surprisingly, very few data has been published regarding the *Sm* effects on cartilage. Santos et al. (1976) have reported retarding effects in both articular and epiphyseal cartilages in growing rabbits. Similar observation were reported in guinea pigs (Ousavaplangchai 1972) and rats (Gimeno 1980). Microscopic and ultrastructural features have been widely studied in normal and rachitic cartilages (Appleton 1988). However, as far as we know, epiphyseal growth plate has not been ultrastructurally studied either in hypervitaminosis D nor in enzootic calcinosis.

The aim of this study was to describe the histological findings in bone tissue and in cartilage of the growth plate and to analyse the ultrastructural features of the growth plate abnormalities after a subacute *Sm* poisoning in rabbits.

MATERIALS AND METHODS

Plant material was collected from several areas of the Province of Buenos Aires, Argentina, and dried at 37°C for 48 h. Leaves were separated from stems and milled to a fine powder. Four white New Zealand rabbits (2-2.5 kg/bw) received 300 mg of dry powdered leaves of *Sm* orally on daily basis during 9 days. The animals received a commercial diet (Cargill, Pilar, Córdoba, Argentina) containing a concentration of 0.87% and 0.78% of calcium and phosphorus, respectively. The grounded leaves were mixed with water and administered intragastrically with a flexible tube. Two rabbits received a placebo during 9 days and were used as controls. The body weight of each animal was recorded twice a week. Clinical signs were observed and recorded every day. The animals were carefully necropsied. Distal femur, proximal tibia and ribs, and primary and secondary spongiosa were processed for histology. Tissues were fixed in 10% buffered formalin, decalcified with EDTA (ethylenediaminetetracetic acid), embedded in paraffin, sectioned at 5µm and stained with haematoxylin and eosin.

For electron microscopy, samples from growth plate of proximal tibia were cut into small cubes of 2-3 mm, fixed in Karnovsky's solution (2.5% glutaraldehyde and 2% paraformaldehyde PBS 0.1M, pH 7.4) containing 0.05% ruthenium red for 2 h. Samples were then washed in buffer containing 0.05% ruthenium red, posfixed in 1% osmium tetroxide containing 0.05% ruthenium red, dehydrated in a gradient of ethanol (50, 70, 80, 90 and 95%) and impregnated in propilen oxidize. Finally, all the samples were embedded in epoxi resin (Poly/Bed 812, Polysciences 18976-2590). Ultrathin sections were cut with a SuperNova ultramicrotome (Reichter-Jung, Austria) and observed using a transmission electronic microscope JEM-1200 FORMER II (JEOL Co. Ltd., Tokyo).

RESULTS

The *Sm*-poisoned rabbits showed anorexia, loss of weight and also diarrhoea and rhinitis. On macroscopic examination, small calcified plaques were observed in the entire length of the aorta. Microscopic calcifications, von Kossa positive, were seen in the aorta, heart and kidneys.

The histologic examination of the long bones revealed that *Sm*-poisoned rabbits had an increase in bone resorption areas in the endosteal surface of the cortical bone (Fig. 1 B) in comparison with control bones (Fig. 1 A). An increase in bone resorption and in the surface covered by osteoblasts were also observed in the

primary and secondary spongiosa of the trabecular bone compartment (Fig. 1 D) differing with what was observed in non-poisoned rabbits (Fig. 1 C). Mesenchymal metaplasia was observed in the periosteum with the formation of cartilage (Fig. 1 E).

Growth plates were narrowed and showed an irregular pattern due to a focal or diffuse reduction in the number of chondrocytes in the proliferative and hypertrophic zones. Moreover, disorganized arrangement of chondrocytes and a shortening of the longitudinal septa in the calcified cartilage zone were also observed (Fig. 1 F).

Ultrastructural studies revealed early degenerative changes in chondrocytes of the proliferative zone, and a reduction in the number and size of proteoglycan granules of the territorial and interterritorial areas in the hypertrophic zone of the cartilage of the growth plate (Fig. 2). The bones of control animals did not show alterations.

DISCUSSION

In the study, the growth plate was irregular and narrow due to a reduction in the number of chondrocytes of the proliferative and hypertrophic zones. This could be explained by an inhibitory effect of 1,25(OH)₂D₃ on cell division (Walters 1992). This feature was described in the skin of *Sm*-poisoned animals (Gimeno et al. 2000). The decrease in the concentration of ruthenium red-positive granules, particularly in the pericellular region of the chondrocytes was accompanied by an increase in matrix calcification. Ruthenium red-positive granules represent an accumulation of unmodified proteoglycan aggregated from which calcium ions are excluded and so preventing calcification (Appleton 1988). It has been reported that proteoglycans are completely desegregated prior to cartilage calcification (Campo & Romano 1986, Yoshioka & Yagi 1989).

The presence of cartilage in the subperiosteum of the ribs in *Sm*-treated rabbits can also be explained by the direct effect of 1,25(OH)₂D₃ stimulus, since mesenchymal metaplasia has been frequently observed in enzootic calcinosis (Collier 1927, Worker & Carrillo 1967, Done et al. 1976, Morris 1978, Barros et al. 1981, Tokarnia et al. 2000).

We have also described an increased bone remodeling in the secondary spongiosa. This finding could be explained by the calcinogenic effect of *Sm* that enhance bone resorption, by increasing the number and activity of osteoclasts as seen with 1,25 (OH)₂D₃ (Baylink et al. 1973, Grise et al. 1990). Other researchers (Riet-Correa 1987, Barros et al. 1996), in contrast, showed that acute *Sm*-poisoning produces a cytotoxic inhibition of osteocytes, osteoblasts and osteoclasts in cortical bone.

Differences in the bone response to *Sm* have been observed both *in vivo* and *in vitro*. Even a dual effect have been reported in relation to the concentration of *Sm* purified extract: *in vitro* studies found a stimulatory effect on bone resorption at low concentrations and a clear inhibition at higher concentrations (Stern et al. 1978).

Vitamin D metabolites also modulate cell differentiation in cartilage. It has been demonstrated that 1,25(OH)₂D₃ promotes chondrocytes differentiation of chondrocytes along the endochondral differentiation cascade (Gerstenfeld et al. 1990).




Fig. 1. Effects of subacute *Solanum malacoxylon* poisoning on different bone structures. (A) Transversal section of the mid-shaft of a rib in a control rabbit. HE, bar 100 μ m. (B) Transversal section of the mid-shaft of a rib of a *Sm*-treated rabbit. Note the prominent eroded surfaces present at the endosteal envelope (arrows). HE, bar 100 μ m. (C) Trabecular bone of a distal femur metaphysis. Control rabbits showed resting surfaces. Toluidine blue, bar 40 μ m. (D) A significant increase in bone remodelling was observed in the *Sm*-poisoned rabbits. Note the eroded surfaces (small arrows), osteoclast surface, and the surfaces covered by osteoid tissue (large arrows). Toluidine blue, bar 40 μ m. (E) Prominent areas of chondrocytes were observed in the periosteum of ribs of *Sm*-treated rabbits, just beneath its outer layer (*). HE, bar 40 μ m. (F) Proximal tibia epiphyseal growth plate of a *Sm*-poisoned rabbit. The growth plate was irregular and narrow due to focal or diffuse reduction of proliferative and hypertrophic chondrocyte zones. The primary spongiosa appeared as a dense metaphyseal band of woven bone, and a reduction in secondary spongiosa development was also seen. HE, bar 40 μ m.




Fig. 2. Effect of subacute *Solanum malacoxylon* poisoning at ultrastructural level. (A) Hypertrophic chondrocytes of control rabbit growth plates. Note the small amount of ruthenium red positive granules in the pericellular region (*). Bar 500nm. (B) High magnification of the territorial zone (*) from A. Note the calcium deposits associated with vesicles and collagen fibers (arrows). Bar 100 nm. (C) Hypertrophic chondrocytes of *Sm*-poisoned rabbit growth plates. Note the increased amount of ruthenium red positive granules in the pericellular region (*) and calcium deposits forming large irregular masses in the territorial zone (arrows). Bar 500nm. (D) High magnification of the territorial zone (*) showed in C. Note the increased amount of calcium deposits associated with vesicles (small arrows) and collagen fibers (large arrow). Bar 200nm.

Chondrocyte response to vitamin D metabolites depends on the state of differentiation. Chondrocytes located at the resting zone respond primarily to $24,25(\text{OH})_2\text{D}_3$, whereas chondrocytes located at the proliferative zone respond primarily to $1,25(\text{OH})_2\text{D}_3$. Vitamin D has direct effects on cartilage and is not mediated by changes in serum or extracellular calcium (Schwartz et al. 1988, Boyan et al. 1992). The $1,25(\text{OH})_2\text{D}_3$ has been shown to have a number of important functions in the endochondral bone formation cascade. It not only stimulates chondrocyte differentiation but also inhibits cell division in chondrocytes of both resting zone and growth zones (Schwartz et al. 1989).

The effects of $1,25(\text{OH})_2\text{D}_3$ on cell differentiation have been extensively studied in both normal and abnormal conditions, e.g. bone disease, psoriasis, and cancer (Clemens et al. 1983, Rice et al. 1992, Walters 1992, Bikle & Pillai 1993). Changes in cell differentiation induced by calcinogenic plants have been

analysed in skin, aorta and lung (Barros & Gimeno 2000, Gimeno et al. 2000, 2004, Gomar et al. 2000, Portiansky et al. 2002).

Altogether, these data suggest that as occurs with $1,25(\text{OH})_2\text{D}_3$ (Bikle et al. 1990), *Sm* may induce different responses or effects, depending on exposure time to the plant, species sensitivity and bone type specificity.

Our data also suggest that the administration of *Sm* may induce a derangement of endochondral ossification by affecting cell differentiation in the epiphyseal growth plate and in the primary spongiosa. Alteration in cell differentiation during the intramembranous ossification may also be present because cartilage tissue was frequently observed beneath the outer layer of the periosteum.

Acknowledgements. The authors would like to thank Mrs. Rosa Villegas for her technical assistance. Financial support was provided in part by grants from the "Agencia Nacional de Promoción Científica", the "Acade-

mia Nacional de Agronomía y Veterinaria" and the "Consejo Nacional de Investigaciones Científicas (CONICET)", Argentina. ELP and EJG are Research Career Members of CONICET.

REFERENCES

- Appleton J. 1988. The ultrastructural distribution of proteoglycans in normal and rachitic growth cartilage from the mandibular condyle of the rat. *Archs Oral Biol.* 33: 379-381.
- Barros S.S. & Gimeno E.J. 2000. Cell differentiation and bone protein synthesis in the lung of sheep affected by spontaneous calcosinosis. *J. Comp. Pathol.* 123:270-277.
- Barros S.S., Russowski D. & Grando S. 1996. Ultra-estrutura das alterações ósseas na intoxicação experimental por *Solanum malacoxylon* em coelhos. *Pesq. Vet. Bras.* 16:81-86.
- Barros S.S., Tabone E., Dos Santos M., Andujar M. & Grimaud J.A. (1981). Histopathological and ultrastructural alterations in the aorta in experimental *Solanum malacoxylon* poisoning. *Virchow Archiv (Cell Pathol.)* 35:167-175.
- Baylink D., Sipe J., Wergedal J. & Whitemore O. 1973. Vitamin D enhance osteocytic and osteoclastic bone resorption. *Am. J. Physiol.* 221:1345-1347.
- Bikle D., Halloran P., McGalliard-Cone C. & Morey-Holton E. 1990. Different responses of trabecular and cortical bone to 1,25(OH)₂D₃. *Am. J. Physiol.* 259:716-722.
- Bikle D.D. & Pillai S. 1993. Vitamin D, calcium, and epidermal differentiation. *Endocr. Rev.* 14:3-19.
- Boyan B., Schwartz Z. & Swain L.D. 1992. *In vitro* studies on the regulation of endochondral ossification by vitamin D. *Crit. Rev. Oral Biol. Med.* 3:15-30.
- Campo R.D. & Romano J.E. 1986. Changes in cartilage proteoglycans associated with calcification. *Calcif. Tissue Int.* 39:175-184.
- Carrillo B.J. 1973. Efecto de la intoxicación de *Solanum malacoxylon* en el sistema óseo. *Revta Investnes Agropec., INTA*, 40:67-77.
- Clemens, T.L., Adams, J.S., Horiuchi, B.A., Cho, H., Tsuchiya, Y., Matsuo, N., Suda, T. & Holick, M.F. 1983. Interaction of 1,25-dihydroxyvitamin-D₃ with keratinocytes and fibroblasts from skin of normal subjects and a subject with vitamin-D-dependent rickets type II: A model for study the mode of action of 1,25-dihydroxyvitamin-D₃. *J. Clin. Endocr. Metab.* 56:824-830.
- Collier W.A. 1927. Zur Kenntnis als Enteeque bezeichneten Krankheit der Rinder in der Provinz Buenos Aires. *Zeitschr. Infektionskrankh. Parasit. Krankh. Hyg. Haust.* 31:81-92.
- Dämmrich K., Döbereiner J., Done S.H. & Tokarnia C.H. 1975. Skelettveränderungen nach Vergiftungen mit *Solanum malacoxylon* bei Rindern. *Zentralbl. Veterinärmed. Reihe A* 22:313-329.
- Döbereiner J. & Dämmrich K. 1974. Skelettveränderungen bei Rindern nach Vergiftungen mit *Solanum malacoxylon* Sendtner. *Verh. Dtsch. Ges. Pathol* 58:323-326.
- Done S.H., Döbereiner J. & Tokarnia C.H. 1976. Systemic connective tissue calcification in cattle poisoned by *Solanum malacoxylon*: a histological study. *Brit. Vet. J.* 132:28-38.
- Esparza M.S., Vega M. & Boland R.L. 1982. Synthesis and composition of vitamin D-3 metabolites in *Solanum malacoxylon*. *Bioch. Bioph. Acta* 719:633-640.
- Gerstenfeld L.C., Kelly C.M. Von Deck M. & Lian J.B. 1990. Effect of 1,25-dihydroxyvitamin D₃ on induction of chondrocyte maturation in culture: extracellular matrix gene expression and morphology. *Endocrinol.* 126:1599-1609.
- Gimeno E.J. 1980. Estudio histopatológico de las alteraciones óseas en ratas carenciadas en vitamina D y ratas intoxicadas con *Solanum malacoxylon*. *Analecta Vet.* 12:61-108.
- Gimeno E.J., Costa E.F., Gomar M.S., Massone A.R. & Portiansky E.L. 2000. Effects of plant induced hypervitaminosis D on cutaneous structure, cell differentiation and cell proliferation in cattle. *J. Vet. Med. A* 47:201-211.
- Gimeno E.J., Portiansky E.L., Gomar M.S., Costa E.F., Massone A.R., Alonso C.R., Dallorso M.E. & Barros S.S. 2004. Calcosinosis in ruminants due to plant poisoning: Contributions on the pathogenesis. Chapter 12. In: Acamovic T., Stewart C.S. & Pennycott T.W. (ed.) *Poisonous Plants and Related Toxins*. CAB International (HB ISBN 0 85199 6140).
- Gomar M.S., Portiansky E.L., Dallorso M.E., Barros S.S., Costa E.F. & Gimeno E.J. 2000. Cell differentiation and bone protein synthesis in the aorta and lung induced by calcinogenic plants, p.435-438. In: Norman A.W., Bouillon R. & Thomasset M. (ed.) *Vitamin D Endocrine System: Structural, Biological, Genetics and Clinical Aspects*. University of California Press, Davis.
- Grise M., MacKay C. & Popoll S. 1990. Effects of 1,25(OH)₂D₃ on osteoclast number and cytochemistry in normal and osteopetrotic rabbits. *Am. J. Anat.* 189:261-266.
- Jubb K.V.F., Kennedy P.C. & Palmer N. 1993. *Pathology of Domestic Animals*. 4th ed. Academic Press, San Diego.
- Krook L., Wasserman R.H., Shively J.N., Tashjian A.H. Jr, Brokken T.D. & Morton J.F. 1975. Hypercalcemia and calcosinosis in Florida horses: implication of the shrub *Cestrum diurnum*, as the causative agent. *Cornell Vet.* 65:26-56.
- Morris K.L.M. 1978. Plant induced calcosinosis. *Vet. Human Toxicol.* 24:34-48.
- Norrdin R.W., Barros C.S.L., Queille ML, Carré M. & Miravet L. 1975. Acute effects of *Solanum malacoxylon* on bone formation rates in growing rats. *Calcif. Tissue Int.* 28:239-243.
- Okada K.A., Carrillo B.J. & Tilley M. 1977. *Solanum malacoxylon* Sendtner: A toxic plant in Argentina. *Econ. Botany* 31:225-236.
- Ousavaplangchai L. 1972. Vergleichende Untersuchungen zur Histopathologie der Intoxikation mit Vitamin D₃ und *Solanum malacoxylon* beim Meerschweinchen. Dissertation, Wien.
- Portiansky EL, Alonso CR, Costa EF & Gimeno EJ. 2002. Collagenous and elastic system fibres in the aorta of cattle poisoned by *Solanum glaucophyllum*. *Vet. Rec.* 150:42-45.
- Puche R.C. & Bingley J.B. 1995. Calcosinosis of Cattle in Argentina. First English edition. National University of Rosario Editors (ISBN No. 950-673-112-8).
- Rice R.H., Qin Q. & Pilato A.L. 1992. Keratinocyte differentiation markers: Involucrin, transglutaminase, and toxicity. *J. Natl Inst. Cancer Monogr.* 13:87-91.
- Riet-Correa F., Schild A.L., Méndez M.C., Wasserman R. & Krook L. 1987. Enzootic calcosinosis in sheep caused by the ingestion of *Nierembergia veitchii* (Solanaceae). *Pesq. Agropec. Bras.* 7:85-95.
- Santos M.N., Nunes V.A., Nunes I.J., Barros S.S., Wasserman R.H. & Krook L. 1976. *Solanum malacoxylon* toxicity: inhibition of bone resorption. *Cornell Vet.* 66:566-589.
- Schwartz Z., Schlader D.L., Swain L.D. & Boyan B.D. 1988. Direct effects of 1,25-dihydroxyvitamin D₃ and 24,25 dihydroxyvitamin D₃ on growth zone and resting zone chondrocyte membrane alkaline phosphatase and phospholipase-A₂, specific activities. *Endocrinol.* 123:2878-2884.
- Schwartz Z., Schlader D.L., Ramirez V., Kennedy M.B. & Boyan B.D. 1989. Effects of vitamin D metabolites on collagen production and cell proliferation of growth zone and resting zone cartilage cells in vitro. *J. Bone Miner. Res.* 4(2):199-207.
- Stern P.H., Ness E.M & DeLuca H.F. 1978. Responses of fetal bones to *Solanum malacoxylon* in vitro: A possible explanation of previous paradoxical results. *Molec. Pharmacol.* 14:357-365.
- Tokarnia C.H., Döbereiner J. & Peixoto P.V. 2000. *Plantas Tóxicas do Brasil*. Editora Helianthus, Rio de Janeiro, p.188-199.
- Walters M.R. 1992. Newly identified actions of vitamin D endocrine system. *Endocr. Rev.* 13:719-764.
- Wasserman R.H., Henion J.D., Haussler M.R. & Mc Cain T.A. 1976. Calcinogenic factor in *Solanum malacoxylon*: evidence that it is 1,25-dihydroxyvitamin D₃-glycoside. *Science* 194:853-855.
- Worker N.A. & Carrillo B.J. 1967. 'Enteeque seco', calcification and wasting in grazing animals in Argentina. *Nature* 215:72-74.
- Yoshioka C & Yagi T. 1989. Electron microscopic study of condylar cartilage of rat mandible stained with ruthenium red: proteoglycans and hypertrophic chondrocytes. *J. Cranifac. Genet. Dev. Biol.* 9:303-314.