










Use of remdesivir for patients with Covid-19: a review article

 Thomás Cavalcanti Pires de Azevedo¹
 Pedro Cavalcanti Pires de Azevedo¹
 Robson Natario Silveira Filho¹
 Arthur Ricardo Vilar Scavuzzi de Carvalho¹
 Murilo Lobo Cezarotti Filho¹
 Fabiano Timbó Barbosa^{1,2,3}
 Célio Fernando de Sousa-Rodrigues^{1,2,5}
 Thiago José Matos-Rocha^{1,5}
 Fernando Wagner da Silva Ramos^{1,4,5}

1. Centro Universitário Cesmac, Maceió, AL, Brasil.

2. Universidade Federal de Alagoas – UFAL, Maceió, AL, Brasil.

3. Hospital Geral do Estado de Alagoas – HGE, Maceió, AL, Brasil.

4. Secretaria de Estado da Saúde de Alagoas – SESAU, Maceió, AL, Brasil.

5. Universidade Estadual de Ciências da Saúde de Alagoas – UNCISAL, Maceió, AL, Brasil.

<http://dx.doi.org/10.1590/1806-9282.66.6.838>

SUMMARY

The etiological agent of COVID-19, which causes severe respiratory diseases such as pneumonia and pulmonary insufficiency, has been confirmed as a new coronavirus, now known as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). There is currently no authorized medication for the treatment of COVID-19. No vaccines have been authorized. Thus, this study aimed at conducting a review of the use of Remdesivir in patients with COVID-19. The following electronic databases were used MEDLINE, SCIELO, LILACS, and PUBMED. On May 1, Remdesivir received emergency use authorization from the Food and Drug Administration. Remdesivir is currently the most promising molecule in the treatment of COVID-19, taking into account its broad antiviral spectrum (considering the genetic sequences of the virus, it is expected to maintain activity against SARS-CoV-2). There is *in vitro* and *in vivo* information available for coronaviruses, as well as an extensive clinical safety database (from a clinical trial of the Ebola virus and in the context of the Monitored Emergency Use of Unregistered and Investigational Interventions - MEURI). Further studies are relevant as available data on the efficacy and safety of Remdesivir against SARS-nCoV-2 are limited.

KEYWORDS: Betacoronavirus. Coronavirus Infections/therapy. Coronavirus.

INTRODUCTION

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) is a newly-emerging human infectious coronavirus, originating in Wuhan, China, that has been spreading rapidly in China and other countries since December 2019¹. SARS-CoV-2 is a β -coronavirus, which is an enveloped non-segmented positive-sense

RNA virus (subgenus sarbecovirus, Orthocoronavirinae subfamily)².

The World Health Organization (WHO) named this novel coronavirus disease COVID-19, and there have been confirmed cases in 189 countries or territories outside China, including Japan, the United States

DATE OF SUBMISSION: 08-May-2020

DATE OF ACCEPTANCE: 23-May-2020

CORRESPONDING AUTHOR: Fernando Wagner da Silva Ramos

Rua Hélio Pradines, 737, Ed. Mont Alverne, Apt. 503, Ponta Verde, Maceió, Alagoas, Brasil - 57035-220

Tel: +55 (82) 99321-3373

E-mail: nandobiomedico@hotmail.com

of America, Italy, Iran, and Brasil³. In Brasil, 52,771 deaths have been caused by COVID-19 and there have been 1,151,479 confirmed cases of the disease recorded nationwide as of June, 23th, 2020⁴.

The clinical features of COVID-19 are varied, ranging from an asymptomatic state to acute respiratory distress syndrome and multi-organ dysfunction. The common clinical features include fever (not in all cases), cough, sore throat, headache, fatigue, myalgia, and breathlessness. Conjunctivitis (pink eye) has also been described⁵.

There is no evidence from randomized clinical trials (RCTs) that any potential therapy improves outcomes in patients with suspected or confirmed COVID-19. This review summarizes current evidence on the use of Remdesivir for COVID-19 and provides a summary of the current clinical experience and treatment guidelines for this epidemic Novel Coronavirus.


METHODS

A literature search was performed in a Medical Literature Analysis and Retrieval System Online (MEDLINE) via PubMed (1966 to January 2020), available through the following link: <https://www.ncbi.nlm.nih.gov/pubmed/>; Scientific Electronic Library Online (SCIELO), available at <https://www.scielo.org/> and Literatura Latino Americana e do Caribe em Ciências da Saúde (LILACS), available through the following link: <https://bvsalud.org/>, using the following terms: 2019-nCoV, COVID-19, SARS-CoV-2, Coronavirus and Treatment, to find articles published from January 5 to April 30, 2020. Moreover, we used the findings of literature retrieved by searching authoritative texts and manual searches in WHO reports. We checked the reference lists of all studies identified by the above methods. Studies were excluded if old data was used, if the topics were inappropriate or not pertinent to the purpose of the study.

Use of Antiviral Remdesivir

The Food and Drug Administration (FDA), the agency that regulates medicines in the United States, has approved the use of Antiviral Remdesivir (GS-5734TM) (Figure 1) in the treatment of severe cases of COVID-19, the disease caused by the new coronavirus (Sars-CoV-2)⁶. The drug, from drugmaker Gilead, was originally developed to fight Ebola, but with no success⁷.

FIGURE 1. CHEMICAL STRUCTURE OF REMDESIVIR.



Remdesivir (also GS-5734) is a monophosphoramidate prodrug of an adenosine analog that has a broad antiviral spectrum including filoviruses, paramyxoviruses, pneumoviruses, and coronaviruses. Remdesivir is a prodrug that is metabolized into its active form GS-441524, an adenine nucleotide analog that interferes with the activity of viral RNA polymerase and promotes evasion of proofreading by viral exoribonuclease, leading to inhibition of viral RNA synthesis. Remdesivir acts early in infection and decreases viral RNA levels in a dose-dependent manner that parallels impairment of viral load *in vitro*⁸.

Remdesivir has demonstrated *in vitro* and *in vivo* activity in animal models against the viral pathogens that cause MERS and SARS, which are coronaviruses structurally similar to SARS-CoV-2, the coronavirus that causes COVID-19^{9,10}.

Concerning its metabolism, it has been shown that, upon intravenous (IV) administration of a 10 mg/kg dose in Rhesus Monkeys, Remdesivir exhibited a short plasma half-life ($t_{1/2} = 0.39$ h) with rapid systemic elimination followed by the appearance of transient systemic levels of a key intracellular intermediate alanine metabolite and more persistent levels of GS-441524 (detectable for over 24h in plasma)¹¹.

Regarding the clinical use of Remdesivir, the results were published by Grein et al.⁹, in which patients received a 10-day course of Remdesivir, consisting of 200 mg administered intravenously on day 1, followed by 100 mg daily for the remaining 9 days of the treatment. It was observed, at the end of the follow-up, that of the 61 patients who received at least one dose of Remdesivir, data from 8 could not be analyzed (including 7 patients with no post-treatment data

and 1 with a dosing error). Of the 53 patients whose data were analyzed, 22 were in the United States, 22 in Europe or Canada, and 9 were in Japan. At baseline, 30 patients (57%) were receiving mechanical ventilation and 4 (8%) were receiving extracorporeal membrane oxygenation.

During a median follow-up of 18 days, 36 patients (68%) had an improvement in oxygen-support class, and 17 of 30 patients (57%) receiving mechanical ventilation were extubated. A total of 25 patients (47%) were discharged, and 7 patients (13%) died; mortality was 18% (6 out of 34) among patients receiving invasive ventilation and 5% (1 of 19) among those not receiving invasive ventilation. In this cohort of patients hospitalized for severe Covid-19 who were treated with compassionate-use Remdesivir, clinical improvement was observed in 36 of 53 patients (68%). Measuring the efficacy of this approach will require ongoing, randomized, placebo-controlled trials of Remdesivir therapy.

In another published study, eligible patients were adults (aged ≥ 18 years) admitted to the hospital with laboratory-confirmed SARS-CoV-2 infection, with an interval from symptom onset to the moment of enrollment of 12 days or less, oxygen saturation of 94% or less on room air, or a ratio of arterial oxygen partial pressure to fractional inspired oxygen of 300 mm Hg or less, and radiologically confirmed pneumonia. Patients were randomly assigned in a 2:1 ratio to intravenous Remdesivir (200 mg on day 1 followed by 100 mg on days 2–10 in single daily infusions) or the same volume of placebo infusions for 10 days. 237 patients were enrolled and randomly assigned to a treatment group (158 to Remdesivir and 79 to placebo); One patient in the placebo group who withdrew after randomization was not included in the ITT population.

Remdesivir use was not associated with any difference in the time for clinical improvement (hazard ratio 1.23 [95% CI 0.87–1.75]). Although not statistically

significant, patients receiving Remdesivir had a numerically faster time to clinical improvement than those receiving placebo, among patients with symptom duration of 10 days or less (hazard ratio 1.52 [0.95–2.43]). In this study of adult patients admitted to hospital for severe COVID-19, Remdesivir was not associated with statistically significant clinical benefits. However, the numerical reduction in the time for clinical improvement in those treated earlier requires confirmation in larger studies¹⁰. At least 23 studies on Remdesivir are currently listed on various trial registers, intending to study 23,500 patients, but fewer than a quarter are double-blind, and some are uncontrolled observational studies.

CONCLUSION

Thusly the studies demonstrated that intravenous doses of Remdesivir were adequately tolerated in patients; this medication was not utilized in patients seriously infected by COVID-19, necessitating control clinical trials. Because of this, more scientific information is needed to reach conclusions about Remdesivir, its benefits, and in what situations it should be used.

Conflicts of Interest

The authors declare that there are no conflicts of interest that may have influenced this work.

Authors' Contributions

TCPA, ARVSC, MLCF, PCPA, and RNSF performed searches in the databases. TJMR, CFSR, FTB, and FWSR selected the articles that would be included in the research. FWSR corrected the writing in English. All authors performed the other parts of the research in an equal way. All authors have reviewed and approved the final text of this article and are responsible for its content.

RESUMO

O agente etiológico da COVID-19, que causa doenças respiratórias graves, como pneumonia e insuficiência pulmonar, foi confirmado como um novo coronavírus, agora conhecido como coronavírus de síndrome respiratória aguda grave 2 (SARS-CoV-2). Não existem atualmente medicamentos autorizados para o tratamento de COVID-19, nem estão também autorizadas quaisquer vacinas. Assim, o estudo teve como objetivo realizar uma revisão sobre a utilização de Remdesivir em pacientes com COVID-19. As seguintes bases de dados eletrônicas foram utilizadas MEDLINE, SCIELO, LILACS e PUBMED. Em primeiro de maio, o Remdesivir recebeu autorização de uso de emergência da Food and Drug Administration. Remdesivir é presentemente a molécula promissora no tratamento da COVID-19 tendo em conta o seu largo espectro antiviral (considerando as sequências genéticas do vírus, é expectável que mantenha atividade contra o SARS-CoV-2). A informação in vitro e in vivo está disponível para os coronavírus, assim como a extensiva base de dados de

segurança clínica (proveniente de ensaio clínico do vírus Ebola e no contexto do Monitored Emergency Use of Unregistered and Investigational Interventions - MEURI). A realização de novos estudos torna-se relevantes uma vez que os dados disponíveis são limitados sobre eficácia e segurança do Remdesivir contra SARS-nCoV-2.

PALAVRAS-CHAVE: Betacoronavirus. Infecções por Coronavirus/terapia. Coronavírus.

REFERENCES

1. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *Nature*. 2020;579:270-3.
2. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727-33.
3. World Health Organization. Coronavirus disease 2019 (COVID-19): situation report – 52. Geneva: World Health Organization; 2020. [cited 2020 Mar 12]. Available from: https://www.who.int/docs/default-source/coronavirus/20200312-sitrep-52-covid-19.pdf?sfvrsn=e2bfc9c0_2
4. BRASIL. Ministério da Saúde. Painel coronavirus. [cited 2020 May 6]. Available from: <https://covid.saude.gov.br/>
5. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-13.
6. Food and Drug Administration. FDA approves emergency use of remdesivir for COVID-19. [cited 2020 May 6]. Available from: <https://www.drugs.com/news/fda-approves-emergency-remdesivir-covid-19-89997.html>
7. Sheahan TP, Sims AC, Graham RL, Menachery VD, Gralinski LE, Case JB, et al. Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic coronaviruses. *Sci Transl Med*. 2017;9(396):eaal3653.
8. Mulangu S, Dodd LE, Davey Jr RT, Mbaya OT, Proschan M, Mukadi D, et al; PALM Consortium Study Team. A randomized, controlled trial of ebola virus disease therapeutics. *N Engl J Med*. 2019;381(24):2293-303.
9. Grein J, Ohmagari N, Shin D, Diaz G, Asperges E, Castagna A, et al. Compassionate use of remdesivir for patients with severe Covid-19. *N Engl J Med*. 2020;NEJMoa2007016. doi: 10.1056/NEJMoa2007016.
10. Wang Y, Zhang D, Du G, Du R, Zhao J, Jin Y, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet*. 2020;395(10236):1569-78.
11. Warren TK, Jordan R, Lo MK, Ray AS, Mackman RL, Soloveva V, et al. Therapeutic efficacy of the small molecule GS-5734 against ebola virus in rhesus monkeys. *Nature*. 2016;531(7594):381-5.

