

An overview of invasive plants in Brazil

RAFAEL DUDEQUE ZENNI^{1,2} and SÍLVIA RENATE ZILLER¹

(received: June 22, 2010; accepted: August 17, 2011)

ABSTRACT – (An overview of invasive plants in Brazil). Alien plants are known to occur in Brazil since the 18th century when African grasses started to be recorded in pastures near Rio de Janeiro. In the beginning of the 19th century two royal decrees (July, 1809 and July, 1810) offered grants and tax exemption to everyone who would introduce plants of economic value. Nowadays, there are 117 plant species recognized as invasive or established and with invasive potential in Brazil and an unknown number of introduced plant species. Some of the most pervasive invasive species are *Artocarpus heterophyllus* Lam. and *Hedychium coronarium* König in tropical ombrophilous forest, *Hovenia dulcis* Thunb. in subtropical ombrophilous forest and subtropical semi-deciduous forest, *Pinus taeda* L. and *Pinus elliottii* Engelm. in subtropical ombrophilous forest and steppe, *Prosopis juliflora* (Sw.) DC. in stepic-savanna, *Tecoma stans* (L.) Juss. ex Kunth in tropical and subtropical semi-deciduous forest, *Melinis minutiflora* P. Beauv. in the Brazilian savannas, and *Eragrostis plana* Nees in the steppe. The purpose of this article is to fill a knowledge gap on alien species that are invasive in Brazil and where they are invading by summarizing data obtained by joint efforts of the Hórus Institute for Environmental Conservation and Development, The Nature Conservancy (TNC), the Inter-American Biodiversity Information Network (IABIN) invasive species thematic network (I3N), and the Brazilian Ministry of Environment (MMA) in the last six years.

Key words - biological invasions, database, invasive alien plants

RESUMO – (Visão geral das plantas exóticas invasoras no Brasil). Alertas sobre espécies exóticas existem no Brasil desde o século XVIII, quando gramíneas africanas começaram a ser registradas em pastagens próximas ao Rio de Janeiro. No início do século XIX dois decretos reais (em julho de 1809 e julho de 1810) ofereciam bônus e isenção de impostos para todos que introduzissem plantas de valor econômico. Atualmente, há 117 espécies de plantas exóticas reconhecidas como estabelecidas com potencial invasor ou invasoras no Brasil, e um número desconhecido de plantas introduzidas. As plantas exóticas invasoras mais relevantes são *Artocarpus heterophyllus* Lam. e *Hedychium coronarium* König na floresta ombrófila densa, *Hovenia dulcis* Thunb. na floresta ombrófila mista e floresta estacional semidecidual do rio Paraná, *Pinus taeda* L. e *Pinus elliottii* Engelm. na floresta ombrófila mista e na estepe, *Prosopis juliflora* (Sw.) DC. na savana-estépica, *Tecoma stans* (L.) Juss. ex Kunth nas florestas estacionais semidecíduais e decíduais, *Melinis minutiflora* P. Beauv. na savana e *Eragrostis plana* Nees na estepe. Este artigo tem como objetivo ajudar a preencher uma lacuna de informação sobre espécies exóticas invasoras no Brasil e os locais onde invadem, apresentando um resumo dos dados obtidos no banco de dados de espécies exóticas invasoras no Brasil criado e mantido nos últimos seis anos por um esforço conjunto do Instituto Hórus de Desenvolvimento e Conservação Ambiental, da The Nature Conservancy (TNC), da Rede Interamericana de Informações sobre a Biodiversidade (IABIN) por meio da rede temática de Espécies Invasoras (I3N) e do Ministério de Meio Ambiente do Brasil (MMA).

Palavras-chave - banco de dados, invasões biológicas, plantas exóticas invasoras

Introduction

Invasive alien species are one of the most severe threats to the conservation of native species, communities and ecosystems (Vitousek *et al.* 1996, Wilcove *et al.* 1998, Traveset & Richardson 2006) and require urgent attention in many parts of the world (Mack *et al.* 2000). However, when data is lacking on which species are invading and where they are invading conservation planning strategies and public awareness are limited and ineffective. A critical first step to raise awareness and increase conservation efficiency is an invasive species

assessment based on existing information (Wittenberg & Cock 2001).

Brazil has the highest plant diversity in the world with two biodiversity hotspots recognized for conservation priorities: *Cerrado* (Savanna) and Atlantic Forest (Myers *et al.* 2000). More than half of the country (*ca.* 8.5 million km² was originally covered by Amazon Forest, the world's largest tropical forest, while the other half was covered by Atlantic forests, steppes, savannas, wetlands and semi-arid ecosystems (Veloso *et al.* 1992). However, little is known about the threats posed by invasive alien species to biodiversity and natural resources within these ecosystems.

The earliest records of an invasive alien plant species in Brazil is of African grasses in pastures near Rio de Janeiro (South-Eastern Brazil) from the early 18th

1. Instituto Hórus de Desenvolvimento e Conservação Ambiental, Servidão Cobra Coral, 111, 88063-513 Florianópolis, SC, Brazil.
2. Corresponding author: rafaeldz@gmail.com

century. Of these first recorded grasses several species are still invasive today such as *Urochloa maxima* (Jacq.) RD. Webster, *Hyparrhenia rufa* (Nees) Stapf, *Urochloa mutica* (Forssk.) T.Q. Nguyen, and *Melinis minutiflora* P. Beauv. (Dean 1996). Data on the source of these introductions is often lacking (*i.e.* deliberate introductions for animal forage or accidental introductions via dry ballast or other vectors), but these species are currently used as forage grasses, with some of them subjected to genetic improvements (see Cook *et al.* 2005).

Early in the 19th century two royal decrees (July, 1809 and July, 1810) offered grants and tax exemptions to everyone who introduced plants with economic value (Dean 1996). Probably due to incentives like these, the French naturalist Auguste de Saint-Hilaire recorded a great diversity of plants upon his visit to São Paulo in 1818. Most of the species he wrote about were brought from Europe but native to many different parts of the world. Among them were the currently invasive species *Sechium edule* (Jacq.) Sw., *Artocarpus heterophyllus* Lam., *Elaeis guineensis* Jacq., and *Ricinus communis* L. (Dean 1996). Saint-Hilaire also documented the spread of *Melinis minutiflora* in the *Cerrado* (Savanna) in 1824 (Lima 2002).

Currently, despite the existence of extensive number of global and regional databases and information networks on invasive species (Meyerson & Mooney 2007), information concerning invasive alien species in Brazil is largely unavailable in the scientific literature (Petenon & Pivello 2008). The purpose of this article is to fill a knowledge gap on species that are invasive in Brazil and where they are invading by summarizing data obtained by joint efforts of the Hórus Institute for Environmental Conservation and Development, The Nature Conservancy (TNC), the Inter-American Biodiversity Information Network (IABIN) invasive species thematic network (I3N), and the Brazilian Ministry of Environment (MMA) in the last six years.

Material and methods

The current article is an overview assessment of invasive plants in Brazil that attempts to summarize existing information gathered from field observations, interviews and literature reviews in the last six years. The data used for compiling a species list was gathered from the I3N Brazil (IABIN Invasive Species thematic network) database (I3N Brazil 2010). This database contains occurrence records for non-native species which are already invasive or have invasive potential in Brazil, but does not provide a full record of the habitats where the species are found. Invasive species' occurrences are organized by phytophysiognomy using the nomenclature proposed by Veloso *et al.* (1992), in Portuguese,

and by physiognomic-ecological classes following UNESCO (1973), in English. These are the physiognomies where the exotic species are considered invasive in Brazil and, on times, species also native to Brazil are considered invasive in physiognomies where the species are not known to occur historically and naturally. The data are available for public consultation and open to criticism and review. The present overview only mentions invasive species in natural habitats, and does not consider data from urban or suburban areas.

The species in the I3N database are classified as invasive based on their aptitude to establish and spread in a new habitat without direct human assistance after their introduction, while introductions are almost always likely mediated by human assistance, deliberate or accidental. The presence of a species in the database, and therefore in this overview, does not provide relationship to direct impacts or threats. We verified species for their history of invasion elsewhere when they were merely listed as established in Brazil.

Local references are related to municipalities or geographical references and to phytophysiognomy according to the Brazilian Classification System for Vegetation (Veloso *et al.* 1992). The phytophysiognomies were then matched to the physiognomic-ecological classes (UNESCO 1973) as the former is an adaptation of the later. We used GIS (Geographic Information System) techniques to merge points of occurrence in the database with corresponding ecoregions (*sensu* Olson *et al.* 2001). We chose not to directly translate the Brazilian classification system for vegetation to English because the translated names would not fully enclose the concepts behind each class and thus could potentially disconnect the information from its primary source.

We filtered the data to select only invasion and establishment records and to remove occurrences not related to municipalities, as well as records of plants in urban and suburban areas. The status of “established” refers to species which are reproducing locally, and “invasive” refers to species that are reproducing and spreading beyond their points of introduction. Although the presence of species was always confirmed by checking the source (*i.e.* a specialist, a scientific publication, or a list of plants), we also used field observations, interviews and literature reviews from the last six years to compile a series of informative case studies that collect most of the knowledge available on the impacts of invasive alien plants in Brazil.

Results

The alien plant species known to invade natural habitats in Brazil are listed in table 1. There were 3,320 database records for “invasion” or “establishment” (figure 1) of 117 different plant species (table 1) in the I3N Brazil database as of December, 2010.

The invasive plants with the greatest number of occurrences recorded are *Urochloa decumbens* (265 records), *Ricinus communis* (253 records), *Tecoma*

Table 1. Invasive alien plants records for the physiognomic-ecological classes (UNESCO 1973) and ecoregion (Olson *et al.* 2001) in Brazil. Data is from the I3N Brazil database at www.institutohorus.org.br. We gathered location references in the database from field observations, interviews and literature reviews from 2003 until 2008. Locations are mostly related to municipalities or geographical reference points. Phytophysiognomies with no correspondence in the original physiognomic-ecological classes are marked with *.

Family/Species	Physiognomic-ecological class	Ecoregion
ACANTHACEAE		
<i>Thunbergia alata</i> Bojer ex Sims	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Bahia coastal forests, Alto Paraná Atlantic forests, Serra do Mar coastal forests
<i>Thunbergia grandiflora</i> Roxb.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Bahia coastal forests, Serra do Mar coastal forests
AGAVACEAE		
<i>Agave sisalana</i> Perrine ex Engelm.	Savanna (or Brazilian savanna), Steppic-savanna*	Cerrado, Caatinga, Pernambuco interior forests
ANACARDIACEAE		
<i>Mangifera indica</i> L.	Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland, Tropical ombrophilous forest, Mangrove forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Caatinga, Bahia coastal forests, Alto Paraná Atlantic forests, Pernambuco interior forests, Southern Atlantic mangroves, Serra do Mar coastal forests, Dry Chaco, Cerrado
APIACEAE		
<i>Ammi majus</i> L.	Steppe	Uruguayan savanna, Alto Paraná Atlantic forests
<i>Centella asiatica</i> (L.) Urb.	Tropical and subtropical semi-deciduous forest, Subtropical ombrophilous forest	Bahia coastal forests, Cerrado
ARECACEAE		
<i>Archontophoenix cunninghamiana</i> H. Wendl. & Drude	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Alto Paraná Atlantic forests, Serra do Mar coastal forests
<i>Elaeis guineensis</i> Jacq.	Evergreen broad-leaved woodland, Tropical ombrophilous forest, Mangrove forest, Salt meadow	Southwest Amazon moist forests, Purus-Madeira moist forests, Pernambuco interior forests, Bahia coastal forests, Southern Atlantic mangroves, Caatinga
<i>Livistona chinensis</i> (Jacq.) R. Br. ex Mart.	Tropical ombrophilous forest	Bahia coastal forests
ASCLEPIADACEAE		
<i>Calotropis gigantea</i> (L.) W. T. Aiton	Salt meadow, Steppic-savanna*	Caatinga
<i>Calotropis procera</i> (Aiton) W. T. Aiton	Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Caatinga, Bahia coastal forests, Atlantic dry forests, Dry Chaco, Southern Atlantic mangroves
<i>Cryptostegia grandiflora</i> R. Br.	Steppic-savanna*	Caatinga
ASTERACEAE		
<i>Chrysanthemum myconis</i> L.	Steppic-savanna*	Caatinga
<i>Cirsium vulgare</i> (Savi) Ten.	Steppe, Tropical ombrophilous forest, Subtropical ombrophilous forest	Uruguayan savanna, Alto Paraná Atlantic forests, Araucaria moist forests
<i>Senecio madagascariensis</i> Poir.	Steppe	Uruguayan savanna
BALSAMINACEAE		
<i>Impatiens walleriana</i> Hook. f.	Steppe, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Savanna (or Brazilian savanna)	Cerrado, Araucaria moist forests, Bahia interior forests, Bahia coastal forests, Alto Paraná Atlantic forests, Serra do Mar coastal forests

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
BIGNONIACEAE		
<i>Spathodea campanulata</i> P. Beauv.	Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Alto Paraná Atlantic forests, Bahia coastal forests, Serra do Mar coastal forests, Araucaria moist forests
<i>Tecoma stans</i> (L.) Juss. ex Kunth	Ecotone Steppe – Deciduous or semi-deciduous forest, Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna)	Uruguayan savanna, Alto Paraná Atlantic forests, Cerrado, Araucaria moist forests, Bahia coastal forests, Serra do Mar coastal forests, Atlantic Coast restingas, Bahia interior forests
CACTACEAE		
<i>Opuntia ficus-indica</i> (L.) Mill.	Tropical and subtropical semi-deciduous forest, Salt meadow, Steppic-savanna*	Caatinga, Bahia coastal forests
CAMPANULACEAE		
<i>Hippobroma longiflora</i> (L.) G. Don	Tropical ombrophilous forest	Araucaria moist forests
CAPRIFOLIACEAE		
<i>Lonicera japonica</i> Thunb. ex Murray	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna)	Bahia interior forests, Araucaria moist forests, Uruguayan savanna
CASUARINACEAE		
<i>Casuarina equisetifolia</i> L.	Steppe, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Araucaria moist forests, Southern Atlantic mangroves, Pernambuco coastal forests, Bahia coastal forests, Serra do Mar coastal forests, Atlantic dry forests, Caatinga, Bahia interior forests
COMBRETACEAE		
<i>Terminalia catappa</i> L.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Mangrove forest, Salt meadow	Bahia coastal forests, Bahia interior forests, Serra do Mar coastal forests
COMMELINACEAE		
<i>Tradescantia zebrina</i> Heynh.	Steppe, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest	Cerrado, Bahia coastal forests, Serra do Mar coastal forests, Araucaria moist forests
CUCURBITACEAE		
<i>Sechium edule</i> (Jacq.) Sw.	Tropical ombrophilous forest	Alto Paraná Atlantic forests, Araucaria moist forests, Serra do Mar coastal forests
CUPRESSACEAE		
<i>Cupressus lusitanica</i> Mill.	Steppe	Cerrado
CYPERACEAE		
<i>Cyperus rotundus</i> L.	Tropical and subtropical semi-deciduous forest, Salt meadow, Alpine meadows, Steppic-savanna*	Bahia interior forests, Bahia coastal forests, Araucaria moist forests, Caatinga, Pernambuco interior forests, Pernambuco coastal forests
<i>Scleria mitis</i> P.J. Bergius	Tropical ombrophilous forest	Bahia coastal forests

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
EUPHORBIACEAE		
<i>Aleurites moluccanus</i> (L.) Willd.	Steppic-savanna*	Caatinga
<i>Euphorbia tirucalli</i> L.	Tropical ombrophilous forest	Bahia coastal forests
<i>Hura crepitans</i> L.	Tropical and subtropical semi-deciduous forest	Bahia coastal forests
<i>Ricinus communis</i> L.	Ecotone Steppe – Deciduous or semi-deciduous forest, Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Ecotone Savanna (or Brazilian savanna) – Savanna (or Brazilian savanna) Estépica, Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Mangrove forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Uruguayan savanna, Araucaria moist forests, Caatinga, Cerrado, Alto Paraná Atlantic forests, Atlantic Coast restingas, Bahia interior forests, Southern Atlantic mangroves, Bahia coastal forests, Serra do Mar coastal forests, Pernambuco interior forests, Atlantic dry forests, Dry Chaco
FABACEAE		
<i>Acacia auriculiformis</i> A. Cunn. ex Benth.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow	Bahia coastal forests, Bahia interior forests, Serra do Mar coastal forests
<i>Acacia farnesiana</i> (L.) Willd.	Tropical and subtropical deciduous forest	Cerrado
<i>Acacia holosericea</i> A. Cunn. ex G. Don	Tropical and subtropical semi-deciduous forest, Mangrove forest	Bahia coastal forests, Bahia interior forests
<i>Acacia longifolia</i> (Andrews) Willd.	Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Ecotone Savanna (or Brazilian savanna) – Savanna (or Brazilian savanna) Estépica, Salt meadow, Steppic-savanna*	Caatinga, Serra do Mar coastal forests, Uruguayan savanna, Atlantic Coast restingas
<i>Acacia mangium</i> Willd.	Ecotone Savanna (or Brazilian savanna) – Ombrophilous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Swamp, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Guianan savanna, Japurá-Solimoes-Negro moist forests, Bahia coastal forests, Bahia interior forests, Guianan moist forests, Amazon-Orinoco-Southern Caribbean mangroves, Serra do Mar coastal forests, Caatinga
<i>Acacia mearnsii</i> De Wild.	Ecotone Steppe – Deciduous or semi-deciduous forest, Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Ecotone Savanna (or Brazilian savanna) – Subtropical ombrophilous forest, Steppe, Steppe Parque, Tropical and subtropical deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Uruguayan savanna, Araucaria moist forests, Cerrado, Alto Paraná Atlantic forests, Serra do Mar coastal forests, Caatinga
<i>Acacia podalyriifolia</i> A. Cunn. ex G. Don	Steppe, Salt meadow	Cerrado, Serra do Mar coastal forests
<i>Albizia falcata</i> (L.) Backer ex Merr.	Tropical ombrophilous forest, Mangrove forest	Bahia coastal forests

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
<i>Clitoria fairchildiana</i> R.A. Howard	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Bahia coastal forests
<i>Crotalaria juncea</i> L.	Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland	Bahia coastal forests, Purus-Madeira moist forests, Southwest Amazon moist forests
<i>Crotalaria spectabilis</i> Roth	Evergreen broad-leaved woodland	Purus-Madeira moist forests, Southwest Amazon moist forests
<i>Leucaena leucocephala</i> (Lam.) de Wit	Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Mangrove forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Alto Paraná Atlantic forests, Caatinga, Bahia coastal forests, Pernambuco interior forests, Cerrado, Bahia interior forests, Serra do Mar coastal forests, Amazon-Orinoco-Southern Caribbean mangroves, Southern Atlantic mangroves, Pernambuco coastal forests, Guianan savanna, Atlantic dry forests, Araucaria moist forests, Dry Chaco, Pantanal, Maranhão Babaçu forests
<i>Parkinsonia aculeata</i> L.	Steppic-savanna*	Caatinga, Pernambuco interior forests
<i>Prosopis juliflora</i> (Sw.) DC.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Savanna (or Brazilian savanna), Steppic-savanna*	Caatinga, Southern Atlantic mangroves, Cerrado
<i>Pueraria phaseoloides</i> (Roxb.) Benth.	Steppe, Evergreen broad-leaved woodland, Tropical ombrophilous forest	Uruguayan savanna, Southwest Amazon moist forests, Japurá-Solimoes-Negro moist forests, Uatuma-Trombetas moist forests, Guianan moist forests
<i>Ulex europaeus</i> L.	Ecotone Ombrophilous forest – Deciduous or semi-deciduous forest, Steppe, Subtropical ombrophilous forest, Salt meadow	Alto Paraná Atlantic forests, Araucaria moist forests, Cerrado, Uruguayan savanna, Atlantic Coast restingas
IRIDACEAE		
<i>Crocasmia × crocosmiiflora</i> (Lemoine) N.E. Br.	Tropical and subtropical semi-deciduous forest, Subtropical ombrophilous forest, Savanna (or Brazilian savanna)	Bahia interior forests, Bahia coastal forests, Alto Paraná Atlantic forests, Araucaria moist forests
LAURACEAE		
<i>Persea americana</i> Mill.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Bahia coastal forests
LILIACEAE		
<i>Asparagus setaceus</i> (Kunth) Jessop	Steppe, Tropical and subtropical semi-deciduous forest, Salt meadow	Uruguayan savanna
<i>Curculigo capitulata</i> (Lour.) Kuntze	Tropical ombrophilous forest	Serra do Mar coastal forests
<i>Dracaena fragrans</i> (L.) Ker Gawl.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow	Bahia coastal forests, Serra do Mar coastal forests, Araucaria moist forests
<i>Ophiopogon japonicus</i> (L. f.) Ker Gawl.	Steppe, Subtropical ombrophilous forest	Araucaria moist forests
<i>Sansevieria trifasciata</i> Prain	Tropical ombrophilous forest, Salt meadow	Bahia coastal forests
LOMARIOPSISIDACEAE		
<i>Nephrolepis cordifolia</i> (L.) C. Presl	Steppe	Cerrado
MALVACEAE		
<i>Sterculia foetida</i> L.	Tropical and subtropical semi-deciduous forest	Bahia coastal forests, Bahia interior forests
<i>Thespesia populnea</i> (L.) Sol. ex Corrêa	Mangrove forest	Southern Atlantic mangroves

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
<i>Urena lobata</i> L.	Evergreen broad-leaved woodland, Tropical ombrophilous forest, Savanna (or Brazilian savanna)	Madeira-Tapajós moist forests, Serra do Mar coastal forests, Cerrado
MELIACEAE		
<i>Azadirachta indica</i> A. Juss.	Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Ecotone Savanna (or Brazilian savanna) – FP de Influência Mar, Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland, Tropical ombrophilous forest, Mangrove forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Caatinga, Pernambuco interior forests, Pernambuco coastal forests, Bahia coastal forests, Madeira-Tapajós moist forests, Southwest Amazon moist forests, Southern Atlantic mangroves, Maranhão Babaçu forests, Atlantic dry forests, Atlantic Coast restingas
<i>Melia azedarach</i> L.	Ecotone Savanna (or Brazilian savanna) – Ombrophilous forest, Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Savanna (or Brazilian savanna)	Chiquitano dry forests, Cerrado, Araucaria moist forests, Alto Paraná Atlantic forests, Serra do Mar coastal forests
MORACEAE		
<i>Artocarpus heterophyllus</i> Lam.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Bahia coastal forests, Southern Atlantic mangroves, Bahia interior forests, Serra do Mar coastal forests, Araucaria moist forests, Caatinga
<i>Morus nigra</i> L.	Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Mangrove forest, Savanna (or Brazilian savanna)	Araucaria moist forests, Serra do Mar coastal forests, Alto Paraná Atlantic forests, Bahia coastal forests
MUSACEAE		
<i>Musa ornata</i> Roxb.	Tropical ombrophilous forest, Steppic-savanna*	Araucaria moist forests, Caatinga
<i>Musa rosacea</i> Jacq.	Tropical ombrophilous forest	Araucaria moist forests
MYRTACEAE		
<i>Eucalyptus robusta</i> Sm.	Campinarana*, Savanna (or Brazilian savanna), Steppic-savanna*	Cerrado, Atlantic Coast restingas
<i>Eugenia malaccensis</i> L. ¹	Tropical ombrophilous forest	Bahia coastal forests
<i>Psidium guajava</i> L.	Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Ecotone Savanna (or Brazilian savanna) – Ombrophilous forest, Steppe, Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna)	Araucaria moist forests, Alto Paraná Atlantic forests, Serra do Mar coastal forests, Cerrado, Bahia coastal forests, Southwest Amazon moist forests, Pernambuco interior forests, Bahia interior forests
<i>Syzygium cumini</i> (L.) Skeels	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Bahia coastal forests, Alto Paraná Atlantic forests, Bahia interior forests, Araucaria moist forests, Cerrado, Atlantic dry forests

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
OLEACEAE		
<i>Ligustrum japonicum</i> Thunb.	Steppe, Subtropical ombrophilous forest	Cerrado, Araucaria moist forests
<i>Ligustrum lucidum</i> W. T. Aiton	Steppe, Tropical and subtropical semi-deciduous forest, Subtropical ombrophilous forest	Cerrado, Alto Paraná Atlantic forests, Araucaria moist forests
<i>Ligustrum vulgare</i> L.	Subtropical ombrophilous forest	Araucaria moist forests
PINACEAE		
<i>Pinus caribaea</i> Morelet	Ecotone Savanna (or Brazilian savanna) – Ombrophilous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow	Chiquitano dry forests, Araucaria moist forests, Tocantins/Pindare moist forests, Cerrado, Serra do Mar coastal forests
<i>Pinus elliottii</i> Engelm.	Ecotone Steppe – Deciduous or semi-deciduous forest, Ecotone Ombrophilous forest – Deciduous or semi-deciduous forest, Ecotone Savanna (or Brazilian savanna) – Ombrophilous forest, Steppe, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Swamp, Salt meadow, Alpine meadows, Savanna (or Brazilian savanna)	Uruguayan savanna, Alto Paraná Atlantic forests, Cerrado, Araucaria moist forests, Serra do Mar coastal forests, Bahia coastal forests, Atlantic Coast restingas
<i>Pinus oocarpa</i> Schiede ex Schldtl.	Tropical and subtropical semi-deciduous forest, Subtropical ombrophilous forest, Savanna (or Brazilian savanna)	Alto Paraná Atlantic forests, Cerrado, Araucaria moist forests
<i>Pinus patula</i> Schldtl. & Cham.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Cerrado, Araucaria moist forests
<i>Pinus taeda</i> L.	Steppe, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Swamp, Salt meadow, Alpine meadows, Savanna (or Brazilian savanna)	Cerrado, Araucaria moist forests, Alto Paraná Atlantic forests, Uruguayan savanna, Serra do Mar coastal forests, Atlantic Coast restingas
PITTOSPORACEAE		
<i>Pittosporum undulatum</i> Vent.	Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Tropical and subtropical semi-deciduous forest, Subtropical ombrophilous forest	Araucaria moist forests, Uruguayan savanna
POACEAE		
<i>Andropogon gayanus</i> Kunth	Tropical and subtropical semi-deciduous forest, Savanna (or Brazilian savanna), Steppic-savanna*	Bahia coastal forests, Caatinga
<i>Arundo donax</i> L.	Savanna (or Brazilian savanna)	Cerrado
<i>Bambusa textilis</i> McClure	Tropical and subtropical semi-deciduous forest	Alto Paraná Atlantic forests
<i>Bambusa vulgaris</i> Schrad. ex J. C. Wendl.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Southern Atlantic mangroves, Pernambuco coastal forests, Bahia coastal forests, Serra do Mar coastal forests, Araucaria moist forests, Caatinga
<i>Cenchrus ciliaris</i> L.	Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Tropical and subtropical semi-deciduous forest, Steppic-savanna*	Caatinga, Pernambuco interior forests, Pernambuco coastal forests, Southern Atlantic mangroves, Maranhão Babaçu forests

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
<i>Cortaderia selloana</i> (Schult. & Schult. f.) Asch. & Graebn.	Tropical ombrophilous forest, Subtropical ombrophilous forest	Araucaria moist forests
<i>Cynodon dactylon</i> (L.) Pers.	Tropical ombrophilous forest, Steppic-savanna*	Bahia coastal forests, Serra do Mar coastal forests, Caatinga
<i>Digitaria decumbens</i> Stent	Tropical and subtropical semi-deciduous forest, Steppic-savanna*	Caatinga, Atlantic Coast restingas
<i>Echinochloa crus-galli</i> (L.) P. Beauv.	Swamp	Cerrado, Caatinga
<i>Eragrostis plana</i> Nees	Steppe, Tropical ombrophilous forest, Subtropical ombrophilous forest, Salt meadow	Cerrado, Uruguayan savanna, Alto Paraná Atlantic forests, Bahia coastal forests, Araucaria moist forests, Bahia interior forests
<i>Hyparrhenia rufa</i> (Nees) Stapf	Tropical and subtropical deciduous forest, Savanna (or Brazilian savanna)	Caatinga, Alto Paraná Atlantic forests, Cerrado, Pantanal
<i>Melinis minutiflora</i> P. Beauv.	Ecotone Ombrophilous forest – Deciduous or semi-deciduous forest, Ecotone Savanna (or Brazilian savanna) – Deciduous or semi-deciduous forest, Campinarana*, Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Mangrove forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Xingu-Tocantins-Araguaia moist forests, Alto Paraná Atlantic forests, Bahia interior forests, Cerrado, Caatinga, Bahia coastal forests, Serra do Mar coastal forests, Araucaria moist forests, Pantanal, Dry Chaco
<i>Melinis repens</i> (Willd.) Zizka	Steppe, Tropical and subtropical deciduous forest, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Araucaria moist forests, Alto Paraná Atlantic forests, Serra do Mar coastal forests, Cerrado, Dry Chaco
<i>Pennisetum clandestinum</i> Hochst. ex Chiov.	Tropical ombrophilous forest, Salt meadow	Bahia coastal forests
<i>Pennisetum purpureum</i> Schumach.	Steppe, Tropical ombrophilous forest, Subtropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Cerrado, Bahia coastal forests, Araucaria moist forests, Serra do Mar coastal forests, Pantanal, Caatinga
<i>Urochloa brizantha</i> (Hochst. ex A. Rich.) R. D. Webster	Ecotone Steppe – Deciduous or semi-deciduous forest, Steppe, Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna)	Alto Paraná Atlantic forests, Cerrado, Uruguayan savanna, Pernambuco coastal forests, Pernambuco interior forests, Madeira-Tapajós moist forests, Japurá-Solimoes-Negro moist forests, Maranhão Babaçu forests, Tocantins/Pindare moist forests, Atlantic Coast restingas, Caatinga, Purus-Madeira moist forests, Guianan savanna, Atlantic dry forests
<i>Urochloa decumbens</i> (Stapf) R. D. Webster	Steppe, Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna)	Cerrado, Bahia interior forests, Madeira-Tapajós moist forests, Southwest Amazon moist forests, Purus-Madeira moist forests, Japurá-Solimoes-Negro moist forests, Bahia coastal forests, Araucaria moist forests, Caatinga, Pernambuco coastal forests, Pernambuco interior forests, Southern Atlantic mangroves, Mato Grosso seasonal forests, Alto Paraná Atlantic forests, Pantanal

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
<i>Urochloa humidicola</i> (Rendle) Morrone & Zuloaga	Campinarana*, Steppe, Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland, Tropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Pantanal, Cerrado, Pernambuco coastal forests, Pernambuco interior forests, Purus-Madeira moist forests, Chiquitano dry forests, Madeira-Tapajós moist forests, Southwest Amazon moist forests, Japurá-Solimoes-Negro moist forests, Mato Grosso seasonal forests, Uatuma-Trombetas moist forests, Amazon-Orinoco-Southern Caribbean mangroves, Guianan savanna, Maranhão Babaçu forests, Tocantins/Pindare moist forests, Araucaria moist forests, Bahia coastal forests, Caatinga, Southern Atlantic mangroves, Atlantic dry forests, Atlantic Coast restingas
<i>Urochloa maxima</i> (Jacq.) R. D. Webster	Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest, Salt meadow, Savanna (or Brazilian savanna), Steppic-savanna*	Cerrado, Caatinga, Dry Chaco, Bahia interior forests, Bahia coastal forests, Serra do Mar coastal forests, Araucaria moist forests, Tocantins/Pindare moist forests, Chiquitano dry forests, Alto Paraná Atlantic forests, Pantanal
<i>Urochloa mutica</i> (Forssk.) T. Q. Nguyen	Tropical ombrophilous forest, Mangrove forest, Salt meadow	Araucaria moist forests
<i>Urochloa plantaginea</i> (Link) R. D. Webster	Salt meadow	Bahia coastal forests
<i>Urochloa ruziziensis</i> (R. Germ. & C. M. Evrard) Crins	Tropical ombrophilous forest, Swamp	Araucaria moist forests
<i>Urochloa stolonifera</i> (Gooss.) Chippindale	Steppic-savanna*	Caatinga
<i>Urochloa subquadripara</i> (Trin.) R. D. Webster	Campinarana*, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Swamp, Mangrove forest, Salt meadow, Savanna (or Brazilian savanna)	Pantanal, Bahia coastal forests, Cerrado, Araucaria moist forests, Guianan moist forests, Uruguayan savanna, Serra do Mar coastal forests, Atlantic Coast restingas, Alto Paraná Atlantic forests
PROTEACEAE		
<i>Grevillea banksii</i> R. Br.	Salt meadow	Bahia coastal forests
<i>Grevillea robusta</i> A. Cunn. ex R. Br.	Steppe, Tropical ombrophilous forest, Swamp, Salt meadow	Cerrado, Bahia coastal forests, Alto Paraná Atlantic forests
PTERIDACEAE		
<i>Pteris vittata</i> L.	Steppe	Cerrado
RHAMNACEAE		
<i>Hovenia dulcis</i> Thunb.	Ecotone Steppe – Deciduous or semi-deciduous forest, Ecotone Ombrophilous forest – Deciduous or semi-deciduous forest, Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest	Uruguayan savanna, Alto Paraná Atlantic forests, Araucaria moist forests, Cerrado, Bahia interior forests, Serra do Mar coastal forests

continue

continuation

Family/Species	Physiognomic-ecological class	Ecoregion
ROSACEAE		
<i>Eriobotrya japonica</i> (Thunb.) Lindl.	Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Subtropical ombrophilous forest	Araucaria moist forests, Cerrado, Alto Paraná Atlantic forests, Bahia coastal forests, Bahia interior forests, Serra do Mar coastal forests
RUBIACEAE		
<i>Coffea arabica</i> L.	Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Steppic-savanna*	Alto Paraná Atlantic forests, Araucaria moist forests, Caatinga
RUTACEAE		
<i>Citrus aurantium</i> L.	Steppe, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Araucaria moist forests, Alto Paraná Atlantic forests, Bahia coastal forests
<i>Citrus limon</i> (L.) Osbeck	Ecotone Ombrophilous forest – Deciduous or semi-deciduous forest, Steppe, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest, Salt meadow	Cerrado, Alto Paraná Atlantic forests, Araucaria moist forests, Bahia coastal forests, Caatinga, Uruguayan savanna
THELYPTERIDACEAE		
<i>Macrothelypteris torresiana</i> (Gaudich.) Ching	Steppe	Cerrado
<i>Thelypteris dentata</i> (Forssk.) E. P. St. John	Steppe	Cerrado
WOODSIACEAE		
<i>Deparia petersenii</i> (Kunze) M. Kato	Steppe	Cerrado
ZINGIBERACEAE		
<i>Hedychium coccineum</i> Buch.-Ham. ex Sm.	Ecotone Ombrophilous forest – Deciduous or semi-deciduous forest, Tropical and subtropical semi-deciduous forest, Tropical ombrophilous forest	Araucaria moist forests, Bahia interior forests, Alto Paraná Atlantic forests
<i>Hedychium coronarium</i> J. König	Ecotone Ombrophilous forest – Deciduous or semi-deciduous forest, Ecotone Tropical ombrophilous forest – Subtropical ombrophilous forest, Ecotone Savanna (or Brazilian savanna) – Ombrophilous forest, Steppe, Tropical and subtropical deciduous forest, Tropical and subtropical semi-deciduous forest, Evergreen broad-leaved woodland, Tropical ombrophilous forest, Subtropical ombrophilous forest, Swamp, Salt meadow, Alpine meadows	Cerrado, Serra do Mar coastal forests, Araucaria moist forests, Bahia interior forests, Caatinga, Alto Paraná Atlantic forests, Bahia coastal forests
<i>Hedychium gardnerianum</i> Sheppard ex Ker Gawl.	Subtropical ombrophilous forest	Araucaria moist forests

¹ Registered in the database as *Syzygium malaccense* (L.) Merr. & L.M. Perry

stans (239 records), *Urochloa humidicola* (225 records), *Azadirachta indica* (220 records), *Prosopis juliflora* (212 records), *Cenchrus ciliaris* (194 records), *Pinus elliottii* (165 records), *Leucaena leucocephala* (155 records), *Pinus taeda* (128 records), and *Hovenia dulcis* (126 records).

However, the invasive plant species covering the largest number of phytophysiological types are *Ricinus communis*, *Hedychium coronarium*, *Melinis minutiflora* (all present in 12 phytophysiological types), *Pinus elliottii* (present in 11), *Acacia mearnsii* (present in 10), *Tecoma*




Figure 1. Localities with at least one occurrence record for invasive plants (gray dots) registered in the I3N-Brazil database as of December 2010. Most localities contain only one record for a single species, but some have many. There is likely a strong sampling bias towards the southern and northeastern regions, while a clear lack of information for central and northern regions. Axes are latitude (vertical) and longitude (horizontal).

stans, and *Azadirachta indica* (both present in 9). Species also worth mentioning due to their relevance as invasives, but present in fewer phytophysionomies are *Artocarpus heterophyllus* in tropical ombrophilous forest, *Hovenia dulcis* in subtropical ombrophilous forest, *Prosopis juliflora* in Steppic-savanna, and *Eragrostis plana* in Steppe.

Phytophysionomies with the highest number of invasive alien plants are, in descending order: tropical ombrophilous forests (69 invasive alien species recorded), tropical and subtropical semi-deciduous forests (60), herbaceous and half-woody salt swamps (45), steppes (40), shrublands or savanna (36) and steppic-savanna (34). As would be expected (figure 1), phytophysionomies in the Amazon region have the lowest numbers of invasive alien plants recorded (13 species) and very few occurrence records (173 occurrences or ~5% of the total).

Most documented invasive alien plant species in Brazil are related to human activities and to accessibility by roads, which are greatly lacking in the Northern region. Additionally, many species were intentionally cultivated at some stage even if the original introduction

was accidental. Examples includes *Eragrostis plana*, a plant accidentally introduced but then cultivated as a forage grass, and *Cyperus rotundus*, cultivated as a medicinal plant.

In order to illustrate the status of biological invasions in Brazil, a few case studies are presented below. We think these case studies also represent a large proportion of the published scientific knowledge available on invasive alien plants for the country.

Eragrostis plana was accidentally introduced as a seed contaminant of *Chloris gayana* Kunth seed shipments in Rio Grande do Sul State (extreme south of Brazil) in the 1950s. After introduction, *E. plana* outcompeted the native species *Aristida* sp. in pastures, which was considered a serious agricultural weed and problem for cattle ranching at the time. *Eragrostis plana* was therefore considered a viable forage alternative (Reis 1993), which led to large promoting and planting of it as a crop species in Southern Brazil. Only later did ranchers notice that the species was too fibrous when mature. Cattle would either not feed on it (Reis 1993) or would not gain weight when feeding on it. The species is currently established on more than two million hectares of mostly degraded or overgrazed steppes in Rio Grande do Sul state, also occurring in smaller extent in Santa Catarina, Paraná, São Paulo, Mato Grosso, Mato Grosso do Sul, Bahia, Tocantins and Pará and in the Distrito Federal (Medeiros & Focht 2007), as well as in Uruguay. The species has reported allelopathic effects, especially on species with late germination cycles, from decomposition of plant tissues (Ferreira *et al.* 2008). It also produces higher root biomass than most native grass species and is more efficient in resource acquisition (*i.e.* water and nutrients) in soil layers between 0 and 10 cm (Abichequer *et al.* 2006). The species is neglected by cattle and thus in highly invaded steppes there are severe economic losses due to the low yield in feeding animals with this forage grass (Marcantonio 2002). Seeds remain viable in the soil more than 24 years when located 20 cm below the soil surface (Medeiros & Focht 2007), potentially making the eradication of the species in invaded sites highly unlikely.

Melinis minutiflora was first recorded in Brazil in 1824, after spreading into deforested areas following several human induced fires (Lima 2002). Nowadays this grass is abundant in many areas, especially in the Cerrado (Pivello *et al.* 1999). The spread of *M. minutiflora* is correlated to a decrease in native species richness in the invaded sites (in experimental plots in the Brasilia National Park the number of native species decreased from 54 in 1997 to 20 in 2001). This has been

attributed to a positive feedback where *M. minutiflora* increases the fuel loads which increase fire temperatures. When seasonal fires reach these higher temperatures they eliminate the seed bank of most species and allow the invasive grass to dominate the burnt areas (Hoffman *et al.* 2004, Martins *et al.* 2004). Moreover, the grass reduces the regeneration of native plants, as shown for *Cecropia pachystachya* Trécul in riparian areas (Morosini & Klink 1997).

Native to Mexico and Southern United States, *Tecoma stans* was introduced in Brazil for landscaping around 1871 and now occupies more than 10,000 hectares of degraded Alto Paraná Atlantic Forests (Bredow *et al.* 2004). It forms dense thickets and resprouts vigorously. The loss of pastures and agricultural areas to invasion induces farmers to abandon areas and generates an annual loss of US\$ 7,500,000 for the cattle industry (Pedrosa-Macedo 2004). It is forbidden to use, transport or commercialize the species since 1997 in Paraná State (Resolução Estadual nº 151/1997).

Prosopis juliflora is a legume tree native to Central America (Palacios 2006) and was introduced as a forage alternative in the 1940's in the *Caatinga* region, Northeastern Brazil (Pegado *et al.* 2006). The Brazilian Federal government promoted the distribution of seedlings of *P. juliflora* in the 1980's, which led to populations scattered across the northeastern states and the *Caatinga* biome. Some reports note that invaded areas have around 87% cover of *P. juliflora*, drastically changing the diversity and structure of native forests (Pegado *et al.* 2006). In invaded riparian areas in Bahia the species represented 60.8% of the forest regeneration (1,255 seedlings ha⁻¹), followed by another invasive alien plant *Parkinsonia aculeata* L. which represented 4.3% (89 seedlings ha⁻¹). Seedlings of other 120 adult native plants occurring in the area shared the other 34.9% of the forest regeneration (P.C.F. Lima & L.H.P. Kiill, unpublished data).

Artocarpus heterophyllus is native to high elevation locations (above 1,110 m) on the Ghats Mountains in India and the Malay Peninsula (Sahni 2000). It was one of the first alien trees introduced in the Atlantic forest region (Dean 1996). The presence of the species is usually associated with human occupation and it benefits from cultivation as a fruit tree and as an ornamental. Areas invaded by *A. heterophyllus* are found to have higher densities of native rats in comparison to natural forests without the alien plant (Bergallo *et al.* 2009). Also, the diversity of seed species dispersed by rats also decreases with the increase of *A. heterophyllus* densities (Bergallo *et al.* 2009). This pattern could

lead to a decrease of native plant abundance and to the increase of *A. heterophyllus* dominance over the long term.

Discussion

Limitations – We found that there is some ability to quantify the species and habitats invaded by reviewing plant occurrence records within the I3N Brazil database. Yet, this database has some limitations in discriminating major and minor invasive plant species within each ecoregion and habitat and across Brazil. Records of occurrences are basically presence records and contain no data on the size and density of occupied areas, or on the dynamics of invasive species at different locations. Therefore, it is not possible to assume that a larger number of records represent a larger area of occurrence. Additionally, as in many records of occurrence in plant databases, it is not possible to determine absences as they are potentially related to the lack of information available or to the presence of the species in a status different than established or invasive (*i.e.* contained, naturalized, present only in urban areas or unknown).

“Established” refers to species which are reproducing locally, and “invasive” refers to species that are reproducing and spreading beyond their points of introduction. Although this definition may be clear, it is difficult for observers to be sure about the status of invasion in the field, especially as these are often side observations done while performing other work in restoration, animal or plant surveys, or just field expeditions. The lack of expertise on biological invasions in Brazil may also lead to confusion between invasive, established or present species in given habitats. For example, it is crucial to differentiate invasions in well preserved natural habitats from invasions on degraded sites, such as road sides or pastures.

The knowledge available on invasive alien floras and plant species is highly heterogeneous in the country (figure 1), thus comparing species presence and/or invasion status across regions or any other spatial scale is not encouraged. However, historical differences among regions and differences in land use might explain some of the differences in numbers of invasive species. While some regions are more studied than others, with the south and southeast having received more attention than the north and central-west, there is also much variation in the level of ecosystem intactness across the landscape. Also, the number, diversity and intensity of pathways and vectors transporting invasive alien species varies tremendously across regions.

It is clear that a country as large as Brazil has an enormous challenge assessing its biota due to its size and diversity of ecosystems and landscapes. This overview assessment of plant invasions in Brazil is far from complete or fully accurate. However, it is an initial step towards a better understanding of biological invasions and provides much-needed information for managers and scientists working on invasion issues in Brazil.

Implications for management – We show that, based on a cursory list of invasive species occurring in Brazil, there are many species that have received little to no attention from biologists. Yet, it is clear that invasive alien plants are important components of environmental change in Brazilian ecosystems based on the trend of decreased biodiversity in invaded areas and more research and action is urgently needed to address drivers and consequences of biological invasions. This research will support the implementation of more accurate management and control strategies as well as contribute towards a better theoretical understanding of biological invasions.

Conflicts of interest between those species deemed economically important and those that may prevent the achievement of environmental conservation goals need to be addressed urgently. Both private companies and governmental agencies promote the spread of invasive species (*i.e.* *E. plana* for pastures, *P. juliflora* for cattle and human feeding, and *Elaeis guineensis* for biofuels) and currently no regulations exist to contain, prevent, or control the spread of species. It is highly desirable that such policies are accompanied by criteria to establish production areas, and that, at the example of certification in Forestry, invasive species that yield economic benefits include prevention and routine control in their management. Some legal limitations are beginning to exist thanks to official invasive species lists issued by the states of Paraná and Santa Catarina, in the south of Brazil, restricting public nurseries to produce seedlings and the use of listed species in restoration and road stabilization efforts.

The country has a fragmented legal framework for dealing with invasive alien species; there are many areas of conflict and many gaps. Existing policies are ineffective in regulating economic activities that are driving the trade of invasive species. Brazilians are beginning to better understand and register the consequences of biological invasions and several promising initiatives are now starting. States are developing strategies for invasive alien species by creating official reference lists of invasive species, controlling and managing invasive alien species in protected areas and restricting the use

of these species for certain purposes. The Ministry of Environment published a national strategy for invasive alien species in 2009 with support from a National Committee on Invasive Species involving different Ministries, environmental management and health agencies. This work needs to be supported and continued, aligned with the Convention on Biological Diversity.

It is also important to notice that most of the widely distributed invasive plants in Brazil (such as *Pinus* spp. and African grasses) are well known for their capacity to thrive under high levels of disturbance. It suggests that several invasions are a consequence of other disturbances rather than agents of disturbance themselves (see Vitousek *et al.* 1996). Therefore, to maintain the quality of many Brazil's natural ecosystems, it is crucial to prevent and control invasions by alien species.

Implications for Ecology – This study shows that numerous opportunities to study the ecological consequences of invasions in extremely rich, diverse and heterogeneous ecosystems and landscapes exist. These opportunities are currently neglected by biologists and ecologists. Until recently, alien plants were ignored in floristic surveys leading to a lack of historical records and quantitative data on the presence of invasive alien plants in reports and in herbarium and museum collections. Most botanists who contribute material to Herbaria do not bother collecting non-native plants, though these would be precious and useful records for early detection, management and for understanding the spread of invasive species by providing historical records on habitat occupation and spread. For example, many species niche models rely on herbarium records to build species' potential distributions models and to predict suitable habitats for invasive species before they get there (*i.e.* Zenni *et al.* 2009).

Many recent (< 70 years) plant introductions have led to completely different colonization patterns. While some species became widespread quickly (*i.e.* *Eragrostis plana* and *Pinus taeda*) others are observed as less able to spread (*i.e.* several *Eucalyptus* species and *Nerium oleander*). The causes of these differences are open research questions, as the behavior of the species varies greatly in different climates, ecosystems and parts of the world.

Richardson *et al.* (2008) recently questioned if pine invasions in South America could become as dramatic as they are in several other Southern Hemisphere countries such as South Africa and New Zealand. We believe the answer to be “yes” from current evidence in a number of countries (Simberloff *et al.* 2010). However, more

research on patterns and drivers of pine invasions in Brazil will much contribute to a more precise answer. So far we can only speculate.

There is equal urgency for increasing research, for qualified ecologists and biologists working on the ecology of invasions, for field managers and decision-makers advancing control work, and for policy-makers supporting legal frameworks in the country and worldwide. Filling the sampling gaps (shown in figure 1) and acquiring knowledge about the dynamics and influences of invasions in different regions and habitats should be invasion biologists' first priority.

Further work – The next steps in invasion efforts should include quantifying invasions of alien plants, evaluating consequences of these invasions on the natural environment, defining priorities and implementing prevention and control strategies when relevant to biodiversity conservation. Thus, it is important that levels of invasion by alien species in different ecosystems are clearly defined as major, minor, incipient, or emerging invaders. Based on these distinctions, priorities should be set according to the potential impacts and the feasibility of eradication or control efforts. The publication of official invasive species lists is an important first step to provide support to managers and institutions to build up prevention and early detection techniques and regional strategies to mitigate the threats posed by biological invasions. Decentralizing these huge task from the federal government to state agencies and to municipalities should greatly help set priorities and understand impacts at the local level, and use these results to multiply the work across the country.

Acknowledgements – We thank everyone who contributed with data and observations during the National Survey on Invasive Alien Species and onward efforts. We thank the Brazilian Ministry of Environment, the Inter-American Biodiversity Information Network (IABIN – I3N), and The Nature Conservancy for funding different parts of this project. We thank Sara Kuebbing for improving the language, Francisco Zenni Guérios for helping with table 1, and two anonymous reviewers for important suggestions.

References

- ABICHEQUER, A.D., MEDEIROS, C.M.O. & SPANNENBERG, P.R.O. 2006. Crescimento e distribuição de raízes de capim-annoni-2: vantagem competitiva em relação ao campo nativo?. *In* Anais da 21ª Reunião do Grupo Técnico em Forrageiras do Cone Sul. Embrapa Clima Temperado, Pelotas, CD-ROM.
- BERGALLO, H.G., RAÍCES, D.S.L., UZÊDA, M.C., ROCHA, C.F.D., ABREU, R.C. & FERREIRA, P.M. 2009. Densidade de jaqueiras e seus efeitos na comunidade de pequenos mamíferos, dispersão de sementes e agregados de solo na Ilha Grande, RJ. *In* Anais do I Congresso Brasileiro sobre Bioinvasão. Universidade Federal do Maranhão, São Luís, CD-ROM.
- BREDOW, E.A., PEDROSA-MACEDO, J.H. & VITORINO, M.D. 2004. Plantas invasoras no Paraná. *In* Princípios e rudimentos do controle biológico de plantas (J.H. Pedrosa-Macedo & E.A. Bredow, eds.). Universidade Federal do Paraná, Curitiba, p.51-99.
- COOK, B.G., PENGELLY, B.C., BROWN, S.D., DONNELLY, J.L., EAGLES, D.A., FRANCO, M.A., HANSON, J., MULLEN, B.F., PARTRIDGE, I.J., PETERS, M. & SCHULTZE-KRAFT, R. 2005. Tropical forages: an interactive selection tool. <http://www.tropicalforages.info> (accessed 2009 Mar 25).
- DEAN, W. 1996. A ferro e fogo: a história e a devastação da mata atlântica brasileira. Companhia das Letras, São Paulo.
- FERREIRA, N.R., MEDEIROS, R.B. & SOARES, G.L.G. 2008. Potencial alelopático do capim-annoni-2 (*Eragrostis plana* Nees) na germinação de sementes de gramíneas perenes estivais. *Revista Brasileira de Sementes* 30:43-50.
- HOFFMANN, W.A., LUCATELLI, V.M.P.C., SILVA, F.J., AZEVEDO, I.N.C., MARINHO, M.S., ALBUQUERQUE, A.M.S., LOPES, A.O. & MOREIRA, S.P. 2004. Impact of the invasive alien grass *Melinis minutiflora* at the savanna-forest ecotone in the Brazilian Cerrado. *Diversity and Distributions* 10:99-103.
- I3N-BRASIL. 2010. Base de dados sobre espécies exóticas invasoras. <http://www.institutohorus.org.br> (accessed 2010 Dez 20).
- LIMA, M.E.A.T. 2002. As caminhadas de Auguste de Saint-Hilaire pelo Brasil e Paraguai. Editora Autêntica, Belo Horizonte.
- MACK, R.N., SIMBERLOFF, D., LONSDALE, W.M., EVANS, H., CLOUT, M. & BAZZAZ, F.A. 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* 10: 689-710.
- MARCANTONIO, G. 2002. Manjedoura cincerro. Federacite, Porto Alegre.
- MARTINS, C.R., LEITE, L.L. & HARIDASAN, M. 2004. Capim-gordura (*Melinis minutiflora* P. Beauv.), uma gramínea exótica que compromete a recuperação de áreas degradadas em unidades de conservação. *Revista Árvore* 28:739-747.
- MEDEIROS, R.B. & FOCHT, T. 2007. Invasão, prevenção, controle e utilização do capim-annoni-2 (*Eragrostis plana* Nees) no Rio Grande do Sul, Brasil. *Revista Agropecuária Gaúcha* 13:1-28.

- MEYERSON, L.A. & MOONEY, H.A. 2007. Invasive alien species in an era of globalization. *Frontiers in Ecology and the Environment* 5:199-208.
- MOROSINI, I.B.A. & KLINK, C.A. 1997. Interferência do capim-gordura (*Melinis minutiflora* Beauv.) no desenvolvimento de plântulas de embaúba (*Cecropia pachystachya*). In *Contribuição ao conhecimento ecológico do cerrado* (L.L. Leite & C.H. Saito, org.). Congresso de Ecologia do Brasil. Universidade Federal de Brasília, Brasília, CD-ROM.
- MYERS, N., MITTERMEIER, R.A., MITTERMEIER, C.G., FONSECA, G.A.B. & KENT, J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403:853-858.
- OLSON, D.M., DINERSTEIN, E., WIKRAMANAYAKE, E.D., BURGESS, N.D., POWELL, G.V.N., UNDERWOOD, E.C., D'AMICO, J.A., ITOUA, I., STRAND, H.E., MORRISON, J.C., LOUCKS, C.J., ALLNUTT, T.F., RICKETTS, T.H., KURA, Y., LAMOREUX, J.F., WETTENGEL, W.W., HEDAO, P. & KASSEM, K.R. 2001. Terrestrial ecoregions of the world: a new map of life on Earth. *Bioscience* 51: 933-938.
- PALACIOS, R.A. 2006. Los mezquites mexicanos: biodiversidad y distribución geográfica. *Boletín de la Sociedad Argentina de Botánica* 41:99-121.
- PEDROSA-MACEDO, J.H. 2004. Plantas invasoras no Paraná. In *Princípios e rudimentos do controle biológico de plantas* (J.H. Pedrosa-Macedo & E.A. Bredow, eds.). Universidade Federal do Paraná, Curitiba, p.115-128.
- PEGADO, C.M.A., ANDRADE, L.A., FÉLIX, L.P. & PEREIRA, I.M. 2006. Efeitos da invasão biológica de algaroba – *Prosopis juliflora* (Sw.) DC. sobre a composição e a estrutura do estrato arbustivo-arbóreo da caatinga no município de Monteiro, PB, Brasil. *Acta Botanica Brasilica* 20:887-898.
- PETENON, D. & PIVELLO, V.R. 2008. Plantas invasoras: representatividade da pesquisa dos países tropicais no contexto mundial. *Natureza & Conservação* 6:65-77.
- PIVELLO, V.R., SHIDA, C.N. & MEIRELLES, S.T. 1999. Alien grasses in Brazilian savannas: a threat to the biodiversity. *Biodiversity and Conservation* 8: 1281-1294.
- REIS, J.C.L. 1993. Capimannoni-2: origem, morfologia, características, disseminação. In *Reunião regional de avaliação de pesquisa comannoni-2*. Embrapa-CPPSUL, Bagé, v.1., p.5-23.
- RICHARDSON, D.M., VAN WILGEN, B.W. & NUÑEZ, M.A. 2008. Alien conifer invasions in South America: short fuse burning? *Biological Invasions* 10:573-577.
- SAHNI, K.C. 2000. *The book of Indian trees*. Oxford University Press, Delhi.
- SIMBERLOFF, D., NUÑEZ, M.A., LEDGARD, N.J., PAUCHARD, A., RICHARDSON, D.M., SARASOLA, M., VAN WILGEN, B.W., ZALBA, S.M., ZENNI, R.D., BUSTAMANTE, R., PEÑA, E. & ZILLER, S.R. 2010. Spread and impact of introduced conifers in South America: lessons from other southern hemisphere regions. *Austral Ecology* 35:489-504.
- TRAVERSE, A. & RICHARDSON, D.M. 2006. Biological invasions as disruptors of plant reproductive mutualisms. *Trends in Ecology and Evolution* 21:208-216.
- UNESCO. 1973. *International Classification and Mapping of Vegetation*. Ecology and conservation, v.6. United Nations Educational, Scientific and Cultural Organization, Paris.
- VELOSO, H.P., OLIVEIRA FILHO, L.C., VAZ, A.M.S.F., LIMA, M.P.M., MARQUETE, R. & BRAZÃO, J.E.M. 1992. *Manual técnico da vegetação brasileira*. Manuais técnicos em geociências, v.1. Fundação Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro.
- VITOUSEK, P.M., D'ANTONIO, C.M., LOOPE, L.L. & WESTBROOKS, R. 1996. Biological invasions as global environmental change. *American Scientist* 84:468-478.
- WILCOVE, D.S., ROTHSTEIN, D., DUBOW, J., PHILLIPS, A. & LOSOS, E. 1998. Quantifying threats to imperiled species in the United States. *Bioscience* 48: 607-615.
- WITTENBERG, R. & COCK, M.J.W. 2001. *Invasive alien species: a toolkit of best prevention and management practices*. CAB International, Wallingford.
- ZENNI, R.D., WILSON, J.R.U., LE ROUX, J.J. & RICHARDSON, D.M. 2009. Evaluating the invasiveness of *Acacia paradoxa* in South Africa. *South African Journal of Botany* 75:485-496.