























# Effects of Selenium-enriched Diets on the Growth Performance, Slaughter Characteristics, and Blood Biochemistry of Rhode Island Red Chicken

## ■ Author(s)

Khan MT <sup>*</sup>	 <a href="https://orcid.org/0000-0002-1963-8608">https://orcid.org/0000-0002-1963-8608</a>
Ahmed S <sup>*</sup>	 <a href="https://orcid.org/0000-0003-0310-3446">https://orcid.org/0000-0003-0310-3446</a>
Azhar M <sup>*</sup>	 <a href="https://orcid.org/0000-0001-9529-677X">https://orcid.org/0000-0001-9529-677X</a>
Asad T <sup>*</sup>	 <a href="https://orcid.org/0000-0003-0858-0196">https://orcid.org/0000-0003-0858-0196</a>
Arslan M <sup>*</sup>	 <a href="https://orcid.org/0000-0002-1160-2942">https://orcid.org/0000-0002-1160-2942</a>
Raziq F <sup>*</sup>	 <a href="https://orcid.org/0000-0003-2021-4201">https://orcid.org/0000-0003-2021-4201</a>
Gondal MA <sup>III</sup>	 <a href="https://orcid.org/0000-0002-7302-5829">https://orcid.org/0000-0002-7302-5829</a>
Rauf M <sup>IV</sup>	 <a href="https://orcid.org/0000-0002-3269-2596">https://orcid.org/0000-0002-3269-2596</a>
Liaqat S <sup>IV</sup>	 <a href="https://orcid.org/0000-0001-6531-5124">https://orcid.org/0000-0001-6531-5124</a>
Bachaya H <sup>VI</sup>	 <a href="https://orcid.org/0000-0001-9202-1076">https://orcid.org/0000-0001-9202-1076</a>
Qumar M <sup>VI</sup>	 <a href="https://orcid.org/0000-0001-6239-2036">https://orcid.org/0000-0001-6239-2036</a>
Faran G <sup>VIII</sup>	 <a href="https://orcid.org/0000-0003-0079-099X">https://orcid.org/0000-0003-0079-099X</a>
Abbasi F <sup>IX</sup>	 <a href="https://orcid.org/0000-0001-9331-6589">https://orcid.org/0000-0001-9331-6589</a>
Farroq Z <sup>X</sup>	 <a href="https://orcid.org/0000-0002-1956-8785">https://orcid.org/0000-0002-1956-8785</a>
Wadood F <sup>XI</sup>	 <a href="https://orcid.org/0000-0001-5716-5533">https://orcid.org/0000-0001-5716-5533</a>
Iqbal ZM <sup>XII</sup>	 <a href="https://orcid.org/0000-0003-1296-6681">https://orcid.org/0000-0003-1296-6681</a>
Abbasi G <sup>XIII</sup>	 <a href="https://orcid.org/0000-0002-2741-994X">https://orcid.org/0000-0002-2741-994X</a>
Bughio E <sup>XIV</sup>	 <a href="https://orcid.org/0000-0002-1409-527X">https://orcid.org/0000-0002-1409-527X</a>
Younas U <sup>XV</sup>	 <a href="https://orcid.org/0000-0002-5107-9391">https://orcid.org/0000-0002-5107-9391</a>
Raja IH <sup>XVI</sup>	 <a href="https://orcid.org/0000-0003-1986-1161">https://orcid.org/0000-0003-1986-1161</a>

\*Affiliations

## ■ Mail Address

Corresponding author e-mail address  
M. T. Khan  
Department of Poultry Science, Cholistan  
University of Veterinary and Animal Sciences,  
Bahawalpur, Punjab, Zip Code 63100,  
Pakistan.  
Phone: +92 300 8360008  
Email: [mtahirkhan@cuvas.edu.pk](mailto:mtahirkhan@cuvas.edu.pk)

## ■ Keywords

Selenium, growth, slaughter characteristics,  
blood chemistry.



Submitted: 16/November/2022  
Approved: 06/September/2023

## ABSTRACT

The goal of the current study was to determine how selenium-supplemented diets affected the growth performance, carcass traits, and blood biochemistry of Rhode Island Red (RIR) chickens. Organic and inorganic selenium (SE) was fed to the birds at 0.30ppm, whereas the control diet was not supplemented with selenium in either organic or inorganic form. A total of 225 day-old RIR chicks were indiscriminately distributed in three groups according to the experimental diets, in a completely randomized design. Each treatment group was repeated 5 times (replicates), and each replicate had 15 birds. Parameters of growth performance, carcass characteristics, and blood biochemistry were assessed. Birds fed an inorganic selenium-supplemented diet had higher feed intake than those fed an organic selenium-supplemented diet or the control diet, whereas birds fed an organic selenium-supplemented diet had higher body weight gains and better feed conversion ratios. Birds fed organic SE in the diet showed higher breast and thigh weight than those receiving inorganic Se, whereas dressing percentage, drumstick yield, and weights of liver, gizzard, heart, and wing were not significantly different across treatments. The organic Se group showed higher values for total protein and globulin in the blood relative to those fed inorganic Se and the control diet. Similarly, organic Se fed birds showed higher blood Se concentration than the other two groups. However, no significant differences in albumin, glucose, cholesterol, triglycerides, and uric acid were observed among the diets. These results lead to the conclusion that organic Se may be utilized in diets to improve the poor performance of RIR chicken.

## INTRODUCTION

Animal protein scarcity is one of the biggest problems in developing nations today (Shafiq, 2016). The chicken industry offers a reasonably affordable supply of meat, eggs, and other necessary nutrients. With an investment of over Rs. 750 billion, Pakistan's poultry industry is one of the most organized sectors of the country's agricultural economy, employing more than 1.5 million people (Pakistan Economic Survey, 2021-2022). Over the past ten years, this industry has grown at an amazing rate of about 7.5 percent per year (Pakistan Economic Survey, 2021-2022). One of the American poultry breeds that has adapted particularly well to the Asian climate is the Rhode Island Red (RIR) chicken (Parveen *et al.*, 2017). They are dependable birds, with a keen sense of curiosity, and are also great mothers. Moreover, they produce very high-quality meat, lay enormous, brown-tinted eggs and have a better level of tolerance to external parasites than other breeds bred in semi-intensive or extensive settings. RIR chicken, on the other hand, has a dismal track record for weight gain. Nevertheless, the performance



of RIR chicken may be enhanced by nutritional interventions such as dietary SE supplementation.

Animals need selenium (Se), a vital antioxidant mineral (Surai, 2002) that has been shown to enhance immunological, antioxidative, and growth performance in broilers (Swain *et al.*, 2000; Wang & Xu, 2008; Wang *et al.*, 2011a; Liao *et al.*, 2012; Skrivan *et al.*, 2012). Glutathione peroxidase (GSH-Px), a vital component of the antioxidative protection system, is one of the proteins or enzymes that includes selenium (Hoffmann & Berry, 2008; Cai *et al.*, 2012; Lee *et al.*, 2014; Dalia *et al.*, 2018). Different types of Se are available for supplemental use in poultry feed, including inorganic sodium selenite or selenite and organic Se yeast (Yuan *et al.*, 2011; Surai & Fisinin, 2014). The latter is included in amino acids and short peptides like selenomethionine, selenocysteine, and selenocystine. Se compounds present in grains and forages share similarities with the organic form originating from yeast. Using the amino acid transport mechanism, animals and poultry have evolved naturally to absorb this form from all parts of the gastrointestinal tract. In comparison to inorganic salts, organic Se is more active, less poisonous, and accumulates at higher levels in organs like the liver, brain, and muscles (Kim & Mahan, 2001; Surai, 2002; Payne & Southern, 2005; Tiwary *et al.*, 2006; Zhang *et al.*, 2008; Attia *et al.*, 2010; Mohapatra *et al.*, 2014; Suchy *et al.*, 2014; Hassan *et al.*, 2021). Chickens' consumption of antioxidant vitamins A, C, and E is intimately related to the absorption and transport of Se. Birds have evolved to metabolize Se and take advantage of its antioxidant qualities because it is a necessary component of the majority of diets (Surai, 2006). Poultry feed, therefore, must contain organic Se to maintain the wellbeing of animals, their effective performance, and high-quality meat (Jiang *et al.*, 2009). It was, therefore, hypothesized that supplementing diets with organic selenium could improve the poor growth and slaughter features of RIR chicks without negatively affecting blood chemistry. The purpose of this study was to compare the effects of various Se sources on the growth performance, carcass traits, and blood biochemistry of RIR chicken.

## MATERIALS AND METHODS

### Ethics

The aforementioned study was carried out in accordance with the recommendations of the Cholistan University of Veterinary and Animal Sciences, Bahawalpur, Ethical Review Committee.

An ethics certificate was obtained from the Office of Research Innovation and Commercialization, CUVAS, Bahawalpur, prior to the start of the trial (Letter No: ORIC-190, Dated: 29-08-22).

### Experiment site, birds, and housing

During the course of 42 days, the current study was carried out at the PRO-MAX Protein Farm in Punjab, Pakistan. 225 day-old RIR chicks were indiscriminately distributed into three groups based on three diets, namely organic (Se-enriched yeast), inorganic (sodium selenite), and control diet, under a completely randomized design (CRD). Each group was repeated 5 times (replicates/pens), with 15 birds in each pen. The experimental birds were placed in an open-sided house, which was built to be well-ventilated. A basal diet (Table 1 and 2) based on corn-soybean meal, containing the same amount of protein and energy, was offered to the birds. It was formulated keeping in view the dietary needs of the birds (NRC, 1994), containing 20-22% of crude protein and 2,800-3200 metabolizable energy (kcal/kg), respectively. Before the start of the trial, the

**Table 1** – Composition of basal diet for the starter phase.

Ingredients (%)	Quantity
Maize	52.200
Soybean meal	29.000
Rice polish	10.272
Canola meal	3.000
Poultry-by-product meal	3.000
Limestone	1.000
Lysine HCL	0.310
MCP <sup>1</sup>	0.300
DL-methionine	0.263
Salt	0.220
Soda	0.100
Threonine	0.100
Betaine HCL	0.075
Vitamin premix <sup>2</sup>	0.055
Mineral premix <sup>2</sup>	0.055
Enramycin	0.030
Coxiril®	0.010
Phytase	0.010
Nutrient	
ME (kcal/kg)	2900
CP (%)	22.000
Moisture (%)	10.500
Ash (%)	5.000
Crude fat (%)	4.000
Crude fiber (%)	3.000
Selenium, ppm	0.04

<sup>1</sup>MCP=monocalcium phosphate.

<sup>2</sup>Provided per kg of diet: vitamin A, 11,000 IU; vitamin D<sub>3</sub>, 2,560 IU; vitamin E, 44 IU; vitamin K, 4.2 mg; riboflavin, 8.5 mg; niacin, 48.5 mg; thiamine, 3.5 mg; d-pantothenic, 27 mg; choline, 150 mg; vitamin B<sub>12</sub>, 33 µg; copper, 8 mg; zinc, 75 mg; manganese, 55 mg; iodine, 0.35 mg.



**Table 2** – Composition of basal diet for the finisher phase.

Ingredients (%)	Quantity
Maize	67.00
Soybean meal	22.00
Poultry-by-product meal	4.000
Rapeseed meal	2.900
Corn gluten 60%	1.700
Limestone	0.800
Lysine HCL	0.333
DL-Methionine	0.224
Salt	0.220
MCP <sup>1</sup>	0.200
Rice polish	0.200
Soda	0.100
Threonine	0.090
Vitamin premix <sup>2</sup>	0.055
Mineral premix <sup>2</sup>	0.055
Betaine HCL	0.050
Enramycin	0.030
Coxiril**	0.010
Phytase	0.010
<b>Nutrient</b>	
ME (kcal/kg)	3150
CP (%)	20.000
Moisture (%)	10.500
Crude fat (%)	4.500
Ash (%)	4.000
Crude fiber (%)	4.000

<sup>1</sup>MCP=monocalcium phosphate.

<sup>2</sup>Provided per kg of diet: vitamin A, 11,000 IU; vitamin D<sub>3</sub>, 2,560 IU; vitamin E, 44 IU; vitamin K, 4.2 mg; riboflavin, 8.5 mg; niacin, 48.5 mg; thiamine, 3.5 mg; d-pantothenic, 27 mg; choline, 150 mg; vitamin B<sub>12</sub>, 33 µg; copper, 8 mg; zinc, 75 mg; manganese, 55 mg; iodine, 0.35 mg.

basal diet was examined to determine the amount of SE it contained. Each pen (1 × 1 m<sup>2</sup>) contained a nipple drinker and two round feeders to provide water and feed *ad-libitum* to the birds. The chicks were brooded at 34 ± 1.1°C temperature with relative humidity set at 62 ± 3%, during the first week. Temperature was gradually decreased to 24°C with RH 65%. Natural day light was provided to birds throughout the study period. The chicks were vaccinated against infectious bronchitis, Newcastle disease, avian influenza H9, and infectious bursal disease according to the instructions given by the company. A deep litter system was prepared before trial, whereon rice husk was spread as bedding material.

## Data collection

### Growth performance and carcass characteristics

The performance parameters of feed intake (FI), body weight gains (BWG), and feed conversion ratio (FCR) were assessed, and the data were then recorded on a weekly basis. The weight gain and FCR were

computed using feed intake and body weight. The deceased birds were removed twice daily, and FCR was modified accordingly. Three broilers with average weights per pen were chosen to collect data on carcass characteristics at the end of the trial. These birds were starved for four hours before being slaughtered in accordance with Halal guidelines, and the carcasses were then defeathered and eviscerated. Breast, thigh, drumstick, wing, liver, gizzard, and heart were dissected and separately weighed in order to gather data on carcass characteristics. The ratio of dressed weight to live weight (excluding skin, shanks, and head) was multiplied by 100 to determine the dressing percentage. Similarly, thigh, drumstick, breast, liver, wing, heart, and gizzard weight were calculated as the ratio between individual weight of each organ and live weight multiplied by 100.

### Blood biochemistry

At the moment of slaughter, measurements of blood indices like albumin, total protein, glucose, globulin, cholesterol, uric acid, and triglycerides were taken. For this, samples of 3 ml of blood from three birds per replicate were taken from the wing vein. An anticoagulant was used inside a single-use syringe measuring 5 ml. Blood vacutainers were used in order to transfer the blood. Serum was extracted from the blood by centrifuging it at 3,000 g for 10 minutes. Serum was stored in 1.5-mL Eppendorf tubes at -20°C for later examination. Blood parameters were determined by spectrophotometric analysis of the serum (Rehman *et al.*, 2017). Prior to analysis, frozen serum was thawed at 4°C.

### Statistical analysis

Analysis of variance (ANOVA) technique with GLM procedure of SAS was applied to analyze data recorded, and means were compared through Duncan's multiple range test by taking each pen as experimental unit.

## RESULTS

### Growth performance

The results indicated significant differences in growth performance parameters among the diets. The birds fed inorganic selenium showed increased feed intake, followed by those fed organic Se and the control diet. On the other hand, birds fed organic Se exhibited higher body weight gain compared with those fed inorganic Se. Similarly, birds fed organic Se showed improved feed conversion ratio, followed by those fed inorganic Se and the control group (Table 3).



**Table 3** – Effect of selenium-supplemented diets on the growth performance of growing Rhode Island Red chicks<sup>1</sup>.

Parameters <sup>2</sup>	Treatments			p-value
	Organic Se	Inorganic Se	Control	
FI (kg)	1.235 <sup>a</sup>	1.280 <sup>ab</sup>	1.175 <sup>b</sup>	0.03
BWG (g)	333.66 <sup>a</sup>	318.60 <sup>b</sup>	280.40 <sup>c</sup>	0.05
FCR	3.70 <sup>c</sup>	4.01 <sup>b</sup>	4.19 <sup>a</sup>	0.02

<sup>a-c</sup>Different letters on treatment means within a row show significant differences ( $p < 0.05$ ).

<sup>1</sup>Data are means of 5 replicates (n=5) with 15 birds per replicate.

<sup>2</sup>FI: feed intake, BWG: body weight gain, FCR: feed conversion ratio.

### Slaughter characteristics

Our results showed that organic SE supplementation increased breast and thigh weights when compared to inorganic SE, but no other metrics, such as dressing percentage, drumstick weight, liver weight, gizzard weight, heart weight, or wing weight, significantly varied between treatments (Table 4).

**Table 4** – Effect of selenium-supplemented diets on the slaughter traits of growing Rhode Island Red chicks<sup>1</sup>.

Parameters <sup>2</sup>	Treatments			p-value
	Organic Se	Inorganic Se	Control	
CY (%)	62.9	61.5	61.2	0.06
BY (%)	22.9 <sup>a</sup>	20.7 <sup>b</sup>	21.5 <sup>ab</sup>	0.005
TY (%)	21.2 <sup>a</sup>	20.2 <sup>b</sup>	20.9 <sup>ab</sup>	0.02
DW (%)	16.4	15.1	15.3	0.0025
LW (%)	3.4	3.7	3.5	0.063
GW (%)	3.2	3.5	3.3	0.055
HW (%)	0.7	0.6	0.7	0.05
WW (%)	8.5	8.1	7.9	0.0172

<sup>a-b</sup>Different letters on treatment means within a row show significant differences ( $p < 0.05$ ).

<sup>1</sup>Data are means of 5 replicates (n=5) with 15 birds per replicate.

<sup>2</sup>CY: carcass yield, BY: breast yield, TY: thigh yield, DW: drumstick weight, LW: liver weight, GW: gizzard weight, HW: heart weight, WW: wing weight.

### Blood biochemistry

When compared to birds fed inorganic Se and the control diet, birds fed organic Se had greater blood levels of total protein and globulin. Additionally, it was discovered that organic Se-fed birds had greater blood Se concentrations than control and inorganic Se-fed birds. However, no differences ( $p > 0.05$ ) between treatments were found for albumin, glucose, cholesterol, triglycerides, or uric acid levels (Table 5).

## DISCUSSION

### Se effect on growth performance

The poultry industry requires efficient growth to maintain business profitability. The weather was hot

and humid when this study was conducted. According to the reports (Siegel, 1995; Melesse *et al.*, 2011; El-Deep *et al.*, 2014; Surai, 2016), high ambient temperature reduces the feed intake, live weight gain, feed efficiency, and immune response of broiler chicken. Moreover, hyperthermia may promote the formation of reactive oxygen species (ROS) (Mujahid *et al.*, 2005). Excessive ROS levels disturb the balance between oxidation and antioxidant defense systems, resulting in lipid peroxidation (Shimizu *et al.*, 2006) and oxidative damage to immune responses (Surai & Fisinin, 2016b), absorption of nutrients (Surai and Fisinin, 2015), and feed conversion efficiency (Belhdj *et al.*, 2015; Vasantha *et al.*, 2016). Therefore, a balance between ROS production and the antioxidant system must be established to maintain immune function, health, and productivity (Surai, 2002). Natural antioxidants including probiotics, vitamins, and trace elements play vital roles in protecting cells from ROS by reducing free radicals and preventing the peroxidation of lipids (Eid *et al.*, 2003; Lin *et al.*, 2006; Eid *et al.*, 2008; Sahin *et al.*, 2009). According to earlier reports, in comparison to its inorganic form, Se in the organic form is more potent (Sahin *et al.*, 2008; Surai, 2016) and eliminates free radicals from metabolic activity more effectively (Papazyan *et al.*, 2006; Sluis, 2007; Hanafy *et al.*, 2009; Attia *et al.*, 2010). Hence, it is possible to assume that including organic Se in the diet of RIR chicks improved growth performance by reducing the negative effects of high ambient temperature and promoting better nutrient utilization.

These results are consistent with reports of other authors (Zhou & Wang, 2011; Markovic *et al.*, 2018; Saleh & Ebeid, 2019; Gul *et al.*, 2021). Contrarily, Se supplementation in broilers has also been observed to not have any effect on growth performance (Payne & Southern, 2005; Ryu *et al.*, 2005; Peric *et al.*, 2009; Cai *et al.*, 2012; Chen *et al.*, 2013; Briens *et al.*, 2014; Chen *et al.*, 2014; Oliveira *et al.*, 2014; Boostani *et al.*, 2015; Chadjo *et al.*, 2015; Liu *et al.*, 2015; Bakhshalinejad *et al.*, 2018; Li *et al.*, 2018; Bakhshalinejad *et al.*, 2019). Furthermore, it has also been demonstrated that broilers' growth performance could be improved through the supplementation of Se using either sodium selenite or Se yeast (Mansoub *et al.*, 2010; Wang *et al.*, 2021).

### Se effect on slaughter traits

Our results showed that organic SE supplementation increased breast and thigh weights when compared



to inorganic SE. The treatment group's increased proportion of breast and thigh tissue was probably caused by the higher absorption and improved development brought on by the organic Se enhancing the metabolism of thyroid hormones that regulated animal growth (Arthur, 1991; Jianhua *et al.*, 2000). These findings are in line with what other writers have reported (Saleh & Ebeid, 2019). On the other hand, broilers given Se-supplemented diets and those fed the control diet showed no differences ( $p>0.05$ ) in the percentages of breast and thigh muscles. Our results showed that Se supplementation had no effect on carcass weights and the percentages of drumstick, liver, gizzard, heart, and wings. Similar outcomes were obtained by other researchers, who discovered no connection between Se supplementation and carcass weights (Choct & Naylor, 2004; Payne & Southern, 2005; Sevcikova *et al.*, 2006; Heindl *et al.*, 2010; Chen *et al.*, 2013; Wang *et al.*, 2021) or carcass parts outputs (Downs *et al.*, 2000; Choct & Naylor, 2004; Payne & Southern, 2005; Sevcikova *et al.*, 2006; Heindl *et al.*, 2010; Cai *et al.*, 2012; Chen *et al.*, 2013; Hada *et al.*, 2013; Chen *et al.*, 2014; Oliveira *et al.*, 2014; Baltic *et al.*, 2016; Bakhshalinejad *et al.*, 2018; Jamnongtoi *et al.*, 2018; Li *et al.*, 2018; Bakhshalinejad *et al.*, 2019). In contrast, organic forms of Se increased eviscerated weight in one study (Choct *et al.*, 2004).

### Se effect on blood biochemistry

Animals' physiological and general health can be assessed using biochemical indicators (Qiu *et al.*, 2021), which are positively connected with their dietary state (Adejumo, 2004). High levels of oxidative stress on blood metabolites and increased generation of reactive oxygen species are associated with selenium deficiency (Zheng *et al.*, 2019). In order to avoid oxidative stress and free radical damage to blood metabolites, an enzyme known as glutathione peroxidase is essential. Selenium indirectly helps to reduce oxidative damage to blood metabolites since selenium is present in this enzyme.

In the current investigation, the levels of total protein and globulin in the blood of the birds fed organic Se were higher than those of the control diet and birds fed inorganic Se. The causes for this were not immediately apparent. Furthermore, it was shown that birds fed organic Se had higher blood Se concentrations than control and inorganic Se-fed birds. Albumin, glucose, cholesterol, triglycerides, and uric acid levels did not differ ( $p>0.05$ ) between treatments (Table 5). This supports earlier findings that selenium

had no significant effects on broiler (Hosseini Mansoub *et al.*, 2010; Rashidi *et al.*, 2010; Kanchana & Jeyanthi, 2010; Habibian *et al.*, 2014; Khalifa *et al.*, 2021) or layer (Lin *et al.*, 2020) serum biochemical markers. On the other hand, it was discovered that feeding broilers Se had a substantial impact on the plasma levels of total cholesterol and triglycerides (Saleh & Ebeid, 2019). In the current study, the organically fed birds had greater blood Se concentrations than the control and inorganically fed birds. The fact that the birds given organic Se had greater blood selenium concentrations is consistent with other previous results (Gul *et al.*, 2021).

**Table 5** – Effect of selenium-supplemented diets on the blood profile of growing Rhode Island Red chicks<sup>1</sup>.

Parameters <sup>2</sup>	Treatments			P-value
	Organic Se	Inorganic Se	Control	
TP (g/dL)	3.4	3.3	3.5	0.05
AB (g/dL)	1.8	1.7	1.7	0.15
GB (g/dL)	1.5	1.5	1.7	0.05
GL (mg/dL)	138.3	140.7	137.2	0.35
CH (m/dL)	160.2	158.7	157.5	0.70
TR (mg/dL)	91.7	90.4	89.6	0.52
UA (g/dL)	3.8	3.9	3.6	0.72
Se (µg/L)	57.35 <sup>a</sup>	50.20 <sup>b</sup>	41.15 <sup>c</sup>	0.002

<sup>a,b</sup>Different letters on treatment means within a row indicate significant differences ( $p<0.05$ ).

<sup>1</sup>Data are means of 5 replicates (n=5) with 15 birds per replicate.

<sup>2</sup>TP: total protein, AB: albumin, GB: globulin, GL: glucose, CH: cholesterol, TR: triglycerides, UA: uric acid.

## CONCLUSIONS

Nutritional supplementation with Se enriched yeast, an organic source of selenium, may help improve the growth and slaughter features of Rhode Island Red chicks without having a detrimental impact on blood biochemistry.

## ACKNOWLEDGEMENTS

The authors extend their appreciation to Researchers Supporting Project number (RSPD2024R-965), King Saud University, Riyadh, Saudi Arabia and the management of the PRO-MAX Protein Farm for facilitating the trial.

## DISCLOSURE OF POTENTIAL CONFLICT OF INTEREST

All authors of the present study declare that there is no financial or any other conflict of interest.



## AUTHOR CONTRIBUTIONS

Conceptualization, M.T.K. and M.A.; methodology, M.T.K., T.A. and M.A.; software, F.R., M.A.G., M.Q. and G.F.; validation, M.R., S.L. and F.A.; formal analysis, M.T.K., Z.M.I., F.W. and I.H.R.; investigation, M.T.K., S.A. and M.A.; data curation, S.A., M.A., T.A. and M.A.; writing—original draft preparation, M.T.K., S.A., S.N., H.A.B., G.A., Z.F. and E.B.; writing—review and editing, U.Y., D.F., F.S.A., Q.N. and M.S.; supervision, M.T.K. and M.A. All authors have read and agreed to the published version of the manuscript.

## REFERENCES

- AAFCO - Association of American Feed Control Officials. Official publication. Champaign: Association of American Feed Control Officials; 1972.
- Adejumo DO. Performance, organ development and hematological indices of rats fed sole diets of graded levels of cassava flour and soybean flour (soygari) as substitutes for energy and protein concentrates. *Tropical Journal of Animal Science* 2004;7:57-63. <https://doi.org/10.4236/jns.2014.514142>
- Arthur JR. The role of selenium in thyroid hormone metabolism. *Canadian Journal of Physiology and Pharmacology* 1991;69:1648-52. <https://doi.org/10.1007/BF02783686>
- Attia YA, Abdalah AA, Zeweil HS, *et al.* Effect of inorganic or organic selenium supplementation on productive performance, egg quality and some physiological traits of dual-purpose breeding hens. *Czech Journal of Animal Science* 2010;55(11):505-19. <https://doi.org/10.17221/1702-CJAS>
- Bakhshalinejad R, Akbari Moghaddam Kakhki R, *et al.* Effects of different dietary sources and levels of selenium supplements on growth performance, antioxidant status and immune parameters in Ross 308 broiler chickens. *British Poultry Science* 2018;59(1):81-91. <https://doi.org/10.1080/00071668.2017.1380296>
- Bakhshalinejad R, Hassanabadi A, Swick RA. Dietary sources and levels of selenium supplements affect growth performance, carcass yield, meat quality and tissue selenium deposition in broilers. *Animal Nutrition* 2019;5(3):256-63. <https://doi.org/10.1016/j.aninu.2019.03.003>
- Baltic MZ, Dokmanovic Starcevic M, Basic M, *et al.* Effects of dietary selenium-yeast concentrations on growth performance and carcass composition of ducks. *Animal Production Science* 2016;57(8):1731-7. <https://doi.org/10.1071/AN16126>
- Belhadj S, Najar T, Ghram A, *et al.* Heat stress effects on livestock: Molecular, cellular and metabolic aspects: A review. *Journal of Animal Physiology and Animal Nutrition* 2015;100(3):401-12. <https://doi.org/10.1111/jpn.12379>
- Boostani A, Sadeghi AA, Mousavi SN, *et al.* Effects of organic, inorganic, and nano-Se on growth performance, antioxidant capacity, cellular and humoral immune responses in broiler chickens exposed to oxidative stress. *Livestock Science* 2015;178:330-6. <https://doi.org/10.1016/j.livsci.2015.05.004>
- Briens M, Mercier Y, Rouffineau F, *et al.* 2-hydroxy-4-methylselenobutanoic acid induces additional tissue selenium enrichment in broiler chickens compared with other selenium sources. *Poultry Science* 2014;93(1):85-93. <https://doi.org/10.3382/ps.2013-03182>
- Cai SJ, Wu CX, Gong LM, *et al.* Effects of nano-selenium on performance, meat quality, immune function, oxidation resistance, and tissue selenium content in broilers. *Poultry Science* 2012;91(10):2532-9. <https://doi.org/10.3382/ps.2012-02160>
- Cantor AH, Scott ML, Noguchi T. Biological availability of selenium in feedstuffs and selenium compounds for prevention of exudative diathesis in chicks. *Journal of Nutrition* 1975;105(1):96-105. <https://doi.org/10.1093/jn/105.1.96>
- Chadio SE, Pappas AC, Papanastatos A, *et al.* Effects of high selenium and fat supplementation on growth performance and thyroid hormones concentration of broilers. *Journal of Trace Elements in Medicine and Biology* 2015;29:202-7. <https://doi.org/10.1016/j.jtemb.2014.09.010>
- Chen G, Wu J, Li C. The effect of different selenium levels on production performance and biochemical parameters of broilers. *Italian Journal of Animal Science* 2013;12(4):486-91. <https://doi.org/10.4081/ijas.2013.e79>
- Chen G, Wu J, Li C. Effect of different selenium sources on production performance and biochemical parameters of broilers. *Journal of Animal Physiology and Animal Nutrition* 2014;98(4):747-54. <https://doi.org/10.1111/jpn.12136>
- Choct M, Naylor AJ, Reinke N. Selenium supplementation affects broiler growth performance, meat yield and feather coverage. *British Poultry Science* 2004;45(5):677-83. <https://doi.org/10.1080/00071660400006495>
- Choct M, Naylor AJ. The effect of dietary selenium source and vitamin E levels on performance of male broilers. *Asian Australasian Journal of Animal Science* 2004;17(7):1000-6. <https://doi.org/10.5713/ajas.2004.1000>
- Dalia AM, Loh TC, Sazili AQ, *et al.* Effects of vitamin E, inorganic selenium, bacterial organic selenium, and their combinations on immunity response in broiler chickens. *BMC Veterinary Research* 2018;14(1):1-10. <https://doi.org/10.1186/s12917-018-1578-x>
- Downs KM, Hess JB, Bilgili SF. Selenium source effect on broiler carcass characteristics, meat quality and drip loss. *Journal of Applied Animal Research* 2000;18(1):61-71. <https://doi.org/10.1080/09712119.2000.9706324>
- Pakistan Economic Survey. Economic survey of Pakistan. Islamabad: Ministry of National Food Security & Research; 2021-2022.
- Eid YZ, Ohtsuka A, Hayashi K. Tea polyphenols reduce glucocorticoid-induced growth inhibition and oxidative stress in broiler chickens. *British Poultry Science* 2003;44(1):350-6. <https://doi.org/10.1080/0007166031000085427>
- Eid Y, Ebeid T, Moawad M, *et al.* Reduction of dexamethasone-induced oxidative stress and lipid peroxidation in laying hens by dietary vitamin E supplementation. *Emirates Journal of Food and Agriculture* 2008;20(2):28-40. <https://doi.org/10.9755/efja.v20i2.5188>
- El-Deep MH, Ijiri D, Eid YZ, *et al.* Effects of dietary supplementation with *Aspergillus awamori* on growth performance and antioxidant status of broiler chickens exposed to high ambient temperature. *Journal of Poultry Science* 2014;51(3):281-8. <https://doi.org/10.2141/jpsa.0130154>
- Gul F, Ahmad B, Afzal S, *et al.* Comparative analysis of various sources of selenium on the growth performance and antioxidant status in broilers under heat stress. *Brazilian Journal of Biology* 2021;83:1-10. <https://doi.org/10.1590/1519-6984.251004>
- Habibian M, Ghazi S, Moeini MM, *et al.* Effects of dietary selenium and vitamin E on immune response and biological blood parameters of broilers reared under thermoneutral or heat stress conditions. *International*



- Journal of Biometeorology 2014;58(5):741-52. <https://doi.org/10.1007/s00484-013-0654-y>
- Hada FH, Malheiros RD, Silva JDT, *et al.* Effect of protein, carbohydrate, lipid, and selenium levels on the performance, carcass yield, and blood changes in broilers. *Brazilian Journal of Poultry Science* 2013;15(4):385-94. <https://doi.org/10.1590/S1516-635X2013000400014>
- Hanafy MM, El-Sheik AMH, Abdalla EA. The effect of organic selenium supplementation on productive and physiological performance in a local strain of chicken. 1-The effect of organic selenium (Sel-Plex™) on productive, reproductive and physiological traits of Bandarah local strain. *Egyptian Poultry Science Journal* 2009;29(4):1061-84. Available from: <http://www.epsaegypt.com/.../9-1158.pdf>
- Hassan F, Mobarez S, Mohamed M, *et al.* Zinc and/or Selenium Enriched Spirulina as Antioxidants in Growing Rabbit Diets to Alleviate the Deleterious Impacts of Heat Stress during Summer Season. *Animals* 2021;11(3):756. <https://doi.org/10.3390/ani11030756>
- Heindl J, Ledvinka Z, Englmaierova M, *et al.* The effect of dietary selenium sources and levels on performance, selenium content in muscle and glutathione peroxidase activity in broiler chickens. *Czech Journal of Animal Science* 2010;55(12):572-8. <https://doi.org/10.17221/2487-CJAS>
- Hoffmann PR, Berry MJ. The influence of selenium on immune responses. *Molecular Nutrition and Food Research* 2008;52(11):1273-80. <https://doi.org/10.1002/mnfr.200700330>
- Hosseini-Mansoub N, Chekani-Azar S, Tehrani AA, *et al.* Influence of dietary vitamin E and zinc on performance, oxidative stability and some blood measures of broiler chickens reared under heat stress (35 °C). *Journal of Agrobiology* 2010;27(2):103-10. <https://doi.org/10.2478/s10146-009-0012-1>
- Jamngotai P, Sivapirunthep P, Chaosap C. Effect of dietary organic and inorganic selenium on carcass composition and meat characteristics of broiler chickens. *International Journal of Agricultural Technology* 2018;14(7):1279-1286. Available from: [http://www.ijat-aatsea.com/pdf/v14\\_n7...](http://www.ijat-aatsea.com/pdf/v14_n7...)
- Jiang ZY, Lin YC, Zhou GL, *et al.* Effects of dietary selenomethionine supplementation on growth performance, meat quality and antioxidant property in yellow broilers. *Journal of Agricultural and Food Chemistry* 2009;57(20):9769-72. <https://doi.org/10.1021/jf902411c>
- Jianhua H, Ohtsuka A, Hayashi K. Selenium influences growth via thyroid hormone status in broiler chickens. *British Journal of Nutrition* 2000;84(5):727-32. <https://doi.org/10.1017/S0007114500002087>
- Kanchana G, Jeyanthi GP. The effect of supplementation of diet with vitamin-E and selenium and their combinations on the performance and lipid profiles of layer chickens. *International Journal of Pharma and Bio Sciences* 2010;1(1):1-11. Available from: <https://www.researchgate.net/publication/286122821>
- Khalifa OA, Al Wakeel RA, Hemed SA, *et al.* The impact of vitamin E and/or selenium dietary supplementation on growth parameters and expression levels of the growth-related genes in broilers. *BMC Veterinary Research* 2021;17(1):251. <https://doi.org/10.1186/s12917-021-02963-1>
- Kim YY, Mahan DC. Comparative effects of high dietary levels of organic and inorganic selenium on selenium toxicity of growing-finishing pigs. *Journal of Animal Science* 2001;79(4):942-8. <https://doi.org/10.2527/2001.794942x>
- Lee SH, Lillehoj HS, Jang SI, *et al.* Effects of in ovo injection with selenium on immune and antioxidant responses during experimental necrotic enteritis in broiler chickens. *Poultry Science* 2014;93(5):1113-21. <https://doi.org/10.3382/ps.2013-03770>
- Li JL, Zhang L, Yang ZY, *et al.* Effects of different selenium sources on growth performance, antioxidant capacity and meat quality of local Chinese Subei chickens. *Biological Trace Element and Research* 2018;181(2):340-6. <https://doi.org/10.1007/s12011-017-1049-4>
- Liao XD, Lu L, Li SF, *et al.* Effects of selenium source and level on growth performance, tissue selenium concentrations, antioxidation, and immune functions of heat-stressed broilers. *Biological Trace Element and Research* 2012;150(1-3):158-65. [https://doi.org/10.1016/S2095-3119\(20\)63432-3](https://doi.org/10.1016/S2095-3119(20)63432-3)
- Lin H, Jiao HC, Buyse J, *et al.* Strategies for preventing heat stress in poultry. *World's Poultry Science Journal* 2006;62(1):71-86. <https://doi.org/10.1079/WPS200585>
- Lin X, Yang T, Li H, *et al.* Interactions between different selenium compounds and essential trace elements involved in the antioxidant system of laying hens. *Biological Trace Element and Research* 2020;193(1):252-60. <https://doi.org/10.1007/s12011-019-01701-x>
- Liu S, Tan H, Wei S, *et al.* Effect of selenium sources on growth performance and tissue selenium retention in yellow broiler chicks. *Journal of Applied Animal Research* 2015;43(4):487-90. <http://doi.org/10.1080/09712119.2014.978780>
- Markovic R, Ćirić J, Drljačić A, *et al.* The effects of dietary Selenium-yeast level on glutathione peroxidase activity, tissue Selenium content, growth performance, and carcass and meat quality of broilers. *Poultry Science* 2018;97(8):2861-70. <https://doi.org/10.3382/ps/pey117>
- Melesse A, Maak S, Schmidt R, *et al.* Effect of long-term heat stress on some performance traits and plasma enzyme activities in naked-neck chickens and their F1 crosses with commercial layer breeds. *Livestock Science* 2011;141(2-30):227-31. <https://doi.org/10.1016/j.livsci.2011.06.007>
- Mohapatra P, Swain RH, Mishra SK, *et al.* Effects of dietary nano-selenium on tissue selenium deposition, antioxidant status and immune functions in layer chicks. *International Journal of Pharmacology* 2014;10(3):160-7. <https://doi.org/10.3923/ijp.2014.160.167>
- Mujahid A, Yoshiki Y, Akiba Y, *et al.* Superoxide radical production in chicken skeletal muscle induced by acute heat stress. *Poultry Science* 2005;84(2):307-14. <https://doi.org/10.1093/ps/84.2.307>
- NRC - National Research Council. Nutrient requirements of poultry. 9<sup>th</sup> rev. Washington: National Academy Press; 1994.
- Oliveira TFB, Rivera DFR, Mesquita FR, *et al.* Effect of different sources and levels of selenium on performance, meat quality, and tissue characteristics of broilers. *Journal of Applied Poultry Research* 2014;23(1):15-22. <https://doi.org/10.3382/japr.2013-00761>
- Papazyan TT, Denev SA, Surai PF. Selenium in poultry nutrition: lessons from research and wild nature. *Krmiva* 2006;48(5):275283.
- Parveen A, Khan SH, Khawaja T, *et al.* Growth performance and haemato-biochemical parameters of different breeds of rural chickens. *Journal of World's Poultry Research* 2017;7(3):114-22.
- Payne RL, Southern LL. Changes in glutathione peroxidase and tissue selenium concentrations of broilers after consuming a diet adequate in selenium. *Poultry Science* 2005;84(8):1268-76. <https://doi.org/10.1093/ps/84.8.1268>
- Peric L, Milosevic N, Zikic D, *et al.* Effect of selenium sources on performance and meat characteristics of broiler chickens. *Journal of Applied Poultry Research* 2009;18(3):403-9. <https://doi.org/10.3382/japr.2008-00017>
- Qiu K, Zheng JJ, Obianwuna UE, *et al.* Effects of dietary selenium sources on physiological status of laying hens and production of selenium-enriched eggs. *Frontiers in Nutrition* 2021;8:726770. <https://doi.org/10.3389/fnut.2021.726770>



- Rashidi AA, Ivani YG, Khatibjoo A, *et al.* Effects of dietary fat, vitamin E and zinc on immune response and blood parameters of broiler reared under heat stress. *Research Journal of Poultry Sciences* 2010;3(2):32-8. <https://doi.org/10.3923/rjpscience.2010.32.38>
- Rehman MS, Mahmud A, Mehmood S, *et al.* Blood biochemistry and immune response in Aseel chicken under free range, semi-intensive, and confinement rearing systems. *Poultry Science* 2017;96(1):226-33. <https://doi.org/10.3382/ps/pew278>
- Rotruck JT, Pope AL, Ganther HE, *et al.* Selenium: biochemical role as a component of glutathione peroxidase. *Science* 1973;179(4073):588-90. <https://doi.org/10.1126/science.179.4073.588>
- Ryu YC, Rhee MS, Lee KM, *et al.* Effects of different levels of dietary supplemental selenium on performance, lipid oxidation, and color stability of broiler chicks. *Poultry Science* 2005;84(5):809-15. <https://doi.org/10.1093/ps/84.5.809>
- Sahin K, Sahin N, Kucuk O, *et al.* Role of dietary zinc in heat stressed poultry: a review. *Poultry Science* 2009;88(10):2176-83. <https://doi.org/10.3382/ps.2008-00560>
- Sahin N, Onderci M, Sahin K, *et al.* Supplementation with organic or inorganic selenium in heat-distressed quail. *Biological Trace Element Research* 2008;122(3):229-37. <https://doi.org/10.1007/s12011-007-8075-6>
- Saleh AA, Ebeid TA. Feeding sodium selenite and nano-selenium stimulates growth and oxidation resistance in broilers. *South African Journal of Animal Science* 2019;49(1):176-83. <https://doi.org/10.4314/sajas.v49i1.20>
- Schwarz K, Foltz CM. Selenium as an integral part of factor 3 against dietary necrotic liver degeneration. *Journal of the American Chemical Society* 1999;115(3):254-6. [https://doi.org/10.1016/S0899-9007\(98\)00183-X](https://doi.org/10.1016/S0899-9007(98)00183-X)
- Sevcikova S, Skrivan M, Dlouha G, *et al.* The effect of selenium source on the performance and meat quality of broiler chickens. *Czech Journal of Animal Science* 2018;51(10):449-57. <https://doi.org/10.17221/3964-CJAS>
- Shafiq M. Comparative phenotypic characterization, performance, antibody response in naked neck chicken [dissertation]. Lahore (PK): University of Veterinary and Animal Sciences; 2016.
- Shimizu N, Hosogi N, Hyon GS, Jiang S, *et al.* Reactive oxygen species (ROS) generation and ROS-induced lipid peroxidation are associated with plasma membrane modifications in host cells in response to AK-toxin I from *Alternaria alternata* Japanese pear pathotype. *Journal of General Plant Pathology* 2006;72:6-15. <https://doi.org/10.1007/s10327-005-0245-9>
- Siegel HS. Stress, strain and resistance. *British Poultry Science* 1995;36(1):3-22. <https://doi.org/10.1080/00071669508417748>
- Skrivan M, Marounek M, Englmaierová M, *et al.* Influence of dietary vitamin C and selenium, alone and in combination, on the composition and oxidative stability of meat of broilers. *Food Chemistry* 2012;130(3):660-4. <https://doi.org/10.1016/j.foodchem.2011.07.103>
- Sluis WVD. Selenium yeast benefits in poultry production. *World's Poultry Science* 2007;8:15.
- Suchy P, Strakova E, Herzig I. Selenium in poultry nutrition: a review. *Czech Journal of Animal Science* 2018;59(11):495-503. <https://doi.org/10.17221/7730-CJAS>
- Surai PF. Natural antioxidants in avian nutrition and reproduction. Nottingham: Nottingham University Press; 2002.
- Surai PF. Selenium in nutrition and health. Nottingham: Nottingham University Press; 2006.
- Surai PF, Fisinin VI. Selenium in poultry breeder nutrition: An update. *Animal Feed Science Technology* 2006;191:1-15. <https://doi.org/10.1016/j.anifeedsci.2014.02.005>
- Surai PF, Fisinin VI. Antioxidant prooxidant balance in the intestine: Applications in chick placement and pig weaning. *Journal of Veterinary Science and Medicine* 2015;3(1):16. <https://doi.org/10.13188/2325-4645.1000011>
- Surai PF. Antioxidant systems in poultry biology: Superoxide dismutase. *Animal Nutrition* 2016a;1(1):8. <https://doi.org/10.21767/2572-5459.100008>
- Surai PF, Fisinin VI. Vitagenes in poultry production: Part 2. Nutritional and internal stresses. *World's Poultry Science Journal* 2016b;72(4):761-72. <https://doi.org/10.1017/S0043933916000726>
- Swain BK, Johri TS, Majumdar S. Effect of supplementation of vitamin E, selenium and their different combinations on the performance and immune response of broilers. *British Poultry Science* 2000;41(3):287-92. <https://doi.org/10.1080/713654938>
- Tiwary AK, Stegelmeier BL, Panter KE, James LF, *et al.* Comparative toxicosis of sodium selenite and selenomethionine in lambs. *Journal of Veterinary Diagnostic Investigation* 2006;18(1):61-70. <https://doi.org/10.1177/104063870601800108>
- Vasanthi SKI, Srividya G, Naidu G, *et al.* Heat stress and its mitigation strategies: a review. *Journal of Veterinary Science & Research* 2016;1(1):1-5. <https://doi.org/10.23880/OAJVSR-16000102>
- Wang YB, Xu BH. Effect of different selenium source (sodium selenite and selenium yeast) on broiler chickens. *Animal Feed Science and Technology* 2008;144(3-4):306-14. <https://doi.org/10.1016/j.anifeedsci.2007.10.012>
- Wang YX, Zhan XA, Yuan D. Effects of selenomethionine and sodium selenite supplementation on meat quality, selenium distribution and antioxidant status in broilers. *Czech Journal of Animal Science* 2011;56(7):305-13. <https://doi.org/10.17221/1296-CJAS>
- Wang CL, Xing GZ, Wang LS, *et al.* Effects of selenium source and level on growth performance, antioxidative ability and meat quality of broilers. *Journal of Integrative Agriculture* 2021;20(1):227-35. [https://doi.org/10.1016/S2095-3119\(20\)63432-3](https://doi.org/10.1016/S2095-3119(20)63432-3)
- Yuan D, Zhan XA, Wang YX. Effects of selenium sources and levels on reproductive performance and selenium retention in broiler breeder, egg, developing embryo, and 1-day-old chick. *Biological Trace Element Research* 2011;144(1-3):705-14. <https://doi.org/10.1007/s12011-011-9111-0>
- Zhang J, Wang X, Xu T. Elemental selenium at nano size (Nano-Se) as a potential chemopreventive agent with reduced risk of selenium toxicity: comparison with se-methylselenocysteine in mice. *Toxicological Sciences* 2008;101(1):22-31. <https://doi.org/10.1093/toxsci/kfm221>
- Zheng S, Zhao J, Xing H, *et al.* Oxidative stress, inflammation, and glycometabolism disorder-induced erythrocyte hemolysis in selenium-deficient exudative diathesis broilers. *Journal of Cellular Physiology* 2019;234(9):16328-37. <https://doi.org/10.1002/jcp.28298>
- Zhou X, Wang Y. Influence of dietary nano elemental selenium on growth performance, tissue selenium distribution, meat quality, and glutathione peroxidase activity in Guangxi Yellow chicken. *Poultry Science* 2011;90(3):680-6. <https://doi.org/10.3382/ps.2010-00977>





## \*AFFILIATIONS

- <sup>i</sup> Department of Poultry Science, Faculty of Animal Production and Technology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
- <sup>ii</sup> Department of Livestock and Dairy Development (Extension), KPK – Pakistan.
- <sup>iii</sup> Institute of Continuing Education and Extension, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
- <sup>iv</sup> Department of Pathology, Faculty of Veterinary Science, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
- <sup>v</sup> Faculty of Veterinary and Animal Sciences, The Islamia University of Bahawalpur-63100, Pakistan.
- <sup>vi</sup> Livestock and Dairy Development Department, Poultry Research Institute, Rawalpindi-46300, Pakistan.
- <sup>vii</sup> Department of Animal Nutrition, Faculty of Veterinary & Animal Sciences, The Islamia University of Bahawalpur, Bahawalpur-63100, Pakistan.
- <sup>viii</sup> Department of Biochemistry, Institute of Biochemistry, Biotechnology and Bioinformatics, The Islamia University of Bahawalpur, Bahawalpur-63100, Pakistan.
- <sup>ix</sup> Department of Zoology, Faculty of Chemical and Biological Sciences, The Islamia University of Bahawalpur, Bahawalpur-63100, Pakistan.
- <sup>x</sup> Department of Zoology, Faculty of Biosciences, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
- <sup>xi</sup> Department of Theriogenology, Faculty of Veterinary Science, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
- <sup>xii</sup> Department of Livestock Management, Faculty of Animal Production and Technology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
- <sup>xiii</sup> Department of Animal Production, Riphah College of Veterinary Sciences, Lahore-54000, Pakistan.
- <sup>xiv</sup> Department of Poultry Production, Shaheed Benazir Bhutto University of Veterinary and Animal Sciences, Sakrand-67210, Pakistan.
- <sup>xv</sup> Department of Animal Nutrition, Faculty of Animal Production and Technology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.

