



■ Author(s)

Uğurlu M¹
Akdağ F¹
Teke B¹
Salman M¹

¹ Ondokuz Mayıs University Veterinary Faculty Department of Animal Breeding and Husbandry, 55220 Atakum, Samsun, Turkey

¹ Ondokuz Mayıs University Veterinary Faculty Department of Animal Nutrition and Nutritional Disease, 55220 Atakum, Samsun, Turkey

■ Mail Address

Corresponding author e-mail address
Mustafa Uğurlu
Faculty of Veterinary Medicine, Ondokuz
Mayıs University, Kurupelit Campus,
Atakum, Samsun, Turkey.
Tel: +90-362-3121919
Email: mustafaugurlu55@hotmail.com

■ Keywords

Embryonal mortality, hatchability, pheasant, protein levels, sex ratio.

Effects of Protein in Diet and Sex Ratio on Egg Production, Egg and Hatching Chick Weight, Fertility, Hatchability and Embryonal Mortality in Pheasants (*Phasianus Colchicus*)

ABSTRACT

This study was performed to investigate the effects of crude protein (CP) in diet and sex ratio on egg production, egg and hatching chick weight and hatchability traits in pheasants (*Phasianus Colchicus*). In this study, the treatment groups were constructed as 15% CP-5:1; 15% CP-7:1; 18% CP-5:1; 18% CP-7:1 for CP and sex (male:female) ratio respectively. The eggs were obtained from 48 weeks old pheasant. In this study, it was determined that there was positive effective on egg production, egg and hatching weight and hatchability of fertile eggs (HFE) for 18% CP. In contrast, fertility (FR) ($p < 0.05$) and hatchability rate of total eggs (HR) ($p > 0.05$), EPEM ($p > 0.05$) and MPEM ($p > 0.05$) tended to decrease with increasing CP. It was found that there was egg production, egg ($p > 0.05$) and hatching weight ($p > 0.05$), FR ($p > 0.05$), HR ($p > 0.05$) and HFE ($p > 0.05$) increased with increasing female for sex ratio. In terms of embryonic mortality, the EPEM ($p > 0.05$) and LPEM ($p > 0.05$) rate decreased with increasing female. For protein levels \times sex ratio, it was found that hatchability traits were adversely affected for 15% CP-1:7. Also, it was determined that the lowest EPEM, MPEM and LPEM were determined for 7:1 in 18% CP group. Based on these results, to improve hatchability traits and a lower embryonic mortality rate, it may be beneficial to use 15% CP-5:1 or 18% CP-7:1 for pheasant breeding.

INTRODUCTION

In poultry breeding, high fertility, hatchability and low embryonic mortality are the main goal for profitability and productivity. Fertility, hatchability and embryonic mortality can be influenced by egg weight (Sczerbinska *et al.*, 1999, Petek *et al.*, 2003, Abiola *et al.*, 2008, Caglayan *et al.*, 2009), length of storage period (Fasenko *et al.*, 2001, Tona *et al.*, 2004, Oral Toplu *et al.*, 2007), age of flock (Elibol *et al.*, 2002, Seker *et al.*, 2004, Sari *et al.*, 2010), breeding type (Ozbey & Esen 2007), composition of hen diet (Cufadar *et al.*, 2010, Felipe *et al.*, 2010, Mohiti-Asli *et al.*, 2012, Praes *et al.*, 2014) and sex ratio (Ipek *et al.*, 2004, Alsobayel & Albadry 2012, Narinc *et al.*, 2013) in poultry species.

The composition of hen diet has an effective factor on egg production (Khajali *et al.*, 2008; Perez-Bonilla *et al.*, 2012), egg weight (Whitehead *et al.*, 1991, Zimmerman 1997; Shim *et al.*, 2013) and hatching characters (Danicke *et al.*, 2000) in poultry.

The protein level of hen diet is one of the factors influencing egg production and egg weight. Some researchers reported the general tendency increase egg production and egg weight with the increase of protein levels in the diets (Gunawardana *et al.*, 2008; King'ori *et al.*, 2010; Mohiti-Asli *et al.*, 2012; Shim *et al.*, 2013). Some other researchers reported that egg production and egg weight was not affected by feeding low crude protein (Cho *et al.*, 2004; Khajali *et al.*, 2008). In



poultry breeding, it is known that energy levels of the diet is important as well as protein levels of the diet. Leeson & Caston (1996) reported that the egg weight decreased when dietary protein content was reduced from 16.80% to 14.40% in isoenergetic hen diets. Li *et al.*, (2013) reported ranges of estimated optimum ME and CP intake were 2591 to 2683 kcal/kg and 15.58% to 16.64% CP for optimizing egg production of Lohmann Brown laying hens. Also, it was reported that diets with high protein and low energy have a negative effect on egg weight and egg production (Li *et al.*, 2013; Steenhuisen & Gous, 2016).

It was reported that there is a positive relationship between egg weight and hatching weight (Özcan *et al.*, 2001; Tona *et al.*, 2004). Therefore, the relationship between egg weight and protein levels of hen diet should be taken into consideration for hatching chick weight. Spratt & Leeson (1987) reported that a reduction in egg weight and consequently in chick weight decrease when dietary protein content was reduced from 16.70% to 12.70% for broiler breeder hens.

The protein levels of breeder diet is one important factor for hatching characters and embryonal mortality. A study with chukar partridge reported the general tendency increase infertility with high protein levels (CP 17%), and in contrary hatchability decreased under the same condition (Cufadar *et al.*, 2010). However, Mohiti-Asli *et al.*, (2012) found that fertility, hatchability of total and fertile eggs, and embryonal mortality of hen diet with high protein (CP 17.40%) were higher than those of hen diet with low protein (CP 14.50%). Some researchers reported that low crude protein (CP 12%) was not effective on fertility (Alsobayel, 1992) and hatchability (King'ori *et al.*, 2010) for local strains.

Sex ratio (male:female) is an important factor for fertility and hatchability. Generally, fertility tends to decrease with increasing sex ratio. Alsobayel & Albadry (2012) reported that sex ratio is efficient on fertility and hatchability of total eggs but not fertile hatchability and embryonal mortality in laying hens. Most researchers reported fertilization and hatching characteristics to decrease with increasing male: female ratio (Deeming & Wadland, 2002; Seker *et al.*, 2004; Narinc *et al.*, 2013). It was recommended that mating ratio of 1:1 and 1:2 produced the highest fertility and hatching characters in japanese quail (Seker *et al.*, 2004; Narinc *et al.*, 2013). Deeming & Wadland (2002) reported that a mating ratio of 8:1 produced a higher rate of egg production, fertility and hatchability compared with a 12:1 ratio.

As in the breeding of all poultry species, the effects of egg weight and length of storage period (Kirikci

et al., 2005, Demirel & Kirikci, 2009, Caglayan *et al.*, 2010), age of flock (Esen *et al.*, 2010; Ozbey *et al.*, 2011) and breeding type (Kirikci *et al.*, 2003; Genc & Ozbey, 2013) on successful hatching are among the most important influences for pheasants. However, there is little published work concerning the effect of protein levels (Monetti *et al.*, 1981) and sex ratio (Deeming & Wadland, 2002) on fertility, hatchability and embryonal mortality.

The National Research Council (NRC, 1994) reported the dietary metabolizable energy and crude protein requirements for breeder pheasants as 2800 kcal/kg ME and 15% CP, also other studies (Rhône Poulenc, 1993) recommended as 2900 kcal/kg ME and 17% CP for breeder pheasant and partridge. Therefore, it is important to differentiate the extent to which changes in hatchability traits are due to changes in protein levels of diet and sex ratio in pheasant.

As a contribution to efforts to increase the efficiency of pheasant breeding, the current study aimed to investigate the effects of the protein levels in the diets and sex ratio on hatchability and embryonal mortality.

MATERIAL AND METHODS

Animals, housing and designation of groups

A total of 360 eggs laid by 48 weeks old pheasant were obtained from the Samsun Gelemen Pheasant Breeding Centre of the Forest and Water Ministry of Turkey. In this study, the nutrition groups were classified as 15% CP and 18% CP. Sex ratio groups were designed as 5:1 and 7:1. After these classifications, a total of 4 groups were constructed, with each nutrition group divided into 2 subgroups for sex ratio. The composition of breeder diet was prepared according to the National Research Council (NRC, 1994). Formulation of breeder diet was performed by using to Microsoft Office Excel in the department of nutrition and nutritional disease. The chemical composition (protein, fat, dry matter and ash) of the experimental basal diets were analyzed according to the Association of Official Analytical Chemists (AOAC, 2000). Ingredient and chemical composition of experimental basal diets were presented in Table 1. The ration was applied on the breeding pheasants during the 4 weeks before laying period. The eggs were collected from small breeding flocks having 1:5 and 1:7 kept in open cages of 4 m x 5 m. The collected pheasant eggs were placed in trays and kept at 18 °C for 7 days. The hatching weights of the eggs were determined by measuring the eggs of all groups one by one in electronic scales delicate to 5g before hatching.



Table 1 – Ingredient and chemical composition of experimental basal diets.

Ingredients (%)	Protein levels	
	15% CP	18% CP
Corn	52.70	46.40
Wheat	19.50	13.50
Soybean (44% CP)	15.00	23.40
Sunflower meal (28% CP)	4.90	7.00
Vegetable oil	0.10	2.10
Dicalcium phosphat	1.80	1.71
DL-methionin	0.08	-
Calcium carbonat	5.25	5.25
Cocsiostat	0.03	0.05
Antioxidant	0.04	0.04
L-lizin hidroklorid	0.03	-
Salt	0.30	0.30
Vitamin-Mineral	0.27	0.25
Energy:protein	180.30	153.42
Analyzed value		
Dry matter %	89.60	90.01
Crude protein %	15.49	18.27
Crude fiber %	5.00	5.08
Ether extract %	3.90	5.00
Total ash %	10.18	9.55
Calculated values		
Metabolizable energy, kcal/kg	2793	2803
Calcium %	2.50	2.46
Available phosphorus %	0.40	0.38
Lysine %	0.68	0.71
Methionine+cysteine %	0.59	0.60

CP: Crude Protein

Hatchery conditions, hatching chick weight, evaluation of incubation results

The selected pheasant eggs were placed in trays and kept at 18 °C and 75% moisture for 7 days, based on their collection days, and then transferred to a cupboard type incubator. The eggs were incubated at 37.7°C and 65% moisture for 21 days in the development section, and at 37.5°C and 90% moisture for the last 3 days (Cetin & Kirikci, 2000). At the end of the 24th day of the incubation period, with monitoring of the hatchings, hatching weights of the chicks determined by measuring the chicks of all groups, one by one in electronic scales delicate to 5g when the chick's bodies dried up after the hatching. Also, the eggs which had not hatched were broken one by one and observed with the naked eye. In that macroscopic examination, the stage of embryo development at death was classified in terms of 3 possible death periods. The classification was done as follows; EPEM: early period embryonic mortality (embryo developed, filling the eggshell, eyes developed); MPEM: middle period embryonic mortality (feathers developed, more of yolk sack external to the

body) and LPEM: late period embryonic mortality (2/3 or whole of yolk sack in the body of embryo) (Akinci *et al.*, 2000).

FR, HR and HFE were calculated as the fertile eggs / total eggs, chick number of hatched / total eggs and chick number of hatched / fertile eggs respectively. Also, EPEM, MPEM and LPEM were calculated as early period embryo mortality / fertile eggs, middle period embryo mortality / fertile eggs and late period embryo mortality / fertile eggs respectively (Aksoy, 1994).

Statistical Analysis

Least square variance analysis was performed for the comparison of egg and hatching weight in the different protein levels, sex ratio and protein levels × sex ratio groups, and determination of the significance of differences between the groups was done with the Duncan test. The Chi-square test was used for the comparison of FR, HR, HFE, EPEM, MPEM and LPEM values for the different protein levels and sex ratio groups (SPSS 1993).

RESULTS

Egg weight and hatching chick weight

The means of egg and hatching chick weight for protein levels and sex ratio groups are presented in Table 2. The egg and hatching chick weight for 15% CP group were lower than those of 18% CP. In respect of sex ratio, the egg and hatching chick weight for 5:1 group were lower than those of 7:1. Differences among egg weights ($p < 0.01$) and hatching chick weight ($p < 0.001$) for protein levels were significant while they were not significant for sex ratio ($p > 0.05$) groups. For the diet of protein levels × sex ratio groups, it was determined that the highest egg weight was for 7:1 in 18% CP, but the highest hatching chick weight

Table 2 – Egg and hatching chick weight by groups (Mean ± S.E.)

Groups	Egg weight(gr)	n	Hatching weight(gr)	n
Protein levels	**		***	
15 %	32.73±0.16	178	22.32±0.17	128
18 %	33.43±0.15	182	23.22±0.17	129
Sex ratio	NS		NS	
5:1	32.87±0.16	173	22.59±0.17	127
7:1	33.29±0.15	187	22.98±0.17	130
Protein levels × sex ratio				
15 %-5:1	32.80±0.23	85	21.92±0.24	66
15 %-7:1	32.65±0.22	93	22.72±0.24	62
18 %-5:1	32.93±0.22	88	23.25±0.24	61
18 %-7:1	33.92±0.22	94	23.19±0.23	68

** : $p < 0.01$, *** : $p < 0.001$, NS: not significant



was 5:1 in 18% CP. In this result, it was determined that protein levels were effective on egg and hatching chick weight but, sex ratio were not effective on them.

Egg production, fertility and hatchability

The mean values for egg production, fertility and hatchability of pheasant eggs for feeding system, sex ratio and diet of protein level × sex ratio interaction are given in Table 3. Egg production was determined as 427 and 450 eggs for 15% CP and 18% CP respectively. It was observed that there was egg production increased with increasing protein level. The highest FR ($p < 0.05$) and HR ($p > 0.05$) were determined for the 15% CP. However, the highest HFE ($p > 0.05$) were for the 18% CP.

In terms of sex ratio, egg production was found as 430 and 447 eggs for 5:1 and 7:1 respectively. It was determined that there was egg production increased with increasing female. The FR, HR and HFE for 5:1 group were determined higher than those of 7:1 group but, there were no significant differences for FR ($p > 0.05$), HR ($p > 0.05$) and HFE ($p > 0.05$).

For the diet of protein levels × sex ratio groups, it was determined that egg production increased with increasing protein level and sex ratio. The highest FR and HR were determined for 5:1 group in 15% CP whereas the highest HFE was determined for 7:1 group in 18% CP group.

Embryonic mortality

The mean values of embryonic mortality for protein levels in diet, sex ratio and protein level in diet × sex ratio interaction are given in Table 4. The total embryonal mortality rate for 15% CP group (16.84%) higher than that of 18% crude protein (10.42%). In the present study, embryonal mortality decreased with increasing

CP for EPEM ($p > 0.05$) and MPEM ($p > 0.05$) whereas embryonal mortality increased for LPEM ($p > 0.05$).

In terms of sex ratio, the EPEM ($p > 0.05$) and LPEM ($p > 0.05$) rate for 5:1 group higher than those of 7:1 group, whereas the highest MPEM ($p > 0.05$) was for 7:1 group.

For interaction groups, it was determined that the lowest EPEM, MPEM and LPEM was determined for 7:1 in 18% CP group.

DISCUSSION

Egg weight and hatching chick weight

Many researchers reported that there were various effects in protein levels of hen diet on the egg weight in poultry species (Cho *et al.*, 2004; Khajali *et al.*, 2008; Gunawardana *et al.*, 2008; Kingori *et al.*, 2010; Shim *et al.*, 2013). Also, it was reported that there is a positive relationship between egg weight and hatching chick weight (Ozcan *et al.*, 2001; Tona *et al.*, 2004). Therefore, the relationship between egg weight and protein levels in hen diet is important for hatching chick weight.

In this study, mean egg weight and hatching chick weight was 32.73 and 22.32 g for 15% CP; 33.43 and 23.22 g for 18% CP, 32.87 and 22.59 g for 5:1; 33.29 and 22.98 g for 7:1 sex ratio respectively (Table 2). In the present study, mean egg weights were similar to those reported by Krystianiak *et al.*, (2005) namely, 31.33 g, and by Kozuszek *et al.*, (2009) namely 32.94 g respectively. When based on the 15% CP egg weight and hatching chick weight, increasing of egg weight and hatching chick weight were determined as 2.13% and 4.03% parallel to increasing crude protein percentage in this study respectively (Table 2). Likewise, many researchers reported the positive effects of dietary protein on egg weight (Gunawardana *et al.*,

Table 3 – Mean hatchability traits for protein levels, sex ratio and protein levels × sex ratio of pheasant eggs

Protein levels	Egg production(n)	Total eggs (n)	Fertile eggs (n)	Chick number (n)	FR (%)	HR (%)	HFE (%)
15 %	427	178	158	128	88.76	71.91	81.01
18 %	450	182	148	129	81.31	70.87	87.16
Chi-square					3.912*	0.047 ^{NS}	2.149 ^{NS}
Sex ratio							
5:1	430	173	150	127	86.70	73.41	86.70
7:1	447	187	156	130	83.42	69.51	83.42
Chi-square					0.760 ^{NS}	0.666 ^{NS}	0.751 ^{NS}
15 %-5:1	211	85	73	66	90.68	77.60	85.70
15 %-7:1	216	93	85	62	87.10	66.70	76.60
18 %-5:1	219	88	73	61	82.95	69.31	83.56
18 %-7:1	231	94	75	68	79.78	72.34	90.66

FR: Fertility, HR: Hatchability Rate, HFE: Hatchability of Fertile Eggs

*: $p < 0.05$, NS: not significant



2008; King'ori *et al.*, 2010; Shim *et al.*, 2013). Also, Spratt & Leeson (1987) reported that an increase in egg weight and consequently in chick weight increased when protein content in isoenergetic diet was changed from 12.70% to 16.70% for broiler breeder hens. The results of this studies indicated that egg and hatching chick weight increased with increasing crude protein in isoenergetic diet.

In laying poultry, there is scarce information about the effects of sex ratio on egg and hatching chick weight. In this study, it was found that there was increase in egg and hatching weight when mating ratio was changed from 1:5 to 1:7 for pheasant. Namely, egg and hatching weight for 7:1 were determined 1.28%, 1.73% higher than those of 5:1 respectively (Table 2). Likewise, a study with silver pheasant (*Lophura nychemera*) reported that an increase in male-female ratio from 1:2 to 1:5 mating ratio showed a positive effect on egg weight (Kaleem *et al.*, 2015). However, these results are lacking explanation for this effect.

For the protein levels of diet × sex ratio groups, some studies reported that diet with high protein and adequate energy had positive effect on egg weight and hatching chick weight (Gunawardana *et al.*, 2008; King'ori *et al.*, 2010; Shim *et al.*, 2013). However, some other studies found that egg weight was not affected by feeding low crude protein diets (Cho *et al.*, 2004; Khajali *et al.*, 2008). These conflicting results may be attributed to the different management and feeding conditions. Leeson & Caston (1996) reported that egg weight increased when dietary protein was increased in isoenergetic hen diets. Also, it was reported that diets with high protein and low energy have negative effect on egg weight. In this study, the highest egg weight and hatching chick weight was determined 18% CP for both of 1:5 and 1:7 sex ratio (Table 2). This result supports that protein levels were effective on egg and hatching weight but, sex ratio was no effective on them.

Egg production, fertility, hatchability and embryonal mortality

In poultry breeding, it has been reported that diet of protein and energy levels are effective factors on egg production (Li *et al.*, 2013; Steenhuisen & Gous, 2016). Some studies reported that egg production increased when dietary protein content was increased (Leeson & Caston, 1996; Praes *et al.*, 2014), also another study reported that diets with low protein exhibit negative effect on egg production (Keshawaraz 1998; Cufadar

et al., 2010) in isoenergetic hen diets. However, it was reported that diets with high protein and low energy have a negative effect on egg production (Li *et al.*, 2013; Steenhuisen & Gous, 2016). In this study, hen diets with different protein levels (15%; 18%) and isoenergetic (2800 kcal/kg) were used. It was determined 5.38% increase in egg production parallel to increasing protein level in this study. Therefore, increasing egg production might be related to the protein level of the diet in this study.

The sex ratio is recommended for commercial pheasant range from 1:7, 1:10 or 1:12 (Deeming & Wadland, 2002). The sex ratio designated in this study were 1:5 and 1:7. In this study, egg production was found as 430 and 447 eggs for 5:1 and 7:1 respectively. It was observed that egg production for 5:1 was lower than that for 1:7 in this study. The low egg production of 5:1 could be due to decreasing of their female number.

For the diet of protein levels × sex ratio groups, in this study, it was determined that egg production increased with increasing protein levels for both protein levels and sex ratio. The highest egg production was found in 18%-7:1 group. It is well known that hen diets with high protein and adequate energy have positive effects on egg production (Leeson & Caston 1996; Praes *et al.*, 2014). It was reported that the egg production increased with female increasing (Deeming & Wadland 2002). This results supported that protein levels of diet and sex ratio had a synergistic effect on egg production.

In the present study, it was determined that the highest FR and HR was determined for 15% CP while the lowest HFE was determined for 15% CP (Table 3). Moreover, the total embryonal mortality rate for 15% CP (16.84%) was higher than that of 18% CP (10.42%) (Table 4). Especially, it was determined that MPEM for 15% CP was higher than that of 18% CP in this study. Supporting this finding, a study with pheasants reported that higher fertility and hatchability was determined in pheasants fed with ration with the poorest protein content (Monetti, 1981). Likewise, Cufadar & Bahtiyarca (2006) reported that diets with low protein (13%), conventional (17%) and higher protein (20%) had no different effect on fertility and hatching characters of breeder chukar partridge. However, another study with chukar partridge reported the general tendency increase infertility with high protein levels (CP 17%), in contrary hatchability decreased under the same condition (Cufadar *et al.*, 2010).



In this study, when the FR, HR and HFE were evaluated for sex ratio, the highest FR, HR and HFE were determined for 5:1 group in this study. Narinc *et al.*, (2013) reported as up to 3 female per male ratio for optimum fertility ratio for japan quails.

Table 4 – Embryonic mortality mean rates for protein levels, sex ratio and protein levels × sex ratio of pheasant eggs

Protein levels	EPEM	MPEM	LPEM
15 %	5.61	7.86	3.37
18 %	2.19	3.29	4.94
Chi-square	2.816 ^{NS}	3.580 ^{NS}	0.556 ^{NS}
Sex ratio			
5:1	4.62	3.46	5.20
7:1	3.20	7.48	3.20
Chi-square	0.488 ^{NS}	2.766 ^{NS}	0.895 ^{NS}
Protein levels × sex ratio			
15 %-5:1	7.05	3.52	2.35
15 %-7:1	4.30	11.82	4.30
18 %-5:1	2.27	3.40	7.95
18 %-7:1	2.12	3.19	2.12

EPEM: Early Period Embryonic Mortality, MPEM: Middle Period Embryonic Mortality, LPEM: Late Period Embryonic Mortality. NS: Non-significant

In this study, when the FR, HR and HFE were evaluated for sex ratio, the highest FR, HR and HFE were determined for 5:1 group in this study. Narinc *et al.*, (2013) reported as up to 3 female per male ratio for optimum fertility ratio for japan quails.

Female birds have larger number of males recruiting and decreasing mating ratio, therefore FR can be increased with decreasing for female:male. It was reported that the FR, HR and HFE for 8:1 was higher than those of 12:1 for pheasants (Deeming & Wadland 2002). Alsobayel & Albadry (2012) reported from 1:6 to 1:14 changed mating ratio hatchability of fertile eggs, early embryonal mortality late embryonal mortality and total embronal mortality decreasing for Baladi Chickens. In the present sudy, the HR and HFE for 5:1 were determined as 5.31% and 3.93% higher than those of 7:1 respectively. The total embryonic mortality was determined as 13.28% and 13.88% for 5:1 and 7:1 respectively (Table 4). However, the MPEM (7.48%) rates was determined for the 1:7. Therefore, the loss of HR and HFE for 7:1 were attributable to an increase in MPEM.

For the protein levels of diets× sex ratio groups, the highest FR and HR was determined for 15% CP and 1:5 sex ratio group while the highest HFE was determined for 18% CP and 1:7 (Table 3). The lowest embriyonal mortality was determined for 18% CP and 1:7 sex ratio group (Table 4). These results show that hatchability in both of protein levels and sex ratio is being determined by rates of fertility rather than embryonic mortality.

Consequently, in terms of protein levels of the diet, the FR and HR for 15% CP was higher than those of 18% CP. In contrast, HFE for 15% CP tended to decrease due to increasing total embryonic mortality. In this study, in terms of sex ratio, the FR, HR and HFE for 5:1 group was higher than those of 7:1. Total embryonic mortality had almost the same value for both sex ratios. However, it was observed that embryonic mortality for 7:1 was lower, except for MPEM. Based on these results, to achieve a higher hatchability rate and hatchability of fertile eggs, and a lower embryonic mortality rate, it may be beneficial to use 15% CP-5:1 or 18%CP-7:1 for pheasant breeding.

ACKNOWLEDGEMENTS

The authors thank the Gelemen Pheasant Breeding Centre of the Forest and Water Ministry of Turkey.

REFERENCES

- Abiola SS, Meshioye OO, Oyerinde BO, Bamghose MA. Effect of size on hatchability of broiler chicks. *Archivos de Zootecnia* 2008;57(217):83-86.
- Aksoy FT. Hen breeding. 2nd ed. Ankara: Şahin Press;1994.
- Akinci Z, Koçak S, Tekerli M, Akcan A. Relationship between egg weight loss rate and embryonic development during incubation in quail eggs. *Tavukçuluk Araştırma Enstitüsü Dergisi* 2001;3(1):31-35.
- Alsobayel AA. Effect of protein rearing diet and age on fertility and hatchability parameters of Saudi Arabian Baladi Chickens. *Journal King Saudia University Agriculture Science* 1992;4(1):47-54.
- Alsobayel AA, Albadry MA. Effect of age and sex ratio on fertility and hatchability of Baladi and Leghorn laying hens. *The Journal of Animal & Plant Science* 2012;22(1):15-19.
- AOAC. Association of Official Analytical Chemistry. Official methods of analysis of AOAC international. 17th ed. Maryland; 2000.
- Çağlayan T, Alaşahan S, Çetin O, Kırıkçı K, Günlü A. Effects of egg weight and length of storage period on chick weight and hatchability performance of pheasant (*Phasianus Colchicus*). *The Journal of Food, Agriculture and Environment* 2010;8:407-410.
- Çetin O, Kırıkçı K. Alternative poultry breeding. Pheasant-Partridge. Konya: Selçuk University Publishing; 2000.
- Cufadar Y, Bahtiyar Y. Effect of dietary protein and aminoacids content on the performance reproductive characteristics and nitrogen excretion in breeding chukar partridge (*Alectoris chukar*). *Selçuk Üniversitesi Ziraat Fakültesi Dergisi* 2006;20:129-136.
- Cufadar Y, Olgun O, Bahtiyar Y, Yıldız AO. Effects of dietary energy and protein on performance, reproduction traits and nitrogen-excretion of breeder chukar partridges (*Alectoris chukar*). *Revue de Médecine Veterinaire* 2010;161(4):151-156.
- Cho YM, Shin IS, Yang CJ. Effects of feeding dried leftover food on productivity on laying hens. *Asian-Australasian Journal of Animal Science* 2004;17:518-522.



- Danicke S, Halle I, Jeroch H, Böttcher W, Ahrens P, Zachmann R, *et al.*, Effect of soy oil supplementation and protein level in laying hen diets on praecaecal nutrient digestibility, performance, fatty acid composition of yolk fat and on other egg quality parameters. *European Journal of Lipid Science and Technology* 2000;102(3):218-232.
- Deeming DC, Wadland ED. Influence of mating sex ratio in commercial pheasant flocks on bird health and the production, fertility and hatchability of eggs. *British Poultry Science* 2002;43(1):16-23.
- Demirel S, Kırıkçı K. Effect of different egg storage times on some egg quality characteristics and hatchability of pheasants (*Phasianus Colchicus*). *Poultry Science* 2009;88:440-444.
- Elibol O, Peak SD, Brake J. Effect of flock age, length of egg storage and frequency of turning during storage on hatchability of broiler hatching eggs. *Poultry Science* 2002;81:945-950.
- Esen F, Özbey O, Genç F. The effect age on egg production, hatchability and egg quality characteristics in Pheasants (*Phasianuscolchicus*). *Journal of Animal and Veterinary Advances* 2010;9(8):1237-1241.
- Fasenko GM, Christensen VL, Wineland MJ, Petite JN. Examining the effects of pre storage incubation of turkey breeder eggs on embryonic development and hatchability of eggs stored for four or fourteen days. *Poultry Science* 2001;80:132-138.
- Felipe L, Santos EC, Tavian AF, Góes PAA, Moraes VMB, Tonhati H, *et al.*, Effect of crude protein levels and organic selenium supplementation in the diets fed during the breeding season on reproductive parameters of red-winged tinamous (*Rhynchotus rufescens*). *Brazilian Journal of Poultry Science* 2010;12(1):63-71.
- Gunawardana P, Roland Sr DA, Bryant MM. Effect of energy and protein on performance, egg components, egg solids, egg quality and profits in molted Hy-Line W36 Hens. *Journal of Applied Poultry Research* 2008;17:390-397.
- Genç F, Özbey O. Effect of various breeding systems on egg production and hatchability characteristics of pheasants (*Phasianus Colchicus*). *Firat University Journal of Health Science (Veterinary)* 2013;27(1):43-47.
- Ipek A, Şahan U, Yılmaz B. The effect of live weight, male to female ratio and breeder age on reproduction performance in Japanese quails (*Coturnix coturnix japonica*). *South African Journal of Animal Science* 2004;34(2):130-134.
- Kaleem M, Ruby T, Khan AA, Rafay M, Abdullah M. Impact of different mating sex ratios on reproductive efficiency of silver pheasant *Lophura nycthemera* (Galliformes, Phasianidae). *Russian Journal of Ecology* 2015;46(3):289-291.
- Kırıkçı K, Tepeli C, Günlü A, Çetin O. Production characteristics of pheasants (*Phasianus Colchicus*) in different breeding regimes. *Turkish Journal of Veterinary and Animal Sciences* 2003;27:907-910.
- Kırıkçı K, Günlü A, Garip M. Some quality characteristics of pheasant (*Phasianus colchicus*) eggs with different shell colors. *Turkish Journal of Veterinary and Animal Sciences* 2005;29:315-318.
- Khajali F, Khoshouie EA, Dehkodi SK, Hematin M. Production performance and egg quality of Hy-line W36 laying hens fed reduced-protein diets at a constant total sulfur aminoacid: lysine ratio. *Journal Applied of Poultry Research* 2008;17:390-397.
- Kingori AM, Tuitoek JK, Muiruri HK, Wachira AM. Effect of dietary crude protein levels on egg production, hatchability and post-hatch offspring performance of Indigenous chickens. *International Journal of Poultry Science* 2010;9(4):324-329.
- Krystianiak S, Kozuszek R, Kontecka H, Nowaczewski S. Quality and ultrastructure of eggshell and hatchability of eggs in relation to eggshell colour in pheasants. *Animal Science and Reports* 2005;23(1):5-14.
- Kozuszek R, Kontecka H, Nowaczowski S, Rosinski A. Storage time and eggshell colour of pheasant eggs vs. the number of blastodermal cells and hatchability results. *Folia Biologica Krakow* 2009;57(3-4):121-130.
- Leeson S, Caston LJ. Response of laying hens to diets varying in crude protein or available phosphorus. *Journal Applied Poultry Research* 1996;5:289-296.
- Li F, Zhang M, Wu XH, Li CY, Yang XJ, Dong Y, *et al.*, Effects of metabolizable energy and balanced protein on egg production, quality and components of Lohmann Brown laying hens. *Journal Applied Poultry Research* 2013;22:36-46.
- MohitiAsli M, Shivazad M, Zaghari M, Rezaizan M, Aminzadeh S, Mateos GG. Effects of feeding regimen, fiber inclusion and crude protein content of the diet on performance and egg quality and hatchability of eggs of broiler breeder hens. *Poultry Science* 2012;91:3097-3106.
- Monetti PG, Marcomini F, Monge F, Tinti P. Protein level in rations and reproduction performance in the pheasant [abstracts]. Rome: AGRIS; 1981.
- NRC. Nutrient requirements of poultry. 9th ed. Washington: National Academy Press; 1994.
- Narinc D, Aygun A, Sari T. Effects of cage type and mating ratio on fertility in Japanese quails (*Coturnix coturnix japonica*) eggs. *Agriculture Science Developments* 2013;2(1): 4-7.
- Oral Toplu HD, Dereli Fidan E, Nazlıgül A. The effects of weight and storage time of hatching egg on hatchability traits and hatching weight in Japanese quails. *Journal of Faculty of Veterinary Medicine, Erciyes University* 2007;4(1):11-16.
- Özbey O, Esen F. The effects of different breeding systems on egg productivity and egg quality characteristics of rock partridge. *Poultry Science* 2007;86:782-785.
- Özbey O, Esen F, Aysöndü MH. The effect of the age of the first egg-laying on the egg production, hatchability and egg quality of pheasants (*Phasianus colchicus*). *Journal of Animal and Veterinary Advances* 2011;10:3196-3200.
- Özcan M, Ekiz B, Güneş H. Effects of egg weight and hatching weight sizes on growth performance in the Japanese quail (*Coturnix coturnix japonica*). *The Journal of the Faculty of Veterinary Medicine, University of Istanbul* 2001;27(2):577-584.
- Perez-Bonilla AS, Novoa J, Garcia M, Mohiti-Asli M, Frika M, Matheos GG. Effects of energy concentration of the diet on productive performance and egg quality of Brown egg-laying hens differing in initial body weight. *Poultry Science* 2012;91:3156-3166.
- Petek M, Baspinar H, Ogan M. Effects of egg weight and length of storage on hatchability and subsequent growth performance of quail. *South African Journal of Animal Science* 2003;33(4):242-247.
- Praes MFFM, Junqueira OM, Pereira AA, Filardi RS, Duarte KF, Sgavioli S, Alva JCR, Dominiques CHP. High-Fiber diets with reduced crude protein for commercial layers. *Brazilian Journal of Poultry Science* 2014;16(2):43-50.
- Rhône Poulenc. Rhodimet feed formulation guide. 6th ed. Rhone Poulenc Animal Nutrition, Antony Cedex; 1993.
- Sarı M, Tilki M, Saatçi M, Işık S, Önk K. Effect of parental age, egg weight and shape index on hatchability traits and liveability in Japanese quail (*Coturnix coturnix japonica*). *Sağlık Bilimleri Veteriner Dergisi, Firat Üniversitesi* 2010;24(2):93-97.
- Şeker I, Kul S, Bayraktar M. Effects of parental age and hatching egg weight of Japanese Quails on hatchability and chick weight. *International Journal of Poultry Science* 2004;3(4):259-265.



Steenhuisen E, Gous RM. The response of broiler breeder hens to dietary balanced protein. *South African Journal of Animal Science* 2016;46(3):269-277.

Szczerbinska D, Zubrecki A. The quail egg weight and their storage period vs. hatching success and rearing performance. *Advances in Agricultural Sciences* 1999;6:91-100.

Shim MY, Song E, Billard L, Aggrey SE, Pesti GM, Sodsee P. Effects of balanced dietary protein levels on egg production and egg quality parameters of individual commercial layers. *Poultry Science* 2013;92:2687-2696.

Spratt RS, Leeson S. Broiler breeder performance in response to diet protein and energy. *Poultry Science* 1987;66(4):683-693.

SPSS. *Statistical Package in Social Science for Windows*. Chicago: Statistical Innovations; 1993b.

Tona K, Onagbesan O, De Ketelaere B, Decuypere E, Bruggemen V. Effects of age of broiler breeders and egg storage on egg quality, hatchability, chick quality, chick weight and chick posthatch growth to forty-two days. *Journal Applied Poultry Research* 2004;13(1):10-18.

Whitehead CC, Bowman AS, Griffin HD. The effects of dietary fat and bird age on the weights of eggs and egg components in the laying hen. *British Poultry Science* 1991;32:565-574.

Zimmerman RA. Management of egg size through precise nutrient delivery. *Journal of Applied Poultry Research* 1997;6:478-482.