



Sources of Human Non-Typhoid Salmonellosis: A Review

■ Author(s)

Freitas Neto OC de¹
Penha Filho RAC²
Barrow P³
Berchieri Junior A⁴

^{1,2,4} Universidade Estadual Paulista "Julio de Mesquita Filho" - campus de Jaboticabal
Departamento de Patologia Veterinária.
³ The University of Nottingham
School of Veterinary Medicine and Science

■ Mail Address

E-mail: oliveirocaetano@yahoo.com.br

■ Keywords

Antimicrobial resistance, beef, food, pork, poultry, *Salmonella* spp., vegetables,

ABSTRACT

Salmonellosis is a worldwide disease caused by bacteria of the genus *Salmonella*. Currently, there are over 2,500 identified serovars of *Salmonella*. A reduced number of these serovars, about eighty, are implicated in most animals and human diseases. Most cases of salmonellosis in humans are associated with the consumption of contaminated food products such as beef, pork, poultry meat, eggs, vegetables, juices and other kind of foods. It may also be associated with the contact between humans and infected pet animals. Therefore, the chain of human salmonellosis is very complex and in most cases the origin of the infection is difficult to establish. The use of antimicrobial agents to treat and to prevent bacterial infections in humans and animals, as well as as growth promoters in animal production, has favoured the selection and transference of resistance genes between different bacteria, including *Salmonella* serovars. Many studies have confirmed the role of foods of animal origin as a source of multi drug-resistant *Salmonella* serovars. For this reason, continuous surveillance of these pathogens along the food chain together with the responsible use of antimicrobial agents is necessary.

BACKGROUND

The bacteria of the genus *Salmonella* are spread worldwide and are responsible for illnesses in human beings and animals. Currently, the genus is divided into two species, *Salmonella enterica* and *Salmonella bongori* (Popoff & Minor 2001). *S. enterica* is divided into six subspecies (*enterica*, *salamae*, *arizonae*, *diarizonae*, *houtenae* and *indica*), each of which has several serovars or serotypes (Table 1). Thus, a serotype is nominated in the following way: *Salmonella enterica* subspecies *enterica* serotype Typhimurium, which may be simplified as *Salmonella* Typhimurium. Nowadays, more than 2,500 serotypes are known and most of them (almost 1,500) belong to subspecies *enterica* (Porwollik *et al.*, 2004; CDC, 2007c).

Table 1. Distribution of serovars by *Salmonella* species.

Species	<i>S. enterica</i>						<i>S. bongori</i>
	<i>enterica</i>	<i>salamae</i>	<i>arizonae</i>	<i>diarizonae</i>	<i>houtenae</i>	<i>indica</i>	
Number of Serovars	1490	500	94	329	72	12	22

Salmonellosis is the disease caused by *Salmonella* serovars. A few serotypes are host-specific. *Salmonella* Pullorum and Gallinarum are responsible for pullorum disease and fowl typhoid in poultry, respectively, *S. Typhi* for typhoid fever in human beings, *S. Dublin* for disease in bovine, etc. However, the others are not host-specific and may infect several animal species including human beings. These serotypes are generally responsible for food-borne diseases and foods of animal origin are the



main source. Historically, *S. Typhimurium* is the most common agent of human food-borne disease, although in the last few decades *S. Enteritidis* has become more common (Braden, 2006; Fernandes *et al.*, 2006). Among all the 2,500 serovars, about eighty are most frequently involved in animal and human salmonellosis. *S. Typhimurium* and *S. Enteritidis* are the most common agents of disease in human beings and animals, but there is also increasing concern about the *Infantis*, *Agona*, *Hadar*, *Heidelberg*, and *Virchow* *Salmonella* serotypes.

Salmonellosis is one of the most important human food-borne diseases. Improvements in the quality of life as a result of hygienic conditions and sewage treatment have helped to prevent typhoid fever, and its occurrence is now frequently related to bad weather conditions like storms resulting in contamination of drinking water with sewage. The world population has increased dramatically, and to address the demand for food it has been necessary to increase production in large scale intensive farms. In these conditions, enteric pathogens like *Salmonella* serovars that have non-specific hosts, once introduced on farms may easily become disseminated among animals, resulting in persistence. Once this happens, it becomes almost impossible to eradicate the pathogens from farms and to eliminate them from products used as food or for preparing foods. To give an idea of the complexity of the chain of food-borne salmonellosis, at the beginning of the 1970s, many cases of human salmonellosis caused by *Salmonella* *Agona* in Europe and in the United States of America were associated with the ingestion of broiler meat from flocks fed with feed containing fish meal from Peru (Clark *et al.*, 1973). Since the 1980s, worldwide outbreaks of human salmonellosis have been caused by *S. Enteritidis* present in eggs and contaminated broiler meal. In this case, breeders and laying hens were often thought to be infected by rodent feces and urine and transmitted the bacterium vertically, resulting in infected progeny. In addition, during the same period, outbreaks of human salmonellosis, resulting from ingestion of pork and beef meat contaminated with *Salmonella* serovars, have been reported in many countries (CDC, 1981; Spitalny *et al.*, 1984; Delpech *et al.*, 1998), with *S. Typhimurium* being the main serotype involved. Pork and beef were probably contaminated during the slaughter process followed by inadequate cooking.

Despite the strong link between food of animal origin and human salmonellosis, people may be infected by other ways, such as by cross contamination

at home and commercial kitchens, through contact with other people, pets - especially dogs, cats and reptiles, as well as by ingestion of vegetables and fruits. This review attempts to provide information about different sources of *Salmonella* serovars which may be involved in human salmonellosis outbreaks. In addition, a basic knowledge about antimicrobial resistance in *Salmonella* serovars isolated from foods of animal origin is also presented since many drugs are used frequently in animal production.

Red meat and red-meat products as a source of *Salmonella* serovars for human beings

Red meat and red-meat products are recognized as an important source of human salmonellosis. Only in 2007, for instance, beef and pork (and products thereof) contaminated with *Salmonella* serovars were responsible for 800 cases of human salmonellosis involving 37 outbreaks in Europe (EFSA, 2009).

Additionally, in 2005, Hess *et al.* (2008) reported that lamb's liver was responsible for an outbreak of *S. Typhimurium* phage type 197 in Australia. In Germany, from 2001 to 2005, epidemiological studies, microbiological testing, and trace-back investigations showed that pork and pork products were involved in human salmonellosis outbreaks (Jansen *et al.*, 2007). Moreover, in Italy, an outbreak of *S. Typhimurium* phage type DT 104A involving 63 cases was reported. In this study, a case-control study was conducted and data were analyzed by multivariate analysis. No food samples were available for testing; however, in northern Italy, two months prior to the outbreak, the veterinary surveillance system identified the first isolation of *S. Typhimurium* DT104A in a pig. Both human (patients) and pig isolates showed indistinguishable pulsed-field gel electrophoresis (PFGE) patterns. It was not possible to trace the pig after the sample collection. The results of the case-control study suggested that the consumption of pork salami was associated with this outbreak, underlining the importance of good manufacturing practices for ready-to-eat foods (Luzzi *et al.*, 2007).

In Denmark an outbreak of human salmonellosis caused by *S. Typhimurium* DT 104 occurred between July and August 2005, and the comparison of strains from patients and food using PFGE, tandem repeat analysis (MLVA), and antibiogram tests showed that the source of the outbreak was imported beef served as carpaccio (Ethelberg *et al.*, 2007). Some months later there was an increase in the number of *S.*



Typhimurium DT 104 cases in the Netherlands (Kivi *et al.*, 2007). These occurrences were found to be epidemiologically associated with the disease in Denmark. A case-control study was conducted with 56 cases and 100 controls, and logistic regression analysis was used for analyzing the results (Ethelberg *et al.*, 2007). From this investigation beef was considered the most probable vehicle of *S. Typhimurium* DT 104. Complementary assessment suggested that imported beef from a third European country had spread this *Salmonella* strain resulting in these outbreaks.

In the USA in 2004, an epidemiological investigation demonstrated that ground beef sold by a national chain of one supermarket was the source of food-borne salmonellosis due to *S. Typhimurium* in nine states. The PFGE patterns of *S. Typhimurium* isolates from patients and ground beef were found to be identical (CDC, 2006).

Many other reports involving human salmonellosis outbreaks associated with consumption of red meat have been recorded in the literature (CDC, 1995a; Davies *et al.*, 1996; Roels *et al.*, 1997; Shapiro *et al.*, 1999; Haeghebaert *et al.*, 2001). In most of cases, the disease was associated with the consumption of contaminated meat or as a result of incorrect or inadequate cooking.

Human non-typhoid salmonellosis caused by the ingestion of dairy products

Salmonella serovars are widespread in the environment and may contaminate a variety of food and food ingredients. Since dairy products are from animal origin, they may also be implicated in human food-borne salmonellosis. Usually milk and milk products are submitted to pasteurization, which kills *Salmonella* serovars. Milk-borne salmonellosis is therefore often related to the consumption of raw or inadequately pasteurized milk. However, *Salmonella* serovars may also contaminate dairy products after the pasteurization process.

S. Typhimurium caused an outbreak in Pennsylvania, USA, in 2007, through the consumption of raw milk. *S. Typhimurium* isolated from human and milk exhibited suspect identical PFGE patterns. Later, an epidemiological investigation confirmed that the source of *S. Typhimurium* was the raw milk and raw milk products from a dairy (CDC, 2007e). An outbreak of *S. Newport* in the USA in 2008 occurred predominantly among Hispanics in northeastern Illinois. *S. Newport* was recovered from patients and from a Mexican-style

aged cheese made with unpasteurized milk (CDC, 2008c). The strains isolated from people and cheese had indistinguishable PFGE patterns. Another outbreak of human salmonellosis associated with cheese was described in Switzerland, Europe, where a nationwide outbreak of gastrointestinal disease caused by *S. Stanley*, a rare serovar in Europe, occurred from September 2006 to February 2007. *S. Stanley* strain was isolated from a local soft cheese and from 94% of affected people. The withdrawal of the cheese from the market resolved the problem (Pastore *et al.*, 2008).

In view of the great number of human salmonellosis associated to the consumption of powdered milk products registered in the 1960's and 1970's, *Salmonella* control criteria were included in the Codex of Hygienic Practice for Foods for infants and children (Codex, 1979). Between 1985 and 2005 at least six outbreaks of salmonellosis, involving up to 250 infants, have been associated with powdered infant formula. For example, in 2005 in France, an outbreak affecting more than 100 infants was associated with powdered infant formula contaminated with *S. Agona*. Many of these outbreaks were identified because the *Salmonella* strains were unique in some way (e.g. a rare serovar). Another common feature of the outbreaks was the low level of *Salmonella* contamination detected in the implicated formula (*Salmonella* serovars may be missed in routine testing). These outbreaks likely represent only a small proportion of the actual number of *Salmonella* infections in infants that have been linked to powdered infant formula (Toyofuku *et al.*, 2006; Cahill *et al.*, 2008).

***Salmonella* serovars in foods from vegetable origin: a matter of increasing concern**

In recent years, the importance of foods of vegetable origin as potential vehicles of gastrointestinal infection has been highlighted. During production, storage or even in retail outlets, *Salmonella* serovars may contaminate vegetables, fruits, juices, and, as they are usually consumed raw, they may also be implicated in human salmonellosis.

Unpasteurized orange juice was responsible for food-borne salmonellosis in 152 people in six states in the USA between May and July 2005 (Jain *et al.*, 2009). Vojdani *et al.* (2008) reviewed fruit juice-associated outbreaks of illness reported to Centers for Disease Control and Prevention (CDC), in Atlanta, USA. From 1995 through 2005, 21 juice-associated outbreaks were



reported to CDC; ten implicated apple juice or cider, eight were linked to orange juice, and three implicated other types of fruit juice. These outbreaks caused 1,366 illnesses, with a average of 21 cases per outbreak (range, 2 to 398 cases). Among the 13 outbreaks of known etiology, five were caused by *Salmonella* serovars.

Three cases of *S. Thompson* infection were registered by the Norwegian reference laboratory in November 2004. The results of the investigation indicated that the consumption of rucola lettuce and mixed salad were responsible for the outbreak (Nygård *et al.*, 2008).

In addition, Little & Gillespie (2008) described general outbreaks of infectious intestinal disease linked to prepared salads reported in England and Wales from 1992 to 2006, as well as European outbreaks associated with prepared salads as a result of international trade. According to the authors, around 23% of these outbreaks were attributed to *Salmonella* serovars. Cross-contamination, infected food handler or inappropriate storage were the most commonly factors associated with vegetable contamination.

In 2002 tomatoes, grown and packed in Virginia state (USA), contaminated with *S. Newport*, caused illness in 510 patients in 26 other states (Kretsinger *et al.*, 2003). Later, in July-November 2005, the same strain (confirmed by PFGE) caused illness in at least 72 patients in 16 states of the USA. A case-control study during the 2005 outbreak confirmed tomatoes as the source. The tomatoes were traced back to Virginia, where the *S. Newport* strain was isolated from pond water used to irrigate tomato fields, suggesting persistent contamination of the fields (Greene *et al.*, 2008). In addition, during 2005-2006, three more outbreaks of *Salmonella* infections associated with eating tomatoes were detected in the USA and Canada. These outbreaks resulted in 387 culture-confirmed cases of salmonellosis, with isolation of *S. Newport*, *S. Braenderup*, and *S. Typhimurium* (CDC, 2007b).

Unicomb *et al.* (2005) described human salmonellosis outbreaks associated with sesame seed-based food. In November 2002, an outbreak of *S. Montevideo* occurred in New South Wales, Australia. Infected persons were interviewed, and epidemiologically linked retail outlets inspected. Imported "tahini", a sesame seed-based food, was rapidly identified as the source of infection. A second outbreak was identified in Australia in June-July 2003 and another in New Zealand in August 2003. Of a total of 68 patients

infected by *S. Montevideo*, 66 were contacted. Fifty-four (82%) reported consumption of sesame seed-based foods. Laboratory analyses demonstrated closely related PFGE patterns in the *S. Montevideo* isolates from human cases and sesame-based foods imported from two countries.

Although it is known that fresh salad vegetables, herbs, seeds and fruit may become contaminated from environmental sources, they have been associated only recently with food-borne diseases. Nowadays, the role of vegetables as vehicle for pathogens transmission has become clear and recurrent, demonstrating that major health problems can arise from consumption of contaminated vegetables if hygiene practices break down.

Pets as a source of *Salmonella* infection for human beings

Many pet animals can harbor and shed *Salmonella* serovars. The close contact between pets and people, especially children and immunosuppressed people, can result in human salmonellosis. Animals commonly infected with *Salmonella* serovars which may pose a risk to humans include amphibians, birds, cats, dogs, fish, guinea pigs, hamsters, horses, mice, rabbits, lizards, snakes, and turtles, although, according to literature reports, the last three (lizards, snakes, and turtles) are responsible for the majority of human salmonellosis outbreaks associated with pets (CDC, 2005; Bruins *et al.*, 2006; Corrente *et al.*, 2006; CDC, 2007f; Bertrand *et al.*, 2008; CDC, 2008a).

The association of reptiles with human salmonellosis was first described in the 1950s (Boycott *et al.*, 1953). Since then, reports have become more frequent. According to Mermin *et al.* (2004), approximately 1.4 million human cases of *Salmonella* infection occur each year in the USA and it has been estimated that 74,000 are the result of exposure to pet reptiles and amphibians. From 1994 to 1995, thirteen health state agencies in the USA reported to CDC human infection cases caused by uncommon *Salmonella* serovars, in which the patients had had direct or indirect contact with reptiles (snakes, lizards and turtles). In most cases, the same *Salmonella* serovar was isolated from the patient and from the reptile (CDC, 1995b). In 2005, at least six cases of human salmonellosis were associated with pet turtles in two states of the USA (CDC, 2005). Also in the USA, in October 2007, the North Carolina Division of Public Health notified to CDC human infections caused by *S. Paratyphi B* var. Java in several states. The results of the epidemiological and



laboratory investigations showed that many of these infections occurred in young children and were associated with exposure to small turtles (CDC, 2008a).

A survey with the purpose of estimating the role of pet reptiles in human salmonellosis was done in Europe (Bertrand *et al.*, 2008). In one study, in 2007-2008 seven human salmonellosis cases in Belgium were described which were related to contact with turtles and snakes. The same survey reported three reptile-associated cases of *Salmonella* infection in the past three years in France: two cases in 2005 and one, imported from China, in 2006. The patients were all young children, aged eight months, three years and four years, respectively. They were infected with a multi-resistant strain of *S. Typhimurium*. The first two cases had contact with, respectively, a snake and an iguana; in the third case an indirect link to a turtle (consumption of turtle soup) was found. At least 14 cases of human salmonellosis associated with reptile contact were identified in Ireland between 2005 and 2008, and 31 cases, also associated with reptiles, were reported in Germany during 2006-2008. Although infections in adults from contact with reptiles have been reported, in most cases infants less than one year old are affected.

In most countries, although cases of salmonellosis are reported within the national surveillance system, the source of infection is not routinely given, and the possible exposure to pet animals is usually only revealed in additional epidemiological investigations. Thus, the true number of cases due to direct or indirect contact with pet animals is likely underestimated.

Other sources of *Salmonella* for human beings

In some cases, salmonellosis may be associated with uncommon sources. It is frequently the result of factors including the ubiquitous distribution of *Salmonella* serovars, usage of contaminated ingredients, incorrect cleaning or disinfection of equipment in the food industry, and infected employees working in restaurant kitchens.

In the USA, an outbreak caused by a *Salmonella* strain with a specific PFGE pattern occurred during January 1- December 31, 2007. In this outbreak, a total of 401 cases from 41 states were identified and 32% of ill persons were hospitalized. A case-control study indicated that illness was associated with consumption of not-ready-to-eat pot pies. Later, the outbreak strain was isolated from 13 samples of unopened pot pies collected from the homes of patients (CDC, 2008b).

Moreover, in November 2006, public health officials of CDC and state health departments detected an increase in the reported incidence of isolates of *S. Tennessee*. In a multi-state case-control study conducted during February 5-13, 2007, illness was demonstrated to be strongly associated with the consumption of two brands of peanut butter produced at the same process plant. A total of 628 persons infected with this outbreak strain of *S. Tennessee* had been reported from 47 states during 2006-2007 (CDC, 2007a). Later, on November 25, 2008, an epidemiological investigation began due to an increase of *S. Typhimurium* isolates that shared the same PFGE patterns. Up to January 2009, 529 persons from 43 states from the USA and one person from Canada had been reported infected with this strain. A total of 116 patients were hospitalized, and the infection might have contributed to eight deaths. Sequential case-control studies indicated significant associations between illness and consumption of peanut butter, and specific brands of peanut butter biscuits. Epidemiological and laboratory findings indicated that peanut butter and peanut paste produced at one plant were the sources of the outbreak. These products were also ingredients in many foods produced and distributed by other companies. This outbreak highlights the complexities of "ingredient-driven" outbreaks and the importance of rapid outbreak detection and investigation (CDC, 2009).

On November 15, 2006, in the USA a case of salmonellosis in an employee of a poultry vaccine facility was notified. When a second case of salmonellosis in another employee at the same facility was reported on November 25, an outbreak investigation was initiated. Results of that investigation suggested that 21 employees of the facility became ill during a 1-month period from exposure to a strain of *S. Enteritidis* that was used in vaccine production. Infection was thought to have resulted from environmental contamination after the spill of a liquid containing a high concentration of *S. Enteritidis*. As a result, it was recommended that the facility improve its infection-control procedures to better protect workers. This outbreak highlights occupational risks that can be associated with the manufacture of veterinary biologicals derived from human pathogens (CDC, 2007d).

During October and November 2006, the Spanish National Reference Laboratory reported a series of isolations of *S. Kottbus* on the island of Gran Canaria. A matched case-control study was conducted and it showed that the ingestion of a brand of bottle water



was related to outbreak. Microbiological and environmental analysis detected *S. Kottbus* in bottles randomly selected from markets, and also in the local factory where the water was bottled. A carrier pigeon loft was found near one water reservoir of the local factory and it could be verified that pigeons frequented this reservoir. Moreover, *Salmonella* spp. was detected in the pigeons, suggesting that they could be contaminating the water reservoir (Palmera-Suárez *et al.*, 2007).

In 2002, an outbreak of *S. Enteritidis* phage-type 8 occurred at a convention center in Dallas, Texas, USA and continued for 6 weeks. From March to April, the hotel kitchen implicated catered for 41 multi-day conferences attended by 9,790 persons. A total of 617 illness reports from residents of 46 states were received. The epidemiological studies showed that sauces or items served with sauces were implicated in outbreak. Eleven food service employees, including one who prepared sauces, had stool cultures that yielded *S. Enteritidis* phage-type 8. Transmission ended with implementation of policies to screen food handlers and exclude those whose stool cultures yielded *S. Enteritidis* (Beatty *et al.*, 2009).

In addition, in Germany, from October 2001 to March 2002 about 439 *S. Oranienburg* reports were registered. Simultaneously, an increase in this strain was noticed in other European countries. A case-control study was conducted, indicating that the consumption of chocolate from one company was associated with illness. Later, two brands from the same company were positive for *S. Oranienburg*. Isolates from humans and from chocolates had indistinguishable PFGE patterns. No source or point of contamination was identified (Werber *et al.*, 2005).

Antibiotic resistance in *Salmonella* serovars: A serious problem in public health

Since the start of the widespread use of antimicrobial agents by humans in the late 1940s, antibiotic resistance as a phenomenon has been observed in almost all bacterial species and against all drugs available. Antimicrobial resistance can increase the morbidity, mortality, and costs associated with disease. Moreover, it has social and economic consequences and requires strong scientific and public health efforts to improve the situation. The increase in the number of resistant and multiresistant (resistant to two and more antimicrobials) strains of bacteria has been recognized by the World Health Organization

(WHO) and health authorities as one of the major problems in public health (Helmuth, 2001).

The use of antibiotics in food animal production systems has resulted in emergence of antibiotic resistant zoonotic bacteria that can be transmitted to humans through the food chain (Walsh & Fanning, 2008). Since the 1960s many studies have focused on this subject and the results are alarming. For instance, a rise in *S. Typhimurium* infection was observed in calves in United Kingdom during 1964-1966 following the adoption of the intensive farming methods (Anderson, 1968). A single phage type of *S. Typhimurium*, type 29, was incriminated as the major pathogen. Attempts to treat and control the disease with a range of antibiotics resulted in the acquisition of transferable multiple drug resistance by *S. Typhimurium* and the transmission of this strain, directly or indirectly, from bovines to man resulted in many human infections.

In Netherlands between 1972 and 1974, almost 50,000 *Salmonella* isolates from several sources (humans, animals, animal products, sewage, etc.) were tested for resistance to ampicillin, chloramphenicol, kanamycin, and tetracycline. In this study, the incidence of resistance to at least one of the above drugs ranged from 39.2% to 45.6%. Moreover, many multidrug-resistant strains of *S. Typhimurium* and *S. Dublin* were isolated from calves and cattle. In 1974, 64.4% of all strains of *S. Typhimurium* from these animals appeared to be resistant to ampicillin, tetracycline, chloramphenicol, and kanamycin, and 25.5% of *S. Dublin* were found to be resistant to chloramphenicol and tetracycline (Voogd *et al.*, 1977).

Holmberg *et al.* (1984) assessed the origin of antimicrobial resistance in infections caused by *Salmonella* serovars in 52 outbreaks investigated by the CDC between 1971 and 1983. According to this study, in 38 outbreaks with identified source, food animals were involved in 11 (69%) of 16 resistant and 6 (46%) of 13 sensitive outbreak strains, *S. Typhimurium* being the main serovar isolated and beef and milk the main products involved. In addition, Cohen & Tauxe (1986) analyzed the occurrence of antimicrobial resistance in *Salmonella* serovars in USA between 1960 and 1980, concluding that the use of antibiotics in animal husbandry played a major role in the emergence or persistence of these resistant *Salmonella* strains.

In 1989 an outbreak of *S. Typhimurium* DT 193 resistant to four antibiotics (ampicillin, streptomycin, sulphonamides, and tetracyclines) occurred in northern



England, affecting 206 persons. An analytical study showed a significant association between illness and consumption of cold roast pork supplied by the butcher's shop. Later, *S. Typhimurium* DT 193 with the same antibiotic resistance pattern as the outbreak strain was isolated from samples of meat bought from the shop and from samples of pig feces taken from the farm supplying the shop. It was concluded that inadequate processing of infected pork meat at the shop may have contributed to this outbreak (Maguire *et al.*, 1993).

Seyfarth *et al.* (1997) studied the frequency of antimicrobial resistance and epidemiological relatedness among 473 isolates of *S. Typhimurium* from human and veterinary sources in Denmark in 1993. In this study, the human strains were clinical isolates from patients with diarrhea (228 isolates) and animal strains were isolated from clinical or subclinical infections in cattle (48 isolates), pigs (99 isolates) and poultry (98 isolates). All strains were tested against 22 different antimicrobial agents used in both human and veterinary medicine with the disc diffusion method. Strains were also phage-typed and the plasmid content determined in all resistant strains. Of 228 human isolates tested, 19.3% of the strains were resistant to one or more antimicrobial agent compared with 10.4% of strains from cattle, 11.1% of strains from pigs and 9.2% of strains from poultry. Multiple resistance was found in 9.2% of the human strains, but in only two of the cattle isolates. Resistance in human strains was most common against tetracycline (13%), ampicillin (12%), sulphamide (12%), streptomycin (10%), and chloramphenicol (8%). The resistance pattern differed somewhat in animal isolates. Poultry strains were usually resistant only to ampicillin, while pig and cattle isolates were most often resistant to sulphamide, tetracycline and streptomycin. Typing of the strains showed that some animal strains and human strains were indistinguishable.

Moreover, Molla *et al.* (1999) assessed the antimicrobial resistance to 10 drugs in 39 *Salmonella* serovars isolated from raw minced beef and chicken (gizzard, liver, and heart) samples in Addis Ababa (Ethiopia). Thirty four isolates (87.2%) were resistant to one or more drugs. The antibiotics to which isolated strains were resistant included nitrofurantoin (48.7%), furazolidone (48.7%) and streptomycin (46.2%). Only four antimicrobials (gentamycin, kanamycin, rifampicin, and sulphamethoxazole-trimethoprim) were effective against all isolate *S. Enteritidis*. About 80% of the *S. Enteritidis* strains showed multiple resistance to up to four

antibiotics followed by *S. Typhimurium* (60.0%) and *S. Dublin* (33.3%).

In 1997, a *S. Typhimurium* DT104 strain resistant to five antimicrobials (ampicillin, chloramphenicol, streptomycin, sulfonamides, and tetracycline) affected 110 people in two outbreaks in USA. A case-control study and laboratory investigations indicated an unpasteurized Mexican-style cheese as the responsible for these outbreaks (Cody *et al.*, 1999).

White *et al.* (2001) identified and characterized strains of *Salmonella* isolated from ground chicken, beef, turkey, and pork purchased at three supermarkets of Washington, USA. The isolates were characterized by serotyping, antimicrobial-susceptibility testing, and phage-typing. Additionally, PCR and DNA sequencing were used to identify resistance integrons and extended spectrum beta-lactamase genes. Of 200 meat samples, 41 (20 %) were contaminated with 13 different serovars. Eighty four percent of the isolates were resistant to at least one antibiotic, and 53% were resistant to at least three antibiotics. Sixteen percent of the isolates were resistant to ceftriaxone. Five isolates of *S. Agona* had resistance to 9 antibiotics, and the two isolates of *S. Typhimurium* DT208 were resistant to 12 antibiotics. Eighteen isolates, representing four serotypes, had integrons with genes conferring resistance to aminoglycosides, sulfonamides, trimethoprim, and beta-lactams.

In another study, Chen *et al.* (2004) analyzed 133 isolates of *Salmonella* serovars recovered from retail meats purchased in the USA and China, assessing the antimicrobial resistance of these strains. In this study, it was demonstrated that seventy-three (82%) of these *Salmonella* serovars isolates were resistant to at least one antimicrobial agent. Resistance to the following antibiotics was common among the USA isolates: tetracycline (68% of the isolates were resistant), streptomycin (61%), sulfamethoxazole (42%), and ampicillin (29%). Eight *Salmonella* isolates (6%) were resistant to ceftriaxone. Fourteen isolates (11%) from China were resistant to nalidixic acid and displayed decreased susceptibility to ciprofloxacin. Moreover, Miko *et al.* (2005) determined antimicrobial resistance of *Salmonella* serovars isolated from German foodstuffs. A total of 319 epidemiologically independent multidrug-resistant isolates comprising 25 different serovars were tested for their antimicrobial susceptibility by broth microdilution. *S. Typhimurium* was the prevalent serovar, and it was isolated from beef and pork meat. The results of this study shown that the most prevalent resistances found in the multidrug-



resistant *Salmonella* serovars from foods were to streptomycin (94%), sulfamethoxazole (92%), tetracycline (81%), ampicillin (73%), spectinomycin (72%), chloramphenicol (48%), and trimethoprim (27%).

According to the USA Food and Drug Administration (FDA) (2006), in 2003, 22.5% of non-typhoid *Salmonella* isolates from humans were resistant to at least one antimicrobial agent and the most common multidrug-resistance phenotype reported was to ampicillin, chloramphenicol, streptomycin, sulfonamides, and tetracyclines, which was detected in 9.3% of isolates tested. While on the veterinary side, 44% of the *Salmonella* samples isolated from animal slaughter and veterinary diagnostic sources were resistant to at least one antimicrobial agent and the phenotype resistant to ampicillin, chloramphenicol, streptomycin, sulfonamides, and tetracyclines was also the most common multidrug resistance profile among veterinary isolates.

Between July 2004 and June 2005, a study was carried out in order to determine the prevalence and antimicrobial resistance of *Salmonella* serovars isolates from processed poultry in the USA. Four hundred eighty pre- and postchill whole broiler chicken carcasses were collected from a poultry processing plant. Water samples also were collected at the entrance and exit of the chiller. The proportions of pre- and postchill carcasses that were positive for *Salmonella* spp. were 88.4% and 84.1%, respectively. Ninety-two percent of water samples collected at the entrance of the chiller were positive for *Salmonella* spp., but all exit water samples were negative. Additionally, *Salmonella* spp. isolated were serotyped and tested for susceptibility to antimicrobials. Thirteen serotypes were identified; the most common were *S. Kentucky* (59.5%) and *S. Typhimurium* (17.8%). Three hundred thirty-nine (79.8%) of the isolates were resistant to at least one antimicrobial, and 53.4% were resistant to three or more antimicrobials. Resistance was most often observed to tetracycline (73.4% of isolates), ampicillin (52.9%), amoxicillin-clavulanic acid (52%), ceftiofur (51.7%), streptomycin (35.2%), and sulfisoxazole (21.8%). These results indicated that a large number of the isolates in poultry processing plants are frequently resistant to commonly used antimicrobials (Parveen *et al.*, 2007).

Valdezate *et al.* (2007) conducted a study in order to determine the distribution and antimicrobial susceptibility of *Salmonella* serovars isolated in 2002 from food in 16 Spanish regions. Fifty one serovars were

detected; the most common were *S. Enteritidis* and *S. Typhimurium*, followed by a group of 49 different serovars. The main sources of *Salmonella* serovars were eggs and their derivatives (21.6% of strains), poultry and related products (16.6%), and seafood (16.3%). Resistance rates for *S. Enteritidis*, *S. Typhimurium*, and the group of 49 serovars were, respectively, 8.3, 69.8, and 13.9% for ampicillin, 3.1, 52.8, and 59% for streptomycin, 40.6, 22.6, and 10.4% for nalidixic acid, 15.6, 71.7, and 31.1% for tetracycline, 7.3, 18.8, and 9.5% for trimethoprim-sulfamethoxazole, 0, 50.9, and 4.3% for chloramphenicol, and 6.2, 71.7, and 17.4% for multiple (at least four) antimicrobials. All the strains remained susceptible to other beta-lactams and fluoroquinolones.

From September 2003 to February 2004, Zewdu & Cornelius (2009) assessed antimicrobial resistance of 98 isolates of *Salmonella* serovars recovered from food in Addis Ababa, Ethiopia. The results of this study revealed that 32 *Salmonella* isolates were resistant to one or more of the 24 antimicrobials tested. The most common resistance was to streptomycin (75%), ampicillin (59.4%), tetracycline (46.9%), spectinomycin (40.6%), and sulfisoxazole (40.6%). The level of antimicrobial resistance was significantly higher for isolates from chicken carcass (62.1%) and pork isolates (22.7%).

In face of this public health problem, it is recommended a careful prescription of antimicrobial agents during veterinary practice, regardless of the purpose of this prescription (therapy, prophylaxis, and growth enhancement). This will increase the efficient use of antimicrobial agents after microbiological identification of the causative pathogen. Moreover, it is important to highlight that the use of any antimicrobial agent should not substitute good hygiene practices and, wherever possible, alternative management methods should be sought and used. In addition, veterinarians should respect and follow the sanitary legislation concerning the use of antimicrobial agents in their country.

Concluding remarks

Members of the genus *Salmonella* are major enteric pathogens of human beings and animals. The increase of the human population and food production increases the potential for dissemination of these ubiquitous microorganisms. The high incidence of human non-typhoid salmonellosis demonstrates the potential harm of the presence of *Salmonella* in food and contact animals to people. In addition to disease caused by



Salmonella serovars, antibiotic resistance can be transferred by other microorganisms that are agents of human disease similar to those involved in hospital-acquired infections. Fortunately, we can learn from undesirable episodes of life. The outbreaks of food-borne *S. Enteritidis* disease related to chickens from 1980's triggered a great deal of discussion related to its control. As a result, in the understanding of the epidemiology of the infection in commercial birds improved. Thus, monitoring program could also be improved resulting in more efficient control programs in the food chain. In addition, nowadays there is an increasing worldwide preoccupation with the quality of the food; many countries have banned the use of antimicrobial agents in the feed and they are encouraging farmers to use non-harmful substances. All measures taken over animal husbandry may minimize the presence of *Salmonella* in food. However, the contribution of the consumers is also necessary by adopting the basic concepts of hygiene during food preparation and storing.

References

- Anderson ES. Drug resistance in *Salmonella* typhimurium and its implications. *British Medical Journal* 1968; 3(5614):333-339.
- Beatty ME, Shevick G, Shupe-Ricksecker K, Bannister E, Tulu A, Lancaster K. *et al.* Large *Salmonella* Enteritidis outbreak with prolonged transmission attributed to an infected food handler, Texas, 2002. *Epidemiology and Infection* 2009; 137(3):417-427.
- Bertrand S, Rimhanen-Finne R, Weill FX, Rabsch W, Thornton L, Perevoscikovs J, van Pelt W, Heck M. *Salmonella* infections associated with reptiles: the current situation in Europe. *Euro Surveillance* 2008; 13(24).
- Boycott JA, Taylor J, Douglas SH. *Salmonella* in tortoiseS. *The Journal of Pathology and Bacteriology* 1953; 65(2):401-411.
- Braden CR. *Salmonella enterica* Serotype Enteritidis and Eggs: A National Epidemic in the United StateS. *Clinical Infectious Diseases: an official publication of the Infectious Diseases Society of America* 2006; 43(4):512-517
- Bruins MJ, de Boer AM, Ruijs GJ. Gastroenteritis caused by *Salmonella* from pet snakes. *Nederlands Tijdschrift Voor Geneeskunde* 2006; 150(41):2266-2269.
- Cahill SM, Wachsmuth IK, Costarrica Mde L, Ben Embarek PK. Powdered infant formula as a source of *Salmonella* infection in infants. *Clinical Infectious Diseases: an official publication of the Infectious Diseases Society of America* 2008; 46(2):268-273.
- CDC. Centers for Diseases Control and Prevention. Multiple outbreaks of salmonellosis associated with precooked roast beef-Pennsylvania, New York, Vermont. *Morbidity and Mortality Weekly Report* 1981; 30(46):569-570.
- CDC. Centers for Diseases Control and Prevention. Multistate outbreak of human *Salmonella* infections associated with exposure to turtles-United States, 2007-2008. *Morbidity and Mortality Weekly Report* 2008a; 57(3):69-72.
- CDC. Centers for Diseases Control and Prevention. Multistate outbreak of *Salmonella* infections associated with frozen pot pies-United States, 2007. *Morbidity and Mortality Weekly Report* 2008b; 57(47):1277-1280.
- CDC. Centers for Diseases Control and Prevention. Multistate outbreak of *Salmonella* infections associated with peanut butter and peanut butter-containing products-United States, 2008-2009. *Morbidity and Mortality Weekly Report* 2009; 58(4):85-90.
- CDC. Centers for Diseases Control and Prevention. Multistate outbreak of *Salmonella* serotype Tennessee infections associated with peanut butter-United States, 2006-2007. *Morbidity and Mortality Weekly Report* 2007a; 56(21):521-524.
- CDC. Centers for Diseases Control and Prevention. Multistate outbreak of *Salmonella* Typhimurium infections associated with eating ground beef-United States, 2004. *Morbidity and Mortality Weekly Report* 2006; 55(7):180-182.
- CDC. Centers for Diseases Control and Prevention. Multistate outbreaks of *Salmonella* infections associated with raw tomatoes eaten in restaurants-United States, 2005-2006. *Morbidity and Mortality Weekly Report* 2007b; 56(35):909-911.
- CDC. Centers for Diseases Control and Prevention. Outbreak of multidrug-resistant *Salmonella* enterica serotype Newport infections associated with consumption of unpasteurized Mexican-style aged cheese-Illinois, March 2006-April 2007. *Morbidity and Mortality Weekly Report* 2008c; 57(16):432-435.
- CDC. Centers for Diseases Control and Prevention. Outbreak of *Salmonella* serotype Typhimurium infection associated with eating raw ground beef-Wisconsin, 1994. *Morbidity and Mortality Weekly Report* 1995a; 44(49):905-909.
- CDC. Centers for Diseases Control and Prevention. Reptile-associated salmonellosis-selected states, 1994-1995. *Morbidity and Mortality Weekly Report* 1995b; 44(17):347-350.
- CDC. Centers for Diseases Control and Prevention. *Salmonella*: annual summary 2005. Atlanta, Georgia: US Department of Health and Human Services; 2007c [cited 2009 jul 14]. Available from: <http://www.cdc.gov/ncidod/dbmd/phlisdata/salmtab/2005/SalmonellaAnnualSummary2005.pdf>.
- CDC. Centers for Diseases Control and Prevention. *Salmonella* serotype Enteritidis infections among workers producing poultry vaccine-Maine, November-December 2006. *Morbidity and Mortality Weekly Report* 2007d; 56(34):877-879.
- CDC. Centers for Diseases Control and Prevention. *Salmonella* Typhimurium infection associated with raw milk and cheese consumption-Pennsylvania, 2007. *Morbidity and Mortality Weekly Report* 2007e; 56(44):1161-1164.



- CDC. Centers for Diseases Control and Prevention. Salmonellosis associated with pet turtles--Wisconsin and Wyoming, 2004. *Morbidity and Mortality Weekly Report* 2005; 54(9):223-226.
- CDC. Centers for Diseases Control and Prevention. Turtle-associated salmonellosis in humans--United States, 2006-2007. *Morbidity and Mortality Weekly Report* 2007f; 56(26):649-652.
- Clark GM, Kaufmann AF, Gangarosa EJ, Thompson MA. Epidemiology of an international outbreak of *Salmonella* Agona. *Lancet* 1973; 2(7827):490-493.
- Chen S, Zhao S, White DG, Schroeder CM, Lu R, Yang H, McDermott PF, Ayers S, Meng J. Characterization of multiple-antimicrobial-resistant *Salmonella* serovars isolated from retail meat. *Applied and Environmental Microbiology* 2004; 70(1):1-7.
- Codex, 1979. Recommended International Code Of Hygienic Practice For Foods For Infants And Children CAC/RCP 21-1979 [cited 2009 jul 14]. Available from: <http://www.euro.who.int/document/fos/FoodsforInfantsandChildren.pdf>.
- Cody SH, Abbott SL, Marfin AA, Schulz B, Wagner P, Robbins K, Mohle-Boetani JC, Vugia DJ. Two outbreaks of multidrug-resistant *Salmonella* serotype typhimurium DT104 infections linked to raw-milk cheese in Northern California. *JAMA: the journal of the American Medical Association* 1999; 281(19):1805-1810.
- Cohen ML, Tauxe RV. Drug-resistant *Salmonella* in the United States: an epidemiologic perspective. *Science* 1986; 234(4779):964-969.
- Corrente M, Totaro M, Martella V, Campolo M, Lorusso A, Ricci M, Buonavoglia C. Reptile-associated salmonellosis in man, Italy. *Emerging Infectious Diseases* 2006; 12(2):358-359.
- Davies A, O'Neill P, Towers L, Cooke M. An outbreak of *Salmonella* typhimurium DT104 food poisoning associated with eating beef. *Communicable Disease Report. CDR Review* 1996; 6(11):159-162.
- Delpech V, McAnulty J, Morgan K. A salmonellosis outbreak linked to internally contaminated pork meat. *Australian and New Zealand Journal of Public Health* 1998; 22(2):243-246.
- EFSA. European Food Safety Authority. The Community Summary Report on Food-borne Outbreaks in the European Union in 2007. *The EFSA Journal* 2009; 271.
- Ethelberg S, Sørensen G, Kristensen B, Christensen K, Krusell L, Hempel-Jørgensen A, Perge A, Nielsen EM. Outbreak with multi-resistant *Salmonella* Typhimurium DT104 linked to carpaccio, Denmark, 2005. *Epidemiology and Infection* 2007; 135(6):900-907.
- Fernandes SA, Tavecchio AT, Ghiliardi AC, Dias AM, Almeida IA, Melo LC. *Salmonella* serovars isolated from humans in São Paulo State, Brazil, 1996-2003. *Revista do Instituto de Medicina Tropical de São Paulo* 2006; 48:179-184.
- FDA. US Food and Drug Administration. National Antimicrobial Resistance Monitoring System- Enteric Bacteria (NARMS): 2003 Executive report. Rockville, MD: US Department of Health and Human Services, FDA, 2006 [cited 2009 jul 14]. Available from:<http://www.fda.gov/downloads/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/UCM061471.pdf>.
- Greene SK, Daly ER, Talbot EA, Demma LJ, Holzbauer S, Patel NJ, Hill TA, Walderhaug MO, Hoekstra RM, Lynch MF, Painter JA. Recurrent multistate outbreak of *Salmonella* Newport associated with tomatoes from contaminated fields, 2005. *Epidemiology and Infection* 2008; 136(2):157-165.
- Haeghebaert S, Duché L, Gilles C, Masini B, Dubreuil M, Minet JC, Bouvet P, Grimont F, Delarocque Astagneau E, Vaillant V. Minced beef and human salmonellosis: review of the investigation of three outbreaks in France. *Euro Surveill* 2001; 6(2):21-26.
- Helmuth, R. Antibiotic Resistance in *Salmonella*. In: Wray C, Wray A, editor. *Salmonella* in domestic animal. Wallingford: CAB International; 2001. p. 89-106.
- Hess IM, Neville LM, McCarthy R, Shadbolt CT, McAnulty JM. A *Salmonella* Typhimurium 197 outbreak linked to the consumption of lambs' liver in Sydney, NSW. *Epidemiology and Infection* 2008; 136(4):461-467.
- Holmberg SD, Osterholm MT, Senger KA, Cohen ML. Drug-resistant *Salmonella* from animals fed antimicrobials. *The New England Journal of Medicine* 1984; 311(10):617-622.
- Jain S, Bidol SA, Austin JL, Berl E, Elson F, Lemaile-Williams M, Deasy M 3rd, Moll ME, Rea V, Vojdani JD, Yu PA, Hoekstra RM, Braden CR, Lynch MF. Multistate outbreak of *Salmonella* Typhimurium and Saintpaul infections associated with unpasteurized orange juice--United States, 2005. *Clinical Infectious Diseases: an official publication of the Infectious Diseases Society of America* 2009; 48(8):1065-1071.
- Jansen A, Frank C, Stark K. Pork and pork products as a source for human salmonellosis in Germany. *Berliner und Münchener tierärztliche Wochenschrift* 2007; 120(7-8):340-346.
- Kivi M, Hofhuis A, Notermans DW, Wannet WJ, Heck ME, Van De Giessen AW, Van Duynhoven YT, Stenvers OF, Bosman A, Van Pelt W. A beef-associated outbreak of *Salmonella* Typhimurium DT104 in The Netherlands with implications for national and international policy. *Epidemiology and Infection* 2007; 135(6):890-899.
- Kretsinger K, Noviello S, Moll M. Tomatoes sicken hundreds: multistate outbreak of *Salmonella* Newport. *Proceedings of the 52nd Annual Epidemic Intelligence Service Conference*; 2003; Atlanta, Georgia. USA.
- Little CL, Gillespie IA. Prepared salads and public health. *Journal of Applied Microbiology* 2008; 105(6):1729-1743.
- Luzzi I, Galetta P, Massari M, Rizzo C, Dionisi AM, Filetici E, Cawthorne A, Tozzi A, Argentieri M, Bilei S, Busani L, Gnesivo C, Pendenza A, Piccoli A, Napoli P, Loffredo L, Trinito MO, Santarelli E, Ciofi degli Atti ML. An Easter outbreak of *Salmonella* Typhimurium DT 104A associated with traditional pork salami in Italy. *Euro Surveill* 2007; 12(4):11-12.
- Maguire HC, Codd AA, Mackay VE, Rowe B, Mitchell E. A large



outbreak of human salmonellosis traced to a local pig farm. *Epidemiology and Infection* 1993; 110(2):239-246.

Mermin J, Hutwagner L, Vugia D, Shallow S, Daily P, Bender J. Reptiles, amphibians and human *Salmonella* infection: a population-based, case-control study. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America* 2004; 38 Suppl 3:S253-261.

Miko A, Pries K, Schroeter A, Helmuth R. Molecular mechanisms of resistance in multidrug-resistant serovars of *Salmonella enterica* isolated from foods in Germany. *The Journal of Antimicrobial Chemotherapy* 2005; 56(6):1025-1033.

Molla B, Kleer J, Sinell HJ Antibiotic resistance pattern of foodborne *Salmonella* isolates in Addis Ababa (Ethiopia). *Berliner und Münchener tierärztliche Wochenschrift* 1999; 112(2):41-43.

Nygård K, Lassen J, Vold L, Andersson Y, Fisher I, Löfdahl S, Threlfall J, Luzzi I, Peters T, Hampton M, Torpdahl M, Kapperud G, Aavitsland P. Outbreak of *Salmonella* Thompson infections linked to imported rucola lettuce. *Foodborne Pathogens and Disease* 2008; 5(2):165-173.

Palmera-Suárez R, García P, García A, Barrasa A, Herrera D. *Salmonella* Kottbus outbreak in infants in Gran Canaria (Spain), caused by bottled water, August-November 2006. *Euro Surveillance* 2007; 12(7):E070712.2.

Parveen S, Taabodi M, Schwarz JG, Oscar TP, Harter-Dennis J, White DG. Prevalence and antimicrobial resistance of *Salmonella* recovered from processed poultry. *Journal of Food Protection* 2007; 70(11):2466-2472.

Pastore R, Schmid H, Altpeter E, Baumgartner A, Hächler H, Imhof R, Sudre P, Boubaker K. Outbreak of *Salmonella* serovar Stanley infections in Switzerland linked to locally produced soft cheese, September 2006 - February 2007. *Euro Surveillance* 2008; 13(37):18979.

Popoff MY, Minor L. Antigenic formulae of the *Salmonella* serovars. 8th ed. Paris: WHO Collaborating Center for Reference and Research on *Salmonella*; 2001.

Porwollik S, Boyd EF, Choy C, Cheng P, Florea L, Proctor E. Characterization of *Salmonella enterica* subspecies I genovars by use of microarrayS. *Journal of Bacteriology* 2004; 186(17):5883-5898.

Roels TH, Frazak PA, Kazmierczak JJ, Mackenzie WR, Proctor ME, Kurzynski TA, Davis JP. Incomplete sanitation of a meat grinder and ingestion of raw ground beef: contributing factors to a large outbreak of *Salmonella* Typhimurium infection. *Epidemiology and Infection* 1997; 119(2):127-134.

Seyfarth AM, Wegener HC, Frimodt-Møller N. Antimicrobial resistance in *Salmonella enterica* subsp. *enterica* serovar Typhimurium from humans and production animals. *The Journal of Antimicrobial Chemotherapy* 1997; 40(1):67-75.

Shapiro R, Ackers ML, Lance S, Rabbani M, Schaefer L, Daugherty J, Thelen C, Swerdlow D. *Salmonella* Thompson associated with

improper handling of roast beef at a restaurant in Sioux Falls, South Dakota. *Journal of Food Protection* 1999; 62(2):118-122.

Spitalny KC, Okowitz EN, Vogt RL. Salmonellosis outbreak at a Vermont hospital. *The Southern Medical Journal* 1984; 77(2):168-172.

Toyofuku H, Kubota K, Morikawa K. Outbreaks of *Salmonella* in infants associated with powdered infant formula. *Bulletin of National Institute of Health Sciences* 2006; 124:74-79.

Unicomb LE, Simmons G, Merritt T, Gregory J, Nicol C, Jelfs P, Kirk M, Tan A, Thomson R, Adamopoulos J, Little CL, Currie A, Dalton CB. Sesame seed products contaminated with *Salmonella*: three outbreaks associated with tahini. *Epidemiology and Infection* 2005; 133(6):1065-1072.

Valdezate S, Arroyo M, González-Sanz R, Ramiro R, Herrera-León S, Usera MA, De la Fuente M, Echeita A. Antimicrobial resistance and phage and molecular typing of *Salmonella* strains isolated from food for human consumption in Spain. *Journal of Food Protection* 2007; 70(12):2741-2748.

Vojdani JD, Beuchat LR, Tauxe RV. Juice-associated outbreaks of human illness in the United States, 1995 through 2005. *Journal of Food Protection* 2008; 71(2):356-364.

Voogd CE, van Leeuwen WJ, Guinée PA, Manten A, Valkenburg JJ. Incidence of resistance to ampicillin, chloramphenicol, kanamycin and tetracycline among *Salmonella* species isolated in the Netherlands in 1972, 1973 and 1974. *Antonie van Leeuwenhoek* 1977; 43(3-4):269-281.

Walsh C, Fanning S. Antimicrobial resistance in foodborne pathogens--a cause for concern?. *Current Drug Targets* 2008; 9(9):808-815.

Werber D, Dreesman J, Feil F, van Treec U, Fell G, Ethelberg S, *et al.* International outbreak of *Salmonella* Oranienburg due to German chocolate. *BMC Infectious Diseases* 2005; 5(1):7.

White DG, Zhao S, Sudler R, Ayers S, Friedman S, Chen S, McDermott PF, McDermott S, Wagner DD, Meng J. The isolation of antibiotic-resistant *Salmonella* from retail ground meatS. *The New England Journal of Medicine* 2001; 345(16):1147-1154.

Zewdu E, Cornelius P. Antimicrobial resistance pattern of *Salmonella* serotypes isolated from food items and personnel in Addis Ababa, Ethiopia. *Tropical Animal Health and Production* 2009; 41(2):241-249.