



Quality Assessment of Marketed Eggs in Bassekabyllie (Algeria)

Technical Note

■ Author(s)

Moula N[†]
Ait-Kaki A^{†*}
Leroy P[‡]
Antoine-Moussiaux N[†]

[†] Department of Animal Production, Faculty of Veterinary Medicine, University of Liege, Liege 4000, Belgium

[‡] Departement of Biochemistry and microbiology, Faculty of natural sciences, University of Mentouri, Constantine, Algeria.

* Nassim Moula and Asma Ait Kaki contributed equally to this publication.

■ Mail Address

Corresponding author e-mail address
Prof. P. Leroy & Dr. N. Antoine-Moussiaux
E-mail: pascal.leroy@ulg.ac.be
& nantoine@ulg.ac.be

■ Keywords

Bejaia, consumer channels, egg quality, egg prices, marketed table eggs, egg grades.

ABSTRACT

Quality variations in retail eggs are widely reported. This study aims at assessing the quality of eggs according to the marketing channel in the department of Bejaia (Algeria). In spring and summer 2012, 3330 eggs were bought in 30 stores divided into 3 categories: 10 supermarkets (1146 eggs), 10 public markets (1048 eggs), and 10 shops (1136 eggs). Egg weights differed significantly between marketing channels with 58.9 ± 0.14 , 61.2 ± 0.13 and 62.8 ± 0.13 g for public markets, shops and supermarkets, respectively ($p < 0.001$). Although shell thickness was similar for all marketing channels, the proportion of damaged eggs was higher in public markets (9.0%), intermediate in food shops (7.3%) and lower in supermarkets (5.7%; $p < 0.05$). The yolk/albumen ratio was significantly higher for eggs from supermarkets (48.0%) compared with the other channels (around 47.4%; $p < 0.05$). The freshness of the eggs, measured by the Haugh method, was lower in public markets (74.3 units), intermediate in shops (77.6 units) and higher in supermarkets (79.9 units; $p < 0.05$). The price of eggs, expressed in Algerian Dinar (AD) per kg, was significantly lower in public markets (124 AD/kg) compared with the two other channels (around 131 AD/kg; $p < 0.05$). It is possible to conclude that egg quality in Bass Kabylie differs significantly among marketing channels, with higher quality observed in supermarkets. The lower quality of eggs in public markets is associated with lower price. Eggs from shops present an intermediate quality. A one-year study would allow studying both the potential seasonal effect and compare intrinsic variability across marketing channels.

INTRODUCTION

Hen's eggs have been traditionally considered as an important source of nutrients for humans (Nau *et al.*, 2003). It is a source of proteins, lipids, minerals and vitamins easily renewable. The egg belongs to the limited category of foods containing the nine amino acids that human cannot synthesize. It was thus chosen by World Health Organization (W.H.O.) as the reference protein source for children (reference 100, which is slightly higher than women's milk) (Nys & Sauveur, 2004). The wide variety of poultry production systems and the low cost price of eggs make them widely accessible to rural and urban population (Moula, 2012). In addition, they are accepted worldwide and are not subject to major cultural or religious prohibitions (Bessadok *et al.*, 2003). Changes in lifestyle and consumption habits, accompanied by the development of the fast food, increased its demand. Indeed, egg proteins are appreciated ingredients in many foods (Mine, 2002).

The production and consumption of eggs in Algeria have undergone a rapid evolution since the 1980s, and continues to increase. Eggs are bought in many different retail spots in Algeria. However, the three



major channels are supermarkets (urban and peri-urban areas), public markets (in urban centers) and small foodshops (urban, peri-urban and rural areas). The aim of this study was to investigate the quality of eggs offered to the consumers in the department of Bejaia. The egg quality traits taken into consideration were egg weight, freshness (Haugh units), yolk color, shell thickness and yolk/albumen ratio. These characteristics are related to egg production (breed, feeding) and marketing (delay, transport care) features.

MATERIALS AND METHODS

A total of 3330 eggs were bought from three different marketing channels: 10 supermarkets, 10 public markets and 10 small food shops. The number of analyzed eggs in each source is listed in Table 1.

Table 1 – Eggs number by marketing channels and seasons

Season	Number of eggs			
	Total	Food shops	Public markets	Supermarkets
Spring	1624	584	494	546
Summer	1706	552	554	600
Total	3330	1136	1048	1146

A random sample was obtained twice from each source at one-week interval during spring and summer. Eggs are mainly sold by batches of 30 units, without any information on the source, laying date, or egg size.

After numbering the eggs, the here-above listed measurements were carried out. Haugh units are an estimate of eggs freshness. These are measured using an electronic scale and a tripod micrometer. Each egg was weighed and broken on a glass surface. The thickness of the albumen was measured with the tripod micrometer, at its maximum height. Haugh units were calculated using the formula described by Haugh (Haugh, 1937): $HU = 100 \log (H - 1.7 w^{0.37} + 7.6)$; where HU = Haugh units, H = Albumen length (mm), W = egg weight (g). Egg yolk color was determined using the DSM Yolk Color scale. After removing shell membranes, the thickness of eggshell was measured on three equator points with an electronic micrometer. The average of these three values was considered for data analysis. Tyler and Geake (1964) reported that the eggshell is thinner but almost uniform on equatorial zone. Eggs were graded according to USDA standards in respect to the studied quality components (USDA, 1975). The eggs price per kg in Algerian Dinar (1DA \approx 0.01€) was also calculated.

The statistical analyses were carried out using SAS software (SAS, 2001). The generalized linear model (GLM) was used to perform an analysis of

variance (ANOVA) of each parameter to determine the differences between the three studied retail sources and their statistical significance. For each parameter, the least squares means (LSM) and the standard errors were calculated. The chi-square test was used to test independence between the qualitative variables and retail sources.

RESULTS AND DISCUSSION

Egg quality is a general term, which refers to several standards that define both internal and external quality. Internal quality refers to egg white (albumen) cleanliness and viscosity, size of the air cell, yolk shape and yolk ratio. Exterior egg quality is defined as texture, color, soundness, cleanliness and shape of the shell. The shell of each egg should be smooth, clean and free from cracks and eggs should be uniform in color, size and shape (Coutts *et al.*, 2006). Egg composition is not a uniform trait and depends on many factors, such as hen breed and egg storage time (Moula *et al.*, 2009; Moula *et al.*, 2010). Variability in the quality of eggs available to consumers has been reported by many investigators (Vanghan and Adams, 1959; Abo Omar & Aref, 2000; Bell *et al.*, 2001). However, little is known about the quality of eggs commercialized in Algeria.

The statistical analysis showed a significant effect of the retail type on egg weight, Haugh unit, shell thickness and price per kg ($p < 0.05$). The season and the interaction between marketing channel and season had no significant effect on egg quality traits ($p > 0.05$) except for egg weight ($p < 0.05$). The season significantly impacted Haugh units ($p < 0.001$), shell thickness ($p < 0.0001$) and price per kg ($p < 0.05$). Marketing channels significantly influenced egg weight ($p < 0.0001$), Haugh unit ($p < 0.0001$), shell thickness ($p < 0.01$) and Y/A ratio ($p < 0.05$). The eggs bought in supermarkets were significantly heavier than those bought in small foodshops and public markets ($p < 0.05$). However, the price per kg was not significantly different between shops and supermarkets (Table 2). The difference of egg weight according to marketing channel was observed in this study (Table 2). Consistent findings were previously obtained by Abo Omar & Aref (2000) and by North (1984), but these authors reported that eggs from shops and public markets were significantly ($p < 0.01$) heavier when compared to supermarkets.

Algeria is a Mediterranean country with a hot summer. This explains the influence of season on egg



Table 2 – Least Squares Means and standard errors of egg weight, Haugh unit, yolk color, yolk/albumen ratio, shell thickness and price per kg.

	Season	Food Shops	Public markets	Supermarkets	Season	Marketing Channels	C*S	R ²
Egg weight (g)	Spring	60.75±0.19 ^{a1}	59.40±0.20 ^{b1}	62.74±0.19 ^{c1}	ns	***	***	.11
	Summer	61.71±0.19 ^{a2}	58.53±0.19 ^{b2}	62.91±0.19 ^{c1}				
	Total	61.22±0.13 ^a	58.94±0.14 ^b	62.83±0.13 ^c				
Y/A ratio	Spring	47.58±0.15 ^{ab1}	47.36±0.16 ^{b1}	47.85±0.16 ^{a1}	ns	*	ns	.01
	Summer	47.01±0.16 ^{a2}	47.69±0.16 ^{b1}	48.11±0.15 ^{b1}				
	Total	47.31±0.11 ^a	47.53±0.11 ^a	47.98±0.11 ^b				
Haugh unit	Spring	81.97±0.60 ^{a1}	79.13±0.65 ^{b1}	81.53±0.58 ^{a1}	***	***	***	.09
	Summer	72.90±0.61 ^{a2}	69.91±0.61 ^{b2}	78.46±0.59 ^{a2}				
	Total	77.56±0.44 ^a	74.26±0.46 ^b	79.92±0.44 ^c				
Yolk color	Spring	11.04±0.07 ^{a1}	11.15±0.07 ^{ab1}	11.23±0.07 ^{b1}	ns	ns	*	.01
	Summer	11.34±0.07 ^{a2}	11.01±0.07 ^{b1}	11.24±0.06 ^{a1}				
	Total	11.19±0.05 ^{ab}	11.07±0.05 ^b	11.23±0.05 ^a				
Shell thickness (.01mm)	Spring	37.36±0.16 ^{a1}	37.45±0.17 ^{a1}	38.06±0.17 ^{b1}	***	**	ns	.02
	Summer	36.28±0.17 ^{a2}	36.80±0.17 ^{b2}	36.91±0.16 ^{b2}				
	Total	36.84±0.12 ^a	37.11±0.12 ^a	37.46±0.12 ^a				
Price/kg (DA)	Spring	131.75±3.62	123.47±3.62	130.77±3.62	*	ns	ns	.32
	Summer	126.71±3.62 ^b	124.03±3.62 ^b	135.26±3.62 ^a				
	Total	129.23±2.53 ^a	123.75±2.53 ^b	133.02±2.53 ^a				

Means followed by the same letter (a,b,c) in the same row are not statistically different (p-value>0.05). Means followed by the same number (1, 2) in the same column are not statistically different (p-value>0.05). ***: p<0.0001; **: p<0.001; *: p<0.05; NS: p>0.05. Season: S; Marketing Channels : C.

quality (Table 2), especially on their freshness (HU units). In fact, egg freshness was significantly higher in spring. The same result was found by Islam *et al.* (2001). Thus, under high temperature conditions, egg production and egg quality decrease (Leeson, 1986) and one of the reasons of these detrimental effects is insufficient feed intake (Leeson and Summers, 1997). Eggs sold in supermarkets in this study, presented the best quality, considering their freshness. In addition, their freshness was higher than those sold in supermarkets of California (HU: 61.1), Illinois (HU: 62.8), Pennsylvania

(64.0), Texas (HU: 59.6), North Carolina (HU: 67.7) and New England (HU: 68.1), as described by Bell *et al.* (2001). No significant effect of marketing channel on the egg price per kg was observed in the study of Abo Omar *et al.* (2000). However, here, the price of eggs in public markets was significantly lower compared with other marketing sources.

Eggweight categories are shown in Table 3. These categories were clustered according to the European classification. The chi-square test showed a significant link between weight classes and marketing channels

Table 3 – Distribution (%) of weight classes

Egg weight classes ¹	Season	Consumer Channels			Mean	Statistical Significance
		Foodshops	Public markets	Supermarkets		
Extra-large	Spring	0.86	-	14.47	5.17	***
	Summer	1.63	2.89	4.83	3.17	***
	Total	1.23	1.53	9.42	4.14	***
Large	Spring	20.03	12.15	25.46	19.46	***
	Summer	28.80	14.62	35.50	26.55	***
	Total	24.30	13.45	30.72	23.09	***
Medium	Spring	76.71	78.95	60.07	71.80	***
	Summer	69.57	67.69	59.67	65.47	***
	Total	73.24	73.00	59.86	68.56	***
Small	Spring	2.40	8.91	-	3.57	***
	Summer	-	14.80	-	4.81	***
	Total	1.23	12.02	-	4.20	***

¹European grades



Table 4 – Distribution (%) of damaged eggs according to the marketing channels.

Damaged eggs	Season	Consumer Channels			Mean	Statistical Significance
		Food shops	Public markets	Supermarkets		
%	Spring	5.14	12.75	5.13	7.45	***
	Summer	9.60	5.60	6.17	7.09	*
	Total	7.31	8.97	5.67	7.27	*

($p < 0.05$). Most of the eggs bought were medium-sized. In the supermarkets, nearly 10% of eggs were extra-large. This category represented less than 2% of eggs in public markets and foodshops (Table 3).

The proportion of damaged eggs was higher in public markets (9.0%), intermediate in food shops (7.3%) and lower in supermarkets (5.7%; $p < 0.05$) (Table 4).

Haugh units were significantly lower in eggs sold in public markets, indicating that eggs were not very fresh ($p < 0.05$; Table 3). This result may be explained by the exposure of eggs to air. Storage conditions, including temperature, humidity, presence of CO_2 , and duration of storage, are also of prime importance for egg quality at retail (Samli *et al.*, 2005). Storage time and temperature appear to be the most crucial factors affecting albumen quality or Haugh unit (HU). A high percentage of the analyzed eggs were AA grade, i.e., according to USDA norms (USDA, 1975) concerning interior quality (Table 5). Yolk color of eggs obtained in different marketing channels was not different, which can be explained by the uniformity of the feed used in poultry farming in Algeria.

As a conclusion, marketing channel significantly influences eggs quality in Bass Kabylie, with higher quality observed in eggs sold in supermarkets. The lower quality of eggs bought in public markets is associated with their lower price. Nevertheless, this lower price should not automatically be ascribed

to the lower quality. Rather, the shorter marketing chains may provide a better explanation. For most of the parameters, eggs from small food shops present intermediate values. Accordingly, this study indicates to the need of quality improvements in short marketing chains. The promotion of such local markets would provide outlets for eggs of local breeds and promote the preservation of biodiversity and inclusive rural development.

REFERENCES

- Abo Omar JM, Aref RO. Some quality characteristics of eggs marketed in North of West Bank (Palestine). *Revue de Médecine Vétérinaire* 2000;151:47-50.
- Bell DD, Patterson PH, Koelkebeck KW, Anderson K E, Darre M J, Carey J B, Kuney DR, Zeidler G. Egg Marketing in National Supermarkets: Egg Quality—Part 1. *Poultry Science* 2001; 80:383-389.
- Bessadok A, Khochilef I, El Gazzah M. Etat des ressources génétiques de la population locale du poulet en Tunisie. *Tropicicultura* 2003;21:167-172.
- Coutts JA, Wilson GC, Fernandez S, Rasales E, Weber G, Hernandez JM. Optimum Egg Quality – A Practical Approach. Sheffield: 5M Publishing; 2006. p.63
- Haugh RR. The Haugh unit for measuring egg quality. *U.S. Egg poultry Magazine* 1937;43: 522-573.
- Islam MA, Bulbul SM, Seeland G, Egg Quality of Different Chicken Genotypes in Summer and Winter. *Pakistan Journal of Biological Sciences* 2001; 4:1411-1414.
- Leeson S, Summers JD. Feeding programs for laying hens. *Guelph: Commercial Poultry Nutrition*; 1997.p.143–206.

Table 5 – Haugh unit values and grade distribution (%) of eggs according to the marketing channels.

Haugh unit value	USDA grade	Season	Distribution of weight classes (%)			Mean	Statistical Significance
			Marketing Channels				
			Foodshops	Public markets	Supermarkets		
>72	AA	Spring	69.69	62.75	76.92	70.01	***
		Summer	18.23	47.29	65.50	56.62	***
		Total	63.20	54.58	70.94		
60-72	A	Spring	22.09	16.40	16.30	18.41	***
		Summer	25.00	23.10	24.00	24.03	***
		Total	23.50	19.94	20.33		
31-60	B	Spring	8.05	17.61	6.78	10.53	***
		Summer	18.30	26.71	10.50	18.29	***
		Total	13.03	22.42	8.73		
<31	C	Spring	0.17	3.24	-	1.05	***
		Summer	0.36	2.89	-	1.06	***
		Total	0.26	8.73	-		



- Leeson S. Nutritional consideration of poultry during heat stress. *World's Poultry Science Journal* 1986;42:69–79.
- Mine Y. Recent advances in egg protein functionality in the food system. *World's Poultry Science Journal* 2002; 58: 31-39.
- Moula N, Antoine-Moussiaux N, Farnir F, Leroy P. Comparison of Egg Composition and Conservation Ability in Two Belgian Local Breeds and One Commercial Strain. *International Journal of Poultry Sciences* 2009;8:768-774.
- Moula N, Antoine-Moussiaux N, Decuypere E, Farnir F, Mertens K, De Baerdemaeker J, Leroy P. Comparative study of egg quality traits in two Belgian local breeds and two commercial lines of chickens. *Archiv Fuer Gefluegelkunde* 2010;74:164–171.
- Moula N. Biodiversité avicole dans les pays industrialisés et en développement: caractérisation et étude des performances de production de races gallines locales. Liège: Presses de la Faculté de Médecinevétérinaire de l'Université de Liège; 2012. 261p.
- Nau F, Anton M, Nys Y. L'œuf de poule: une mine de molécules a activitésbiologiques. Cinquièmes Journées de la Recherche Avicole; 2003 Mar 26-27; Tours.
- North MO. Commercial chicken production manual. Connecticut: AVI Publishing; 1984. 710 p.
- Nys Y, Sauveur B. The nutritional value of eggs. *INRA Productions Animales* 2004;17:385-393.
- Samli HE, Agha A, Senkoylu N. Effects of Storage Time and Temperature on Egg Quality in Old Laying Hens. *The Journal of Applied Poultry Research* 2005;14:548–553.
- SAS INSTITUTE. SAS/STAT user's guide. Version 9. Cary, NC; 2001.
- Tyler C, Geake FH. Eggshell strength and its relationship to thickness, with particular reference to individuality in the domestic hen. *British Poultry Science* 1964;5:3-18.
- USDA. Egg grading manual. Washington: Agricultural Marketing Services, 1975.
- Vanghan MC, Adams JL. Quality of eggs offered to consumers in Lincoln, Nebraska. *Poultry Science* 1959;38:1257

