



Review

■ Author(s)

Wang Z^I
Deng X^{II}
Wang A^{III}
Liu R^I

^I College of Animal Science and Technology, Northwest A&F University, Yangling 712100, China

^{II} National Engineering Laboratory for Animal Breeding and the Key Laboratory of Animal Genetics and Breeding of the Ministry of Agriculture, China Agricultural University, Beijing, 100193, China

^{III} State Key Laboratory of Direct-Fed Microbial Engineering, Beijing 100192, China

■ Mail Address

Corresponding author e-mail address
Ruifang Liu
College of animal science and technology,
Northwest A&F University, Xinong Road,
22, Yangling, Shaanxi, 712100, China.
Phone/Fax: +86 (029) 87092102
E-mail: liuruifang79@163.com

■ Keywords

Chicken, blue egg, biliverdin, HMOX1, differential expression.

High Expression of HMOX1 in Blue-Shelled Chickens is Associated with a TG Haplotype

ABSTRACT

HMOX1 is an important gene in biosynthesis of the eggshell pigment of blue eggs. Previous studies found that *HMOX1* is highly expressed in the shell gland of hens laying blue eggs (BlueH) compared with hens laying brown eggs (BrownH); however, the reasons for the differential expression are unclear. In this study five single nucleotide polymorphism (SNP) in *HMOX1* were genotyped in 111 BlueH and 115 BrownH. The association of haplotypes of these SNP with the blue egg phenotype was tested. Haplotype-specific expression of *HMOX1* was detected in the shell gland. The interaction of sequence variants and transcription factors was analyzed using electrophoretic mobility shift assay (EMSA). A TG haplotype covering upstream 1.4 kb region of *HMOX1* was significantly associated with blue eggs ($p < 0.05$). Furthermore, the birds ($n=12$) with the haplotype expressed 3.8 fold more transcripts than those ($n=12$) without the haplotype ($p < 0.05$). After re-sequencing a 2.2 kb region harboring the TG haplotype, a total of 26 SNP were found, of which a SNP was predicted to create a binding site of Nrf2, a transcription factor initiating *HMOX1* expression. However, subsequent EMSA failed to confirm the Nrf2-DNA interaction. Taken together, the data suggested that the TG haplotype is not directly involved in regulation of *HMOX1* expression; a regulatory mutation located near the haplotype and linked with the haplotype may exist and be responsible for the differential expression of *HMOX1*.

INTRODUCTION

Birds are the only class of vertebrates that have evolved eggshell pigmentation (Cassey *et al.*, 2010). The eggshell colors have been proposed to serve a wide variety of visual, physical, physiological, and behavioral functions, such as cryptic coloring, resisting harmful solar radiation, thermal protection, reinforcing eggshell strength, sexual selection, etc. (Reynolds *et al.*, 2009). Thus, appearance of eggshell colors represents an important event of bird evolution. Blue-green coloration is a common eggshell color among birds. It is present not only in wild birds, but also in domestic fowls (i.e., several indigenous chicken breeds, such as Chilean Araucano, Chinese Dongxiang, and Lushi chickens) (Kennedy & Verves, 1973; Wang *et al.*, 2013). The blue-green coloration is believed to function as a signal of female health to attract males involved in the supervision of the offspring (Morales *et al.*, 2010).

Blue egg is colored by biliverdin, a bile pigment derived from the oxidative degradation of heme (Lang & Well, 1987). Two studies have confirmed that hens laying blue eggs (BlueH) have significant higher levels of biliverdin in eggshell and shell gland than those laying brown eggs (BrownH) (Zhao *et al.*, 2006; Wang *et al.*, 2009). However, molecular mechanisms underlying this difference are poorly understood. Blue-green eggshell color, also called as oocyan, is genetically controlled by a dominant mutant (*O*) (Punnett, 1933). In the latest study, Wang



et al. (2013) identified the *O* as a retrovirus (EAV-HP) insertion in *SLCO1B3*. Because OATP1B3 (gene protein) is a membrane transporter that can mediate transportation of biliverdin (Hagenbuch & Gui 2008), it is speculated that formation of blue egg may be relevant to abnormal transportation of the eggshell pigment in shell gland (Wang *et al.*, 2013).

In addition of *SLCO1B3*, two studies found higher *HMOX1* expression in the shell glands of BlueH than of BrownH (Wang *et al.*, 2010; Wang *et al.*, 2011). Because heme oxygenase-1 (HO-1, protein encoded by *HMOX1*) is the rate-limiting enzyme catalyzing oxidative degradation of heme into biliverdin (Maines, 1997), the above result implied that *HMOX1* can play a crucial role in biosynthesis of eggshell biliverdin.

The aim of this study was to further explore the regulatory mechanisms underlying high expression of *HMOX1* in the shell glands of BlueH. The association of *HMOX1* haplotype with blue egg phenotype was tested in the groups of BlueH and BrownH. Haplotype-specific expression of *HMOX1* was detected in the shell glands. The effect of sequence variants on *HMOX1* expression was studied using electrophoretic mobility shift assay.

MATERIAL AND METHODS

Birds and Sample Collection

In this study, Dongxiang chickens were used to explore how the regulatory mechanisms of *HMOX1* expression work. Dongxiang is a well-known blue egg-laying breed from Dongxiang city, Jiangxi province, China. Although blue egg phenotype has been specifically selected in the Dongxiang chickens, the trait has not been fixed in the group, in which most birds lay blue-green eggs, but there are still minor individuals laying light brown eggs. A total of 111 BlueH and 115

BrownH were used for haplotype association analysis. All birds were provided by Hualv Dongxiang chicken protection farm, Jiangxi province, China.

Twelve birds with TG haplotype (TG_) and 12 birds without the haplotype (XX) were euthanized approximately three to five h prior to ovulation. Shell glands were collected and immediately stored in liquid nitrogen for subsequent expression analysis. Use and care of birds in this study was approved by the Northwest A&F University Ethics Committee.

SNP genotyping and haplotype analysis

Five SNPs were used to construct the *HMOX1* haplotype. These SNPs, covering over 7 kb of the *HMOX1* genome sequence, are sufficient to capture most of the variation in *HMOX1*, as assessed using the Tagger program of Haploview 4.2 (Broad Institute, Cambridge, MA). Information on the SNPs is summarized in Table 1.

Blood samples were collected by wing vein puncture. Genomic DNA was extracted from the blood using a standard phenol-chloroform method. The SNPs were genotyped by polymerase chain reaction-restriction fragment length polymorphism. Restriction enzymes were summarized in Table 1. All fragments were resolved on 2.5 % agarose gels or 12% nondenaturing polyacrylamide gels and visualized after an ethidium bromide or silver staining.

All allele frequencies were tested for confirmation with the Hardy-Weinberg equilibrium. Construction and association analysis of the haplotype were completed using Haploview v4.2 under default values. Haplotypes with frequencies lower than 0.01 were discarded. In "sliding windows" analysis, 2-SNP, 3-SNP, 4-SNP window sizes were used to detect whether

Table 1 – Information on the five SNPs used in *HMOX1* haplotype association

Name	dbSNP Accession No ¹	Allele	SNP positions in UCSC ²	SNP positions in <i>HMOX1</i>	primer sequence (5'-3')	Restriction enzyme
HSNP1	rs314037822	G/T	Chr1:51956353	5' upstream region	F: AGCTCCAGATGGAAGAGG R: GGTGATAGAGGAGGACAGC	<i>Hinf</i> I
HSNP2	rs315838673	A/G	Chr1:51954906	intron1	F: CGCTCCTCCTCTCACCTCTG R: GCCGTGCTGTACACCCTAC	<i>Sau</i> I
HSNP3	rs313454440	G/A	Chr1:51954831	exon2, synonymous variant	F: GACCGGGAGGAGAAGCATGG R: ACACGGCTCTGGAACACC	<i>Sac</i> I
HSNP4	rs3137366	T/C	Chr1:51952346	exon3, V122A	F: GCAGACAAAAGACACACC R: GCTGTAACCCACAACCTC	<i>HPY</i> 99I
HSNP5	rs13866562	A/G	Chr1: 51949356	3'UTR	F: ACTGGGCTCTGCCACTAATA R: GAGCAAACAGAGCTCCGGGAG GTGCA ³	<i>Apa</i> L I

¹ dbSNP database: <http://www.ncbi.nlm.nih.gov/snp>

² SNP positions refer to galGal4 assembly of UCSC database (<http://genome.ucsc.edu>).

³ The reverse primer for rs13866562 was modified to create a specific sequence of restriction enzyme of *Apa*L I. The modified base is indicated in bold.



some regions can carry stronger association with blue egg than others.

Haplotype-specific expression analysis

Total RNA was extracted with TRIzol reagent (TianGen Co., Beijing, China). Two micrograms of total RNA were reverse transcribed in a final volume of 25 μ L, containing 1 μ L of 20 μ M Oligo (dT)15 primer, 5 μ L of first-strand buffer (250 mM Tris-HCl, 375 mM KCl, 15 mM MgCl₂, 50 mM DTT), 5 μ L of 10 mM dNTP, 1 μ L of 25 U Recombinant RNasin® Ribonuclease Inhibitor, 1 μ L of 200 U/ μ L M-MLV reverse transcriptase (Promega, Wisconsin, USA), and RNase-free water to final volume.

Real-time PCR was performed in an ABI PRISM 7900 instrument (Applied Biosystems Corp., Foster, CA). Amplification of 1 μ L of cDNA was carried out in a total volume of 20 μ L, according to the RealMasterMix (SYBR Green I) manual (Tiangen Corp., Beijing, China). Every sample was evaluated in triplicate. Data were analyzed by the 2^{- $\Delta\Delta$ CT} method (Livak & Schmittgen, 2001). The XX group was designed as calibrator. *GAPDH* was used as endogenous reference to normalize the amount of cDNA added to the PCR. The amount of *HMOX1* transcripts, normalized to *GAPDH* and relative to the calibrator, was given by 2^{- $\Delta\Delta$ CT}. The forward and reverse primers were 5'-ATCGCATGAAAACAGTC-CAG-3' and 5'-CAAATAAGCCCACGGCGAC-3' for *HMOX1* and 5'-ATACACAGAGGACCAGGTTG-3' and 5'-AAACTCATTGTCATACCAGG-3' for *GAPDH*.

Electrophoretic Mobility Shift Assays (EMSA)

The two biotin-labeled oligonucleotides used in EMSA reaction were 5'-CGCGGCTCTGCTCT**TC**CTCCTGCTGCCCGG-3' and 5'-CGCGGCTCTGCTCT**AA**CTCCTGCTGCCCGG-3', corresponding to two alleles (shown in bold and underlined letters) of TC and AA at TSNP21, respectively. Nuclear proteins were extracted from the shell glands using the Nuclei Protein Extraction Kit (Viagene Biotech Inc, Beijing, China), following the manufacturer's protocol. Protein concentration was determined with the BCA Protein Assay Kit (Cwbio Co. Beijing, China). Five μ g of nuclear proteins and 2.5 ng of biotin-labeled probes were incubated at room temperature for 20 min in binding mixture containing 1.5 μ L of 10 \times binding buffer and 1 μ g of poly(dI)•poly(dC) in a total volume of 15 μ L. In antibody supershift assays, 2 μ g of Anti-Nrf2 rabbit polyclonal antibody (Abcam Co., Shanghai, China) were added to the reaction mixture and incubated for 20 min at room temperature prior to electrophoresis. In competition assays, firstly a 100-fold molar excess of unlabeled oligonucleotides was added

to the binding mixture, then incubated for 30 min at room temperature. Secondly, labeled probes were added and subjected to additional 20 min incubation at room temperature. DNA-protein complexes were resolved on a 5 % nondenaturing polyacrylamide gel in 0.5 \times TBE at 80 V for 2-3 h. After electrophoresis, the complexes were transferred to a membrane and detected with horseradish peroxidase-conjugated streptavidin. Finally, blots were visualized using EMSA Chemiluminescent system (Viagene Biotech Inc, Beijing, China).

Statistical analysis

For comparison of *HMOX1* expression between birds with TG haplotype and those without the haplotype, the null hypothesis that there was no difference between two groups was tested by unpaired *t* test. Statistical significance was assumed at *p* < 0.05. All statistical analyses were done by SAS® V8.2 statistical software package (PC-SAS) (Cary, NC, USA).

RESULTS

Association of *HMOX1* haplotypes with blue egg

In order to elucidate whether there are sequence variants responsible for high expression of *HMOX1* in shell gland of BlueH, the association of *HMOX1* haplotypes, consisting of 5 SNPs, with blue egg phenotype was analyzed. Five major haplotypes were observed in BlueH and BrownH, of which only the GAGTG haplotype showed a significant association with blue eggs (*p*<0.05, Table 2). In order to explore whether some regions of *HMOX1* can carry a stronger association, 4-, 3- and 2-SNP "sliding window" analysis were performed. As the "sliding window" size was reduced, three haplotypes (TGGT, TGG, and TG) showed increasing association with blue egg, with the TG haplotype showing the strongest association with blue eggs (Table 3).

Table 2 – Frequencies of haplotypes constructed by the five SNPs

Haplotype ¹	Frequencies		Chi-square <i>P</i> value
	BrwonH ³	BlueH ³	
GGACA	0.218	0.216	0.9571
TAGTA	0.183	0.177	0.8696
TAGCG	0.113	0.158	0.1704
GAGCA	0.139	0.082	0.0578
GAGTG²	0.050	0.107	0.0212
all rare<0.05	0.285	0.217	0.0936

¹ Only haplotypes with frequencies \geq 0.05 are shown. Rare haplotypes with frequencies < 0.05 were combined.

² A significant result is highlighted in bold.

³ BrownH = hens laying brown eggs, BlueH = hens laying blue eggs

**Table 3** – Results of 2, 3 and 4-SNP “sliding window” analysis


Configurations	Haplotype ¹	Frequencies		Chi-square P value
		BrownH ²	BlueH ²	
HSNP2-HSNP3-HSNP4-HSNP5	GGTG	0.010	0.039	0.0432
HSNP2-HSNP3-HSNP4-HSNP5	GGCA	0.007	0.039	0.0206
HSNP2-HSNP3-HSNP4	GGC	0.006	0.040	0.017
HSNP1-HSNP2-HSNP3-HSNP4	TGGT	0.025	0.063	0.0489
HSNP1-HSNP2-HSNP3	TGG	0.026	0.077	0.0133
HSNP1-HSNP2	TG	0.029	0.099	0.0022

¹ Only haplotypes significantly associated with blue egg are shown in the table.

² BrownH = hens laying brown eggs, BlueH = hens laying blue eggs

Haplotype-specific expression analysis

The TG haplotype covers upstream 1.4 kb region of *HMOX1*, which is traditionally a key region for the regulation of gene expression. A haplotype-specific expression assay was performed using hens birds (n=12) with the TG haplotype (TG₋) and hens (n=12) without the haplotype (XX) to investigate the role of the haplotype in the regulation of *HMOX1* expression. The expression data showed that the TG₋ group expressed 3.8-fold more transcripts than the XX group in shell gland (p<0.05, Figure 1).

Figure 1 – Haplotype-specific expression analysis in shell gland


TG₋ group consisted of birds (n=12) with the TG haplotype, correspondingly XX for ones (n=12) without the haplotype. The results were presented as mean folds of *HMOX1* expression relative to the XX group. The TG group expressed approximate 3.8 fold more transcripts than the XX group (* p<0.05).

Effects of sequence variants within the TG-covering region on transcription factor-DNA interactions

A 2.2 kb region of Chr1:51954712-51956937, harboring the TG haplotype, was re-sequenced using 10 BlueH and 10 BrownH to identify crucial regulatory variants. A total of 26 SNPs were found in the region, most of which showed significant association with

blue eggs (Table 4). After performing a prediction of transcription factor binding sites (TFBS) using CONSITE software (<http://consite.genereg.net/>) (LENHARD *et al.* 2003), 11 SNPs of these SNPs were found to create a putative TFBS (Table 4), with special attention paid to TSNP21 because the blue egg-associated TC allele of the TSNP21 can create a putative TFBS of Nrf2. It has been well established that Nrf2 is a transcription factor initiating *HMOX1* expression under oxidative stress (Campbell *et al.*, 2013).

An EMSA assay, therefore, was performed, using nuclear extracts from the shell gland and probes corresponding to TC and AA alleles in order to investigate the authenticity of Nrf2-mediated expression. Similar shifted bands were detected in lanes 2 and 6 with TC and AA probes, suggesting that there was no obvious difference in transcription factor-binding activity between two TSNP21 alleles (Figure 2). After performing a supershift assay with anti-Nrf2 antibody, we also failed to observe supershifted bands in lanes 3 and 7 (Figure 2), which, therefore, implied that the protein-DNA complexes in lanes 2 and 6 may be produced by the binding of other proteins, instead of Nrf2, to the oligonucleotides.

Figure 2 – EMSA validated Nrf2-TSNP21 interaction

Nrf2 is a well-established transcription factor initiating *HMOX1* expression under oxidative stress. A putative Nrf2-binding element was created by the TC allele of the TSNP21 locus. EMSA was performed to validate the protein-DNA interaction using nuclear extracts from the shell gland and two probes corresponding to TC and AA alleles of TSNP21. The positions of free probe and shifted DNA-protein complexes are indicated with arrows. In lanes 1 and 5, no shifted band was observed when no nuclear extracts added; in lanes 2 and 6, the same shifted bands were observed using two probes, indicating that there was no obvious difference in transcription factor-binding activity between two alleles; in lanes 3 and 7, the shifted bands were not supershifted by Nrf2 antibody, indicating that the observed complexes in lanes 3 and 7 can consist of other protein, instead of Nrf2, and oligonucleotides; in lane 4 and 8, the shifted bands can be competed out by 100 × molar excess of unlabelled probes (cold probe), confirming that the complex consists of two target probes and unknown proteins.


Table 4 – Effect of SNPs within a TG-covering region of Chr1: 51954712-51956937 on transcript factor binding sites

Name	dbSNP accession No.	Position at Chr. 1 ¹	Position at <i>HMOX1</i>	Alleles	Chi-square P value ²	Effect of SNP on TFBS ³
TSNP1	rs314913481	51956882		A/G	p<0.05	no
TSNP2	rs312661490	51956662		A/G	p<0.05	no
TSNP3	new	51956591		T/C	p<0.05	no
TSNP4	new	51956521		G/A	p<0.05	no
TSNP5	rs315124462	51956427		C/T	p<0.05	C→T, FREAC-4
TSNP6	new	51956403		A/C	p<0.05	C→A, Thing1-E47
TSNP7	rs316233291	51956396		C/G	p<0.05	G→C, Myf
TSNP8	rs314166379	51956380		A/G	p<0.05	
HSNP1	rs314037822	51956353	5' upstream region	G/T	p<0.05	
TSNP9	rs315494484	51956346		C/T	p<0.05	
TSNP10	new	51956248		C/T	p<0.05	
TSNP11	new	51956226		C/T	p<0.05	
TSNP12	new	51955536		A/G	p>0.05	G→A, Max
TSNP13	new	51955474		C/T	p>0.05	C→T, c-FOS
TSNP14	rs14824170	51955419		C/T	p>0.05	T→C, Spz1
TSNP15	rs317039830	51955410		C/G	p>0.05	
TSNP16	new	51955322		C/T	p>0.05	T→C, ARNT
TSNP17	rs314520877	51955099		A/C	p<0.05	
TSNP18	new	51954974		A/G	p<0.05	
TSNP19	new	51954966		A/G	p<0.05	A: AML-1, G: Myf
TSNP20	new	51954944	intron1	C/T	p<0.05	C→T, Spz1
TSNP21	new	51954940-51954941		TC/AA	p<0.05	AA TC, Nrf2
HSNP2	rs315838673	51954906		A/G	p<0.05	
TSNP22	rs313881629	51954876		T/C	p<0.05	
TSNP23	rs313454440	51954831	Exon2, synonymous variant	A/G	p>0.05	
TSNP24	rs314101757	51954807	Exon2, synonymous variant	A/G	p>0.05	G→A, CHOP, Nrf2

¹ SNP positions at chromosome1 were given according to chicken genome galGal4 assembly in UCSC database (<http://genome.ucsc.edu/cgi-bin/hgGateway>).

² Association of SNP alleles with blue egg was tested by Chi-square.

³ A putative transcript factor binding sites (TFBS) was created with alternation of alleles, which was predicted using the CONSITE software (<http://asp.ii.uib.no:8090/cgi-bin/CONSITE/consite>).

DISCUSSION

Biliverdin, a bile pigment, previous regarded as a waste product of heme breakdown, actually has an important role in scavenging free radicals (Bulter & Mcgraw, 2013). In birds, the bile pigment also plays the role of pigmentation of the eggshell with blue-green color (Duval *et al.*, 2013). Biliverdin is derived from the oxidative degradation of heme, with the formation of equimolar CO and Fe²⁺. In this reaction, heme oxygenase (HO) is the rate-limiting enzyme (Maines, 1997). Previous studies have found that HO-1 (one of three HO isozymes) have a higher level of expression in shell the gland of BlueH than BrownH (Wang *et al.*, 2010; Wang *et al.*, 2011), which therefore suggested that HO-1 is most likely to play a role in the expression of the blue egg phenotype as well. However, the regulatory mechanism underlying the differential expression remains unclear. This study found a TG haplotype to be significantly associated with blue egg phenotype. Subsequent haplotype-specific expression

analysis further revealed that the haplotype has a higher expression activity, which implied that a cis-regulation variant may exist in the *HMOX1* and be responsible, or partly at least, for the differential expression of *HMOX1* between BlueH and BrownH.

To find out the variant, we sequenced a 2.2 kb sequence covering the TG haplotype and found a suspicious SNP, which can create a putative binding site of Nrf2. However, EMSA failed to verify the authenticity of the Nrf2-DNA interaction. This urges us to rethink the role of the TG haplotype in regulation of *HMOX1* expression. It is plausible that the phenotype-associated haplotype itself is not involved directly in regulation of *HMOX1* expression; it may only serve as a marker indicating that an adjacent mutation, situated in linkage disequilibrium with it, may mediate the high *HMOX1* expression in BlueH. The inference obtained a support from studies to *cis*-acting elements of *HMOX1*. HO-1 is an inducible cytoprotective protein in the defense system against oxidative stress (Calay & Mason, 2014). A long list



of *cis*-regulatory elements, such as ARE, HeRE, HSE, CdRE, MARE, MPRE, and (GT)_n repeats, were found to be involved in regulation of HMOX1 expression in human, rat and chicken (Martin *et al.*, 2004; Chang *et al.*, 2004; Shibahara *et al.*, 1989; Shibahara *et al.*, 2003; Shan *et al.*, 2004). Thus, it is conceivable that any variants in these *cis*-elements may be responsible for the differential expression as well.

In the case of trans-acting factors three MAPK pathways of ERK, JNK, and p38 are known to be involved in up-regulation of HO-1 expression (Martin *et al.*, 2004). In addition, trans-acting protein BACH1 is a repressor of Nrf2-mediated HO-1 expression. Its inactivation is a prerequisite for HO-1 induction (Reichard *et al.*, 2007). Thus, any function or expression alterations in these trans-regulatory factors can also become the reasons for the differential expression of HMOX1.

Chicken oviduct is a highly vascular organ. When oocytes leave the ovary and enter the oviduct, the metabolism of the oviduct is stimulated into activity; blood flow through it increases; albumen is largely synthesized and secreted (Kennedy & Ververs, 1973; Baird *et al.* 1975). Following the process, a large amount of reactive oxygen species will be produced, causing deleterious effects on cellular proteins, lipids, and DNA (Alonso-Alvarez *et al.*, 2004). HO-1 is a crucial protein in resisting oxidative injury (Calay & Mason, 2014). Thus, its high expression in BlueH may imply that appearance of blue egg is a result of bird adaptive evolution in the defense against oxidative stress during lay.

Based on previous studies of HMOX1, this work attempted further to reveal the reasons for the high expression of the gene in BlueH. Here, a TG haplotype was found to represent high expression activity. Because the haplotype appears more frequently (0.099) in BlueH than in BrownH (0.029), the data explained the reasons for high expression of HMOX1 to some extent. However, given the complex regulatory mechanisms of HMOX1, it remains an enormous challenge to define which of the variants is responsible for its differential expression. It, therefore, is necessary further to explore functional variants in an extended region and several trans-acting factors in the future.

ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China (Grant No. 31401051) and China Postdoctoral Science Foundation (2014M550510).

REFERENCES

- Alonso-Alvarez C, Bertrand S, Devevey C, Prost J, Faivre B, Sorci G. Increased susceptibility to oxidative stress as a proximate cost of reproduction. *Ecology Letters* 2004;7(5): 363-368.
- Baird T, Solomon SE, Tedstone DR. Localization and characterization of eggshell porphyrin in several avian species. *British Poultry Science* 1975;16(2):201-208.
- Bulter MW, McGraw KJ. Immune function is related to adult carotenoid and bile pigment levels, but not to dietary carotenoid access during development, in female mallard ducks. *Journal of Experimental Biology* 2013;216(Pt 14):2632-2640.
- Calay D, Mason JC. The multifunctional role and therapeutic potential of HO-1 in the vascular endothelium. *Antioxidant & Redox Signaling* 2014;20(11):1789-1809.
- Campbell MR, Karaca M, Adamski KN, Chorley BN, Wang XT, Bell DA. Novel hematopoietic target genes in the NRF2-mediated transcriptional pathway. *Oxidative Medicine and Cellular Longevity* 2013;2013(2013):1-12.
- Cassey P, Portugal SJ, Maurer G, Ewen JG, Boulton RL, Hauber ME, Blackburn TM. Variability in avian eggshell colour: a comparative study of museum eggshells. *PLoS ONE* 2010;5(8):e12054.
- Chang KW, Lee TC, Yeh WI, Chung MY, Liu CJ, Chi LY, Lin SC. Polymorphism in heme oxygenase-1 (HO-1) promoter is related to the risk of oral squamous cell carcinoma occurring on male areca chewers. *British Journal of Cancer* 2004;91(8):1551-1555.
- Duval C, Cassey P, Mikšik I, Reynolds SJ, Spencer KA. Condition-dependent strategies of eggshell pigmentation: an experimental study of Japanese quail (*Coturnix coturnix japonica*). *Journal of Experimental Biology* 2013;216(Pt4):700-708.
- Hagenbuch B, Gui C. Xenobiotic transporters of the human organic anion transporting polypeptides (OATP) family. *Xenobiotica* 2008;38(7-8):778-801.
- Kennedy GY, Ververs HG. Eggshell pigments of the Araucano fowl. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* 1973;44(1):11-25.
- Lang MR, Well JW. A review of eggshell pigmentation. *Worlds Poultry Science Journal* 1987;43(3):238-246.
- Lenhard B, Sandelin A., Mendoza L, Engstrom P, Jareborg N, Wasserman WW. Identification of conserved regulatory elements by comparative genome analysis. *Journal of Biology* 2003;2(2):13.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 2001;25(4):402-408.
- Maines MD. The heme oxygenase system: a regulator of second messenger gases. *Annual Review of Pharmacology and Toxicology* 1997;37:517-554.
- Martin D, Rojo AI, Salinas M, Diaz R, Gallardo G, Alam J, *et al.* Regulation of heme oxygenase-1 expression through the phosphatidylinositol 3-kinase/Akt pathway and the Nrf2 transcription factor in response to the antioxidant phytochemical carnosol. *Journal of Biological Chemistry* 2004;279(10):8919-8929.
- Morales J, Torres R, Velando A. Parental conflict and blue egg coloration in a seabird. *Naturwissenschaften* 2010;97(2):173-180.
- Punnett RC. Genetic study in poultry-IX. The blue egg. *Journal of Genetics* 1933;27(3):465-470.



- Reichard JF, Motz GT, Puga A. Heme oxygenase-1 induction by NRF2 requires inactivation of the transcriptional repressor BACH1. *Nucleic Acids Research* 2007;35(21):7074-7086.
- Reynolds SJ, Martin GR, Cassey P. Is sexual selection blurring the functional significance of eggshell coloration hypotheses? *Animal Behaviour* 2009;78(1):209-215.
- Shan Y, Lambrecht RW, Bonkovsky HL. Identification of key elements that are responsible for heme-mediated induction of the avian heme oxygenase-1 gene. *Biochimica et Biophysica Acta-Gene Structure and Expression* 2004;1679(2):87-94.
- Shibahara S, Sato M, Muller RM, Yoshida T. Structural organization of the human heme oxygenase gene and the function of its promoter. *European Journal of Biochemistry* 1989;179(3):557-563.
- Shibahara S, Nakayama M, Kitamuro T, Uono-Fujimori R, Takahashi K. Repression of heme oxygenase-1 expression as a defense strategy in humans. *Experimental Biology and Medicine* 2003;228(5):472-473.
- Wang XT, Zhao CJ, Li JY, Xu GY, Lian LS, Wu CX, Deng XM. Comparison of the total amount of eggshell pigments in Dongxiang brown-shelled eggs and Dongxiang blue-shelled eggs. *Poultry Science* 2009;88(8):1735-1739.
- Wang XT, Zhao CJ, Li JY, Xu GY, Lian LS, Wu CX, Deng XM. Heme Oxygenase-1 is important to the formation of eggshell biliverdin in chicken. *Journal of Applied Animal Research* 2010;38(2):229-232.
- Wang ZP, Liu RF, Wang AR, Li JY, Deng XM. Expression and activity analysis reveal that heme oxygenase (decycling) 1 is associated with blue egg formation. *Poultry Science* 2011;90(4):836-841.
- Wang ZP, Qu LJ, Yao JF, Yang XL, Li GQ, Zhang YY, *et al.* An EAV-HP insertion in 5' flanking region of SLCO1B3 causes blue eggshell in the chicken. *PLoS Genetics* 2013;9(1):e1003183.
- Zhao R, Xu GY, Liu ZZ, Li JY, Yang N. A study on eggshell pigmentation: biliverdin in blue-shelled chickens. *Poultry Science* 2006;85(3):546-549.

