



## Effect of Antimicrobials on *Salmonella* Spp. Strains Isolated from Poultry Processing Plants

### ■ Author(s)

Mion L<sup>1</sup>  
Parizotto L<sup>1</sup>  
Calasans M<sup>1</sup>  
Dickel EL<sup>1</sup>  
Pilotto F<sup>1</sup>  
Rodrigues LB<sup>1</sup>  
Nascimento VP<sup>1</sup>  
Santos LR<sup>1</sup>

<sup>1</sup> Curso de Medicina Veterinária –  
Universidade de Passo Fundo

### ■ Mail Address

Corresponding author e-mail address  
Luciana Ruschel dos Santos  
Curso de Medicina Veterinária –  
Universidade de Passo Fundo – BR 285  
99052-900. São José, Passo Fundo, RS, Brazil  
E-mail: [luruschel@upf.br](mailto:luruschel@upf.br)

### ■ Keywords

Antimicrobials, poultry, sanitizers, *Salmonella* spp.

### ABSTRACT

The routine use of antimicrobials in animal production for the treatment of infections, disease prevention, or as growth promoters is a predisposing factor for the development and dissemination of antimicrobial resistance. In food industries, sanitizers are used for the control of microbial colonization, and their efficacy depends on contact time and on the dilution of the products used. The present study assessed the effect of 12 antimicrobials and four commercial sanitizers on 18 *Salmonella* spp. strains isolated from poultry processing plants. None of the evaluated antimicrobials was 100% effective against the tested *Salmonella* spp. strains; however, 94% of the isolates were susceptible to ciprofloxacin, 77% to amoxicillin + clavulanic acid and to ampicillin, and 72% to enrofloxacin, whereas 100% of the isolates were resistant to penicillin G, 16% to tetracycline, and 11% to sulfonamide. The tested *Salmonella* spp. strains were 100% inhibited by peracetic acid after five minutes of contact, 0.5% by quaternary ammonium after 15 minutes, and 85.7% by chlorhexidine after 15 minutes. The results indicate the importance of testing of efficacy of antimicrobials used in animal production and in public health to monitor their action and the development of resistance.

### INTRODUCTION

The routine use of antimicrobials in animal production for the treatment of infections, disease prevention, and as growth promoters, may lead to the development and dissemination of antimicrobial resistance, which is later transmitted to human beings in the food chain (EFSA, 2014).

The sanitation standard operating procedures (SSOP) applied in processing plants aim at controlling the dissemination of microorganisms and the formation of biofilm on the surfaces with which the food has contact; however, bacteria of the genus *Salmonella* are known for their adhesion to surfaces and resistance to sanitizers (Rodrigues *et al.*, 2013). The most commonly products used in SSOP in poultry processing plants are peracetic acid, quaternary ammonium, and chlorhexidine. The selection and use of sanitizers are crucial for the reduction of microbial counts. However, the degree of hygiene and type of surfaces of processing plants bacterial growth (Baltreme, 2014), and should be known when testing the efficacy of sanitizers applied in such facilities.

This study evaluated the efficacy of 12 antimicrobials and four commercial sanitizers on 18 *Salmonella* spp. strains isolated from poultry processing plants.

### MATERIALS AND METHODS

*Salmonella* spp. strains were isolated from seven broiler processing plants under federal inspection services in the state of Rio Grande do



Sul, Brazil, between 2012 and 2014. A total of 1,071 samples were collected, out of which 18 (1.68%) were positive for *Salmonella* spp. The samples positive for *Salmonella* spp were determined in cloacal swabs (n=6), transport crate swabs after washing (n=3), carcasses after plucking (n=1), carcasses after the first wash (n=1), plucked and washed carcasses (n=1), eviscerated carcasses (n=1), at the exit of the chiller (n=1), after final wash (n=1), carcasses chilled at 4°C (n=1), carcasses frozen at -12° for 24 hours (n=1), and carcasses frozen at -12°C for 60 days (n=1). The samples were processed at the Laboratory of Bacteriology and Mycology of the Veterinary Hospital of the University of Passo Fundo (HV-UPF), Brazil, using *Salmonella* Enteritidis ATCC 13076 as positive control.

Cloacal swabs were taken from 300 broilers in each collection, using one swab for every two birds, making up a pool of 50 swabs that were stored in flasks containing 50 mL of buffered peptone water (BPW 1.0%, HiMedia®). In the laboratory, pools were homogenized and 10-mL aliquots were used for *Salmonella* spp isolation.

Transport crates were sealed with official labels and sampled by rubbing 3M® sponge sticks with neutralizing buffer across the inner side of the crate before and after washing and disinfection. The sponges were placed the collection bags provided by the manufacturer and 50 mL of 1.0% BPW was added; a 10-mL aliquot was used for *Salmonella* spp isolation.

Birds and carcasses were placed in individual plastic bags sealed with official labels and rinsed with 400 mL of 1.0% BPW, and a 10-mL aliquot was used for analyses.

For the isolation of *Salmonella* spp., 10-mL aliquots of 1.0% BPW were incubated at 37±1°C for 16 to 20 hours. Subsequently, 1 mL was inoculated in 9 mL tetrathionate broth (Merck®) and incubated at 37±1°C for 24±3 hours, and 100 µL were inoculated in 9.9 mL Rappaport-Vassiliadis broth (Merck®) and incubated at 41.5±1°C for 24±3 hours. The samples were then streaked onto Rambach Agar (Merck®) and Brilliant Green Agar supplemented with Novobiocin (Merck®), and incubated at 37±1°C for 24±3 hours. Suspected *Salmonella* spp. colonies were transferred to Triple Sugar Iron (TSI) agar (Merck®), Lysine Iron Agar (Merck®), Sulfite Indole Motility (SIM) medium (Merck®), Urea Broth (Merck®), and then confirmed by Poly O Antiserum (Probac®).

For the antimicrobial sensitivity tests (CLSI, 2012), *Salmonella* spp isolates were incubated in BHI broth (HiMedia®) at 36±1°C for 16 to 18 hours. A


suspension equivalent to a 0.5 McFarland standard was obtained by dilution in BHI broth and used for the inoculation of tested bacteria onto Mueller-Hinton agar (Oxoid®) plates. The assessed antimicrobials (Laborclin®) were selected because are regularly used to treat human salmonellosis and farm animals, either as therapeutic agents or as growth promoters, and included amoxicillin + clavulanic acid (30µg), ampicillin (10µg), ceftiofur (30µg), chloramphenicol (30µg), enrofloxacin (5µg), streptomycin (10µg), gentamicin (10µg), neomycin (30µg), benzyl penicillin (10u), sulfonamide (300µg), ciprofloxacin (5µg), and tetracycline (30µg). After incubation at 36±1°C for 16 to 18 hours, results were interpreted according to a specific table (Laborclin®) and as recommended by Clinical and Laboratory Standards Institute (USDA, 2012). Multiresistance was determined according to the National Antimicrobial Resistance Monitoring System criteria (USDA, 2012) were applied, according to which multiresistance is the resistance to three or more classes of antimicrobials.

For testing the efficacy of sanitizers, according to Beltrame (2009), sanitizers were tested at the dilutions routinely used in processing plants, as follows: 0.5% chlorhexidine, 0.5% and 1% quaternary ammonium, and 1% peracetic acid, were utilized. The bacteria previously stored at -20°C in BHI broth with 20% glycerin were inoculated in BHI broth (HiMedia®) and incubated at 37°C for 18 hours. Subsequently, 100µL of the *Salmonella* spp. culture were added to sterile tubes containing 9 mL of the sanitizer at the tested concentration and 1 mL of UHT whole milk (to simulate the presence of organic matter). After contact times of 1, 5, 10 and 15 min, 10-µL aliquots were transferred to 5.0 mL of BHI broth and incubated for 96 hours at 37°C. The bacteria were considered resistant (R) when the culture medium presented turbidity, film formation on the surface, or precipitation, and susceptible (S) in the absence of turbidity. In this case, positive samples were seeded onto brilliant green agar plates to check bacterial viability.

## RESULTS AND DISCUSSION

None of the tested antimicrobials was 100% effective against the isolated *Salmonella* spp. (Figure 1).

Chloramphenicol is used to treat human salmonellosis, due to its low cost and adequate therapeutic response (Alecrim *et al.*, 2002). Colla *et al.* (2012) reported 25.6% resistance and 64.1% of intermediate resistance to this drug in *Salmonella* Typhimurium and Panama serovars, respectively,



**Figure 1** – Antimicrobial sensitivity of 18 *Salmonella* spp. samples isolated from poultry processing plants.

isolated from swine carcasses, while Mion *et al.* (2014) observed 100% efficacy of chloramphenicol, enrofloxacin, ciprofloxacin, and streptomycin against *Salmonella* Heidelberg isolated from poultry processing plants between 2005 and 2009. However, in the present study, only ciprofloxacin was more than 90% effective against *Salmonella* spp isolated between 2012 and 2014, suggesting a possible development of resistance of this bacterium during this period.

Tetracyclines are one of the most widely classes of antimicrobials therapeutically in livestock (Wilson, 2004). In the present study, 61.1% of the isolates were resistant to this drug, indicating that its continuous use has increased microbial resistance, consequently reducing the available therapeutic options. Tetracyclines and aminoglycosides (gentamicin and streptomycin), amoxicillin with clavulanic acid, ampicillin, and ciprofloxacin are considered by the World Health Organization to be critically important for human medicine (WHO, 2011). The resistance of bacteria to such drugs has increased (EFSA, 2014), and represent a substantial cost to public health as resistant bacteria are more harmful to patients than susceptible strains of the same species (Balsalobre *et al.*, 2014). Therefore, increasing rates of multi-resistant bacteria represent a potential public health hazard (Chiappini *et al.* 2002, Zimmermann 2008).

In a survey carried out in 2010 in the European Union, *Salmonella* spp. isolated from poultry meat and showed 27%, 24%, 24%, 21%, 20%, 4%, 3% and 2% resistance to sulfonamide, ciprofloxacin, nalidixic acid, ampicillin, tetracycline, cefotaxime,

chloramphenicol, and gentamicin respectively, which are similar to the rates detected in *Salmonella* strains isolated from humans. In the European Community, fluoroquinolones are used as first-line treatment against salmonellosis in adults, while third-generation cephalosporins are used in children, yielding resistant strains due to misuse and resulting in inefficient therapy, also causing multiresistance (EFSA 2014).

Drug multiresistance is described by the National Antimicrobial Resistance Monitoring System (USDA, 2012) as resistance to three or more classes of antimicrobials, and the ACSSuT R-type stands for the resistance to ampicillin (A), chloramphenicol (C), streptomycin (S), sulfonamide (Su), and tetracycline (T) (Reis *et al.*, 2011). In the present study, a *Salmonella* spp. isolate from a cloacal swab presented intermediate resistance to ampicillin and resistance to ceftiofur, chloramphenicol, streptomycin, gentamicin, neomycin, penicillin g, sulfonamide, and tetracycline, suggesting that broiler flocks on the farm may host ACSSuT multiresistant *Salmonella* strains (Figure 2).

Considering the possible transfer of such resistance to the human population, the European Union banned the use of antibiotics as growth promoters in animal production as of January 2006. Because Brazil is one of the main poultry exporters to the EU, Brazilian companies needed to comply with EU legislation (Lorenço *et al.*, 2007). In July 2009, the Brazilian Ministry of Agriculture (Brasil, 2012) banned the use of amphenicols, tetracycline, beta-lactams (benzylpenicillins and cephalosporins), quinolones, and sulfonamides as performance enhancers or




Figure 2 – Multiresistance of 18 *Salmonella* spp. samples isolated from poultry processing plants according to the number of antibiotics (ATBs)

as food preservers, limiting their use to veterinary treatments. Finally, in May 2012, the use of spiramycin and erythromycin as performance enhancers in animal production was also banned (Brasil, 2012).

The *in-vitro* assessment of the effect of commercial sanitizers against *Salmonella* spp. strains showed that only peracetic acid was 100% effective after 5 min of contact (Figure 3). Peracetic acid quickly acts against bacteria and it is more effective against biofilm formation because of its higher inhibitory potential of bacteria and in a shorter time of exposure compared with other sanitizers (Rodrigues *et al.*, 2013). The stronger inhibitory effect of 1% quaternary ammonium (42.9% to 100%) compared with 0.5% (57.2% to 85.7%), as shown in Figure 3, emphasizes the need to test the efficacy of disinfectants sold by different manufacturers, as commercial formulations may present different dilutions of the active ingredients. The weak effect of chlorhexidine may be due to its continuous use in dental, medical, and veterinary treatments since the 1950s, producing resistance because of its prolonged use and its inappropriate contact time and dilutions (Colla *et al.*, 2014). These results highlight that the effect of sanitizers against *Salmonella* strains is associated with practical situations, particularly with the presence of organic matter, contact time, and with the resistance developed by bacteria to the most common disinfectants.

## CONCLUSION

None of the tested antimicrobials was 100% effective against the *Salmonella* spp. strains. Peracetic acid presented the best sanitizing performance. These results show that strategies to reduce contamination risks in the food chain based in careful use of antimicrobial drugs and monitoring of resistance to antimicrobials and sanitizers in animal production need to be applied.

## ACKNOWLEDGMENTS

We thank the Research Incentive Foundation of the State of Rio Grande do Sul (FAPERGS – Process 12/2312-4) for the financial support.

## REFERENCES

- Alecrim WD, Loureiro ACSP, Moraes RS, Monte RL, Lacerda MVG. Febre tifóide: recaída por resistência antimicrobiana. Relato de caso. Revista da Sociedade Brasileira de Medicina Tropical 2002;35:661-663.
- Balsalobre LC, Dropa M, Matté MH. An overview of antimicrobial resistance and its public health significance. Brazilian Journal of Microbiology 2014;45:1-5.
- Beltrame CA. Avaliação da eficiência de sanitizantes utilizados pelas indústrias de alimentos. Erechim: Universidade Regional Integrada do Alto Uruguai e das Missões; 2009.




Figure 3 – Effect of commercial sanitizers and different contact times against 18 *Salmonella* spp. samples isolated in poultry processing plants.



- Beltrame CA. Avaliação da adesão e remoção de biofilmes de *Listeria monocytogenes* e *Escherichia coli* em diferentes superfícies e indução de resistência bacteriana, com diferentes sanitizantes [tese]. Erechim (RS): Universidade Regional Integrada do Alto Uruguai e das Missões; 2011.
- Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa nº 14, de 18 de maio de 2012. Proíbi em todo o território nacional a importação, fabricação e o uso das substâncias antimicrobianas espiramicina e eritromicina com finalidade de aditivo zootécnico melhorador de desempenho na alimentação animal [cited 2012 maio 28]. Available from: [http://www.agricultura.gov.br/arq\\_editor/file/CRC/IN%20142012%20%20proibe%20antimicrobianos%20espiramicina%20e%20eritromicina%20melh%20desemp.pdf](http://www.agricultura.gov.br/arq_editor/file/CRC/IN%20142012%20%20proibe%20antimicrobianos%20espiramicina%20e%20eritromicina%20melh%20desemp.pdf)
- Chiappini E, Galli L, Pecile P, Vierucci A, Martino M. Results of a 5-year prospective surveillance study of antibiotic resistance among *Salmonella* enterica isolates and ceftriaxone therapy among children hospitalized for acute diarrhea. *Clinical Therapeutics* 2002;24:1585-1594.
- Colla FL, Rodrigues LB, Dickel EL, Borsoi A, Nascimento VP, Santos LR. Avaliação *in vitro* de clorexidina, amônia quaternária e ácido peracético frente a amostras de *Salmonella* Heidelberg isoladas de abatedouro avícola em 2005 e 2009. *Pesquisa Veterinária Brasileira* 2012;32:289-292.
- Colla FL, Mion L, Parizotto L, Santos LA, Pilotto F, Rodrigues LB, et al. Perfil de sensibilidade aos antimicrobianos e eficácia de sanitizantes frente aos isolados de *Salmonella* spp. oriundos de carcaças suínas no Rio Grande do Sul. *Pesquisa Veterinária Brasileira* 2014;34:320-324.
- EFSA. Scientific opinion of the panel on biological hazards on a request from the European food safety authority on foodborne antimicrobial resistance as a biological hazard [cited 2014 Ago 20]. *Esa Journal* 765:1-87, 2005. Available from: [www.efsa.europa.eu/efsajournal](http://www.efsa.europa.eu/efsajournal).
- Lorençon L, Nunes RV, Pozza PC, Pozza MSS, Appelt MD, Silva WTM. Utilização de promotores de crescimento para frangos de corte em rações fareladas e peletizadas. *Acta Scientiarum* 2007;29:151-158.
- Mion L, Colla FL, Cisco IC, Webber B, Diedrich LN, Pilotto F, et al. Perfil de resistência a antimicrobianos por *Salmonella* Heidelberg isoladas de abatedouro avícola em 2005 e 2009. *Acta Scientiae Veterinariae* 2014;42:1197.
- Reis EMF, Rodrigues DP, Almeida ACF, Hofer E. Prevalence of R-type ACSSuT in strains of *Salmonella* serovar Typhimurium DT193 isolated from human infections in Brazil. *Pan American Journal of Public Health* 2011;29:387-392.
- Rodrigues LB, Santos LR, Rizzo NN, Tagliari VZ, Trenhago G, Oliveira AP, et al. *Salmonella* and *Listeria* from stainless steel, polyurethane and polyethylene surfaces in the cutting room of a poultry processing plant. *Acta Scientiae Veterinariae* 2013;41:1164.
- USDA. Performance standards for antimicrobial susceptibility testing: twenty-second information supplement. *Clinical and Laboratory Standards Institute M100-S22* 2012;32(3):1-184.
- USDA. National Antimicrobial Resistance Monitoring System (NARMS): Enteric bacteria, animal arms [Animal Arm Annual Report]. Athens: Department of Agriculture, Agricultural Research Service; 2012.
- Wilson IG Antimicrobial resistance of *Salmonella* in raw retail chickens, imported chicken portions, and human clinical specimens. *Journal of Food Protection* 2004;67:1220-1225.
- WHO - World Health Organization. Critically important antimicrobials for human medicine. Copenhagen: Report of the 3rd WHO expert meeting, 2011 [cited 2014 ago22]. Available from: [http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485_eng.pdf)
- Zimmermann FC, Pinheiro D, Cesco MAO, Guayba J, Borsoi A, Fortes BFB, et al. Resistência antimicrobiana em amostras de *Salmonella* Hadar isoladas de carcaças de frango [cited 2014 Jan 24]. *Anais do 35º Congresso Brasileiro de Medicina Veterinária; 2008; Porto Alegre. Rio Grande do Sul. Brasil.* Available from: <http://www.sovergs.com.br/conbravet2008/anais/cd/resumos/R0750-1.pdf>

