

Complications that increase the time of Hospitalization at ICU of patients submitted to cardiac surgery

Complicações que aumentam o tempo de permanência na unidade de terapia intensiva na cirurgia cardíaca

Artur LAIZO¹, Francisco Eduardo da Fonseca DELGADO², Glauco Mendonça ROCHA³

RBCCV 44205-1169

Abstract

Objective: To present the complications that prolong the stay in the Intensive Care Unit (ICU) of the patients undergoing cardiac surgery.

Methods: We retrospectively analyzed records of 85 patients undergoing cardiac surgery during the period of March to May 2009, at the Santa Casa de Misericórdia de Juiz de Fora (MG), and, of these patients we studied 14 (16.47%) who stayed for more than five days in ICU. Amongst the 85 patients, there were three deaths, being two patients operated on an emergency basis, which increases morbidity, and one patient who remained in the hospital and under mechanical ventilation (MV) for 21 days.

Results: The study showed that complications that increased the length of stay in ICU were respiratory and metabolic disorders, according to the literature.

Conclusion: The complications that increase length of stay in ICU are related to respiratory function, chronic obstructive pulmonary disease, smoking, lung congestion, prolonged switch-off from MV, diabetes, infections, kidney failure, stroke and hemodynamic instability.

Keywords: Postoperative complications. Cardiovascular surgical procedures. Length of stay. Intensive care units.

Resumo

Objetivo: Apresentar as complicações que aumentam a permanência na Unidade de Terapia Intensiva (UTI) dos pacientes submetidos à cirurgia cardíaca.

Métodos: Foram analisados, retrospectivamente, 85 prontuários de pacientes submetidos à cirurgia cardíaca, no período de março a maio de 2009, na Santa Casa de Misericórdia de Juiz de Fora (MG) e, destes pacientes, foram estudados 14 (16,47%) que permaneceram por mais de 5 dias na UTI. Dentre os 85 pacientes, houve três óbitos, sendo dois pacientes operados em caráter de urgência, o que aumenta a morbidade, e um paciente que permaneceu internado e em ventilação mecânica (VM) por 21 dias.

Resultados: O estudo demonstrou que as complicações que aumentaram o tempo de internação na UTI foram respiratórias e metabólicas, de acordo com a literatura.

Conclusão: As complicações que aumentam o tempo de permanência na UTI são as relacionadas à função respiratória, doença pulmonar obstrutiva crônica, tabagismo, congestão pulmonar, desmame da VM prolongado, diabetes, infecções, insuficiência renal, acidente vascular encefálico e instabilidade hemodinâmica.

Descritores: Complicações pós-operatórias. Procedimentos cirúrgicos cardiovasculares. Tempo de internação. Unidades de terapia intensiva.

1. Master in General Surgery, UFMG, Professor, University Presidente Antonio Carlos - Campus VI - Juiz de Fora.
2. Ph.D. in Animal Science at UFMG, Professor of UNIPAC - JF.
3. Medicine Academic of UNIPAC - campus VI - Juiz de Fora.

Correspondence address: Artur Laizo
Avenida Rio Branco, 2644, apto 302 - Centro - Juiz de Fora, MG, Brazil.
E-mail: artur93@hotmail.com

Work performed at Santa Casa de Misericórdia de Juiz de Fora, Juiz de Fora, MG, Brazil.

Article received on December 17th, 2009
Article accepted on April 26th, 2010

INTRODUCTION

Cardiac surgeries are major surgeries broadcasted worldwide, among them stands out mostly coronary artery bypass (CABG) and valve replacement [1-4].

Patients undergoing cardiac surgery undergo a series of examinations and preoperative tests, so that there are no surprises and complications. The procedure has high morbidity and has complications related to the preoperative and cardiopulmonary bypass (CPB) used during the operation, requiring that patients undergoing these procedures are well prepared hemodynamically and psychologically for the postoperative [5-10].

The morbidity and mortality in the postoperative of cardiac surgeries is of great interest, prompting several protocols for postoperative management [1-4]. The ischemic asymptomatic episodes are very frequent and seem to imply a worse prognosis [11,12]. Thoracotomy is an intervention that will limit movements, and the position in bed leads to pain in varying degrees.

The patient undergoing cardiac surgery shall remain on mechanical ventilation (MV) in the immediate postoperative period until he covers total lucidity, in some cases for even longer, and should remain in bed for three to six weeks [3,12-14]. Because of hemodynamic instability that can happen in the immediate postoperative period, the patient may need inotropic support (VAD) and must remain in the Intensive Care Unit (ICU) while there is need [15-17]. The controls of vital data, such as capillary blood glycemia and monitoring are performed on an hourly basis, which can also bring discomfort to the patient [2, 18-21].

Systemic diseases, especially respiratory and metabolic diseases such as chronic obstructive pulmonary disease (COPD) and diabetes, may increase the length of stay in ICU and in some cases, increase the duration of mechanical ventilation [3,9,18]. Patients with COPD present, if not well-conditioned and have not stopped smoking for at least four weeks before surgery, greater difficulty in oxygenation and, therefore, remain longer in the VM [22,23]. Diabetes is controlled with capillary blood glycemia hourly and regular insulin according to glycemia. There are cases where it is necessary to apply intravenous insulin therapy and the support of endocrinology in the ICU [24,25].

Diseases such as hypertension, autoimmune diseases, peripheral vascular disorders, metabolic syndrome (MS) and others need to be controlled and require greater care in the immediate postoperative period [26].

The aim of this paper is to describe the complications that increase length of stay in ICU in the immediate postoperative period of cardiac surgeries.

METHODS

We evaluated medical records of 85 patients undergoing cardiac surgery at Santa Casa de Misericordia de Juiz de Fora - MG (SCMJF) in the surgical ICU during the period of March to May 2009.

The study was approved by the Ethics and Research Committee of Santa Casa de Misericordia de Juiz de Fora, with Protocol 0079/09.

Statistical analysis

We calculated the mean and standard deviation of length of stay and patients' age (Figures 1 and 2), to verify, descriptively, the behavior of the sample according to the period of hospitalization of patients. The inferential statistics were made with the program SPHINXS LEXICA V. 5.1., using the test: ANOVA *one-way* with a significance level of 5% to see if there was difference between the times of hospitalization in relation to surgical procedures and the comorbidities, respectively.

RESULTS

Among the 85 patients, 52 (61.17%) were male and 33 (38.82%) female. All underwent cardiac surgery, namely: coronary artery bypass grafting, 51 (60%), aortic valve replacement, 22 (25.88%), mitral valve replacement, seven (8.23%), and two patients underwent mitral valve replacement and aortic valve, repair of atrial septal defects, one (1.17%); correction of ventricular septal defect, one (1.17%), mitral valvuloplasty, one (1.17%), aneurysmectomy of left ventricle, one (1.17%), non-surgical angioplasty, one (1.17%) - (Figure 1).




Fig. 1 - Type of heart surgery performed. CABG: coronary artery bypass grafting, AVR: aortic valve replacement; MVR: mitral valve replacement, ASD: atrial septal defect, VSD: ventricular septal defect

In these patients, the mean age was 59.09 ± 14.40 years and the duration of ICU stay was 1-21 days with a mean of 4.16 ± 3.76 days. One (1.17%) patient remained in ICU for 1 day, 39 (45.88%) for 2 days, 12 (14.11%) for 3 days, 10 (11.76%) for four days, nine (10.58%) for 5 days and 14 (16.47%) for over 5 days (Figures 2 and 3).

The results show that the average length of stay in ICU were not significantly different ($F=0.41$; $1-P = 7.61\%$) between them in according to the surgical procedures performed ($P < 0.05$), which highlights the fact that these do not determine significant variations in the length of stay (Figure 4).




Figure 4 - Relation of length of stay in ICU (days) and intercurrent comorbidities in the immediate postoperative




Fig. 2 - Length of Stay in ICU in the postoperative per patient

The results of this study show that the average length of stay in ICU were significantly different ($F = 2,469.09$; $1-P \Rightarrow 99.99\%$) between them in relation to intercurrent comorbidities in the immediate postoperative period ($P < 0.05$), which highlights the fact that these determine significant variations in the length of stay.

With regard to comorbidities that kept the 14 patients admitted for longer we had: four patients with refractory hypertension (HBP) and pulmonary congestion, three remained for six days and one for seven days; a patient with perioperative acute myocardial infarction (AMI) stayed for seven days. The patient who stayed for eight days showed pulmonary congestion, hypertension and required MV for more than 24 hours. The patient who remained nine days had AMI, MRF and required MV for more than 24 hours. The patient who stayed 10 days had pulmonary congestion, hypertension and required MV for more than 24 hours. The patient who stayed for 12 days showed mediastinitis and sternal wound dehiscence. The patient who stayed 13 days had COPD, pulmonary congestion, hypertension and required MV for more than 24 hours.

The patient who remained hospitalized for 14 days was maintained on MV for 10 days due to respiratory failure and pneumonia after surgery and still mediastinitis with wound dehiscence; the patient who was hospitalized for 17 days, age 50, she presented severe COPD and went into acute respiratory failure (ARF) in the second postoperative day, having to remain on MV for more than 10 days, including being subjected to tracheostomy. The patient who was hospitalized for 18 days, age 72, showed decreased level of consciousness due to ischemic cerebral vascular accident (CVA) in the immediate postoperative and acute respiratory failure, staying in MV for 15 days, the patient who stayed 21 days hospital, aged 70 years, had peroperative instability, which required the use of intra-aortic balloon in the immediate postoperative period for more than 72 hours. He




Figure 3 - Relation between length of stay in ICU (days) and surgical procedures performed

also presented perioperative AMI, pulmonary congestion, need for mechanical ventilation throughout the hospital stay, had pneumonia associated with mechanical ventilation (MVAP), evolving to tracheostomy, renal failure, requiring hemodialysis and died.

Among the 85 patients, there were also two (2.35%) deaths of patients in the PO, who underwent emergency surgery with ventricular failure before surgery. The internment time ranged from 1 to 3 days. Bianco et al. [16] refer to increased morbidity in emergency surgeries.

DISCUSSION

Surgical interventions are part of the current therapy for heart disease [5,27]. The most prevalent type of surgery is CABG, in this study performed in 54 patients, confirming the published literature, followed by valve replacement [6,18,27,28]. The valve repair has become the procedure of choice for treatment of mitral valve disease. Many published studies show that valvuloplasty is preferable to valve replacement [28,29].

One of the factors that may increase morbidity and mortality in the PO, especially in the elderly is the duration of CPB [16,18,30,31]. The prolonged CPB, in addition to systemic inflammation, predisposes to postoperative complications, especially respiratory [3,4,7,16,31-36] and the need to use the IAB reflects hemodynamic instability [2,3,16,30, 31,37]. CABG surgery without CPB was initially performed in 1967 by Kolessov. While seemingly difficult to achieve, the arterial anastomosis with a beating heart may have greater benefits [3,4,32,36]. It was noticed, also, higher morbidity in patients undergoing emergency surgery in relation to elective surgery [7,16,27,37].

The complications of cardiac surgery may be related to pre-existing diseases such as previous pulmonary diseases, COPD, asthma, smoking [3,7,23,25], advanced age [7,16,18,23,30], poor nutritional status [38], obesity, diabetes [16,39], among others [25]. Among the most common complications we found atrial fibrillation and flutter, pulmonary congestion and stroke [7,15,22]. Jaeger et al. [24] showed that 14% of patients presented perioperative AMI. The factors influencing the onset of AMI include: reoperation, injury to the left coronary artery, female gender [29], preoperative unstable angina, a higher number of grafts and prolonged cardiopulmonary bypass time [16,24,31,32,37].

Thrombotic complications in CABG are presented as major causes of mortality. The perioperative myocardial infarction can cause not only enzyme elevations without clinical relevance, as well as low cardiac output, malignant tachyarrhythmias and death [4,36,37,40]. Other factors are diabetes mellitus and hypertension [9,24,25]. Mediastinitis is one of the most serious complications of median

sternotomy. Contamination of the wound may be the main determinant. Mediastinitis is defined as an infection of deep tissues of the wound associated with osteomyelitis of the sternum, which can also involve the retrosternal space. The factors most related to mediastinitis are obesity, COPD and diabetes mellitus [25,41]. The incidence of mediastinitis in sternotomy is 5% and death caused by mediastinitis is around 15% [25,41].

Other factors of increased incidence of postoperative complications are associated with the natural process of aging associated with increased prevalence of chronic degenerative diseases. Due to increased life expectancy, the number of people over 70 years who require cardiac surgery has increased [7,25,30]. Cardiac surgery in the elderly is associated with high morbidity and mortality due to the high prevalence of comorbidities and lower functional reserve of these patients [4,30,37]. Low body mass index (BMI) may increase the frequency of postoperative complications in the elderly undergoing coronary artery bypass grafting [2,30,38,42].

Metabolic syndrome (MS) is an association of cardiovascular risk factors that is closely linked with increased insulin resistance. The patients with MS, an average of 10% among women and 84% among men, are at increased risk by up to four times to death from coronary artery disease and three more times to death of any kind. As SM is defined as increased blood pressure (> 130x85 mmHg), fasting glycemia above 110 mg / dl, triglycerides > 150, HDL cholesterol <40 mg / dl and waist circumference > 102 cm in men and > 88 cm in women [26]. There was in this study, no patients with MS featured in evolution, but both obese and diabetic patients underwent surgery.

Obesity is directly related to the incidence of cardiovascular diseases [6]. Obesity provides a greater risk of decreased lung compliance [23], leads to the restriction on capacity and lung volumes that causes changes in gas exchange, increasing the chances of developing atelectasis and respiratory infections [6] while increasing the risk of mediastinitis [25].

According to Fernandes et al. [5], the length of hospital stay in cardiac surgery is on average around 6.6 days preoperatively with a mean of 5.4 ± 5.9 days in ICU. In this study, we chose to study those patients who stayed in ICU for more than five days. The pre-operative must be done well, with exams and guidance to patients, and this may require longer hospitalization [5]. The work of the respiratory therapy should begin preoperatively with the patient orientation on the work of breathing that should be done before and after surgery [3,6]. The patient that is well oriented in the preoperative treatment will better accept the treatment and the breathing exercises in physiotherapy and mechanical ventilation in the immediate postoperative period [6].

CONCLUSION

The complications that increase length of stay in ICU are mainly those related to respiratory function, COPD and smoking, pulmonary congestion, prolonged mechanical ventilation, infection, kidney failure, stroke and hemodynamic instability, such as hypertension, arrhythmias and acute myocardial infarction.

REFERENCES

1. Azzolin KO, Castro I, Feier F, Pandolfo F, Oderich C. Valor prognóstico do índice de performance miocárdica no pós-operatório de cirurgia de revascularização miocárdica. *Arq Bras Cardiol.* 2006;87(4):456-61.
2. Feier FH, Sant'Anna RT, Garcia E, Bacco F, Pereira E, Santos M, et al. Influências temporais nas características e fatores de risco de pacientes submetidos a revascularização miocárdica. *Arq Bras Cardiol.* 2006;87(4):439-45.
3. Lopes CR, Brandão CMA, Nozawa E, Auler Jr JOC. Benefícios da ventilação não-invasiva após extubação no pós-operatório de cirurgia cardíaca. *Rev Bras Cir Cardiovasc.* 2008;23(3):344-50.
4. Lobo Filho JG, Leitão MCA, Lobo Filho HG, Soares JPH, Magalhães GA, Leão Filho CSC, et al. Cirurgia de revascularização coronariana esquerda sem CEC e sem manuseio da aorta em pacientes acima de 75 anos. *Rev Bras Cir Cardiovasc.* 2002;17(3):208-14.
5. Fernandes AMS, Mansur AJ, Canêo LF, Lourenço DD, Piccioni MA, Franchi SM, et al. Redução do período de internação e de despesas no atendimento de portadores de cardiopatias congênitas submetidos à intervenção cirúrgica cardíaca no protocolo da via rápida. *Arq Bras Cardiol.* 2004;83(1):18-26.
6. Olmos SC, Graço BM, Oliveira SP, Figueiredo LC, Sasseron AB, Cardoso AL, et al. Tempo de internação hospitalar relacionado à fisioterapia respiratória no pré-operatório de cirurgia cardíaca eletiva. *Arq Med ABC.* 2007;32(Supl 2):S23-5.
7. Campagnucci VP, Silva AMRP, Pereira WL, Chamlian EG, Gandra SMA, Rivetti LA. EuroSCORE e os pacientes submetidos a revascularização do miocárdio na Santa Casa de São Paulo. *Rev Bras Cir Cardiovasc.* 2008;23(2):262-7.
8. Pierri H, Wajngarten M, Serro-Azul JB, Pivotto L, Nusbacher A, Gebara O, et al. Efeito da cirurgia de revascularização miocárdica sobre os episódios isquêmicos assintomáticos. *Arq Bras Cardiol.* 1998;70(4):265-9.
9. Liakopoulos OJ, Choi YH, Haldenwang PL, Strauch J, Wittwer T, Dörge H, et al. Impact of preoperative statin therapy on adverse postoperative outcomes in patients undergoing cardiac surgery: a meta-analysis of over 30,000 patients. *Eur Heart J.* 2008;29(12):1548-59.
10. Virani SS, Nambi V, Razavi M, Lee VV, Elayda M, Wilson JM, et al. Preoperative statin therapy is not associated with a decrease in the incidence of postoperative atrial fibrillation in patients undergoing cardiac surgery. *Am Heart J.* 2008;155(3):541-6.
11. Costa MB, Magalhães SS, Caldas LR, Barral MM, Souza MC, Paula RB. Valor do teste ergométrico na detecção da isquemia miocárdica silenciosa em pacientes diabéticos. *HU Revista.* 2008;34(2):107-12.
12. Dias CMCC, Maiato ACCA, Baqueiro KMM, Figueiredo AMF, Rosa FW, Pitanga JO, et al. Resposta circulatória à caminhada de 50 m na unidade coronariana na síndrome coronariana aguda. *Arq Bras Cardiol.* 2009;92(2):135-42.
13. Merin O, Ilan M, Oren A, Fink D, Deeb M, Bitran D, et al. Permanent pacemaker implantation following cardiac surgery: indications and long-term follow-up. *Pacing Clin Electrophysiol.* 2009;32(1):7-12.
14. Auler Júnior JO, Carmona MJ, Silva MH, Silva AM, do Amaral RV. Haemodynamic effects of pressure-controlled ventilation versus volume-controlled ventilation in patients submitted to cardiac surgery. *Clin Intensive Care.* 1995;6(3):100-6.
15. Alcade RV, Guaragna JC, Bodanese LC, Castro I, Sussenbach E, Noer R, et al. Alta dose de amiodarona em curto período reduz incidência de fibrilação e flutter atrial no pós-operatório de cirurgia de revascularização miocárdica. *Arq Bras Cardiol.* 2006;87(3):236-40.
16. Bianco ACM, Timerman A, Paes AT, Gun C, Ramos RF, Freire RBP, et al. Análise prospectiva de risco em pacientes submetidos à cirurgia de revascularização miocárdica. *Arq Bras Cardiol.* 2005;85(4):254-61.
17. Garwood S. Renal insufficiency after cardiac surgery. *Semin Cardiothorac Vasc Anesth.* 2004;8(3):227-41.
18. Silva AMRP, Campagnucci VP, Pereira WL, Rosa RF, Franken RA, Gandra SMA, et al. Revascularização do miocárdio sem circulação extracorpórea em idosos: análise da morbidade e mortalidade. *Rev Bras Cir Cardiovasc.* 2008;23(1):40-5.

19. Haase-Fielitz A, Haase M, Bellomo R, Lambert G, Matalanis G, Story D, et al. Decreased catecholamine degradation associates with shock and kidney injury after cardiac surgery. *J Am Soc Nephrol.* 2009;20(6):1393-403.
20. Chan AW. Treatment of severe functional mitral regurgitation: is cardiac surgery always indicated? *J Invasive Cardiol.* 2007;19(10):E286-9.
21. Kilger E, Weis F, Briegel J, Frey L, Goetz AE, Reuter D, et al. Stress doses of hydrocortisone reduce severe systemic inflammatory response syndrome and improve early outcome in a risk group of patients after cardiac surgery. *Crit Care Med.* 2003;31(4):1068-74.
22. Lúcio EA, Flores A, Blacher C, Leães PE, Lucchese FA, Ribeiro JP. Efetividade do metoprolol na prevenção de fibrilação e flutter atrial no pós-operatório de cirurgia de revascularização miocárdica. *Arq Bras Cardiol.* 2003;82(1):37-41.
23. Ambrozim ARP, Cataneo AJM. Aspectos da função pulmonar após revascularização do miocárdio relacionados com risco pré-operatório. *Rev Bras Cir Cardiovasc.* 2005;20(4):408-15.
24. Jaeger CP, Kalil RAK, Guaragna JCVC, Petracco JB. Fatores preditores de infarto do miocárdio no período perioperatório de cirurgia de revascularização miocárdica. *Rev Bras Cir Cardiovasc.* 2005;20(3):291-5.
25. Guaragna JC, Facchi LM, Baião CG, Cruz IBM, Bodanese LC, Albuquerque L, et al. Preditores de mediastinite em cirurgia cardíaca. *Rev Bras Cir Cardiovasc.* 2004;19(2):165-70.
26. Pimenta E, Passarelli Júnior O, Borelli F, Souza MG, Gun C, Amato V, et al. Síndrome metabólica em pacientes submetidos à cirurgia de revascularização miocárdica: prevalência e marcador de morbi-mortalidade no período intra-hospitalar e após 30 dias. *Arq Bras Cardiol.* 2007;88(4):413-7.
27. Araújo WF, Gerola LR, Kin HC, Pereira Filho A, Vargas GF, Catani R, et al. Tratamento cirúrgico das valvopatias aórticas com bioprótese de pericárdio bovino sem suporte: resultados imediatos. *Arq Bras Cardiol.* 2006;87(1):22-7.
28. Segalote RC, Pomerantzeff PMA, Brandão CMA, Stolf NAG. Cirurgia de preservação da valva aórtica em idosos com estenose aórtica. *Rev Bras Cir Cardiovasc.* 2008;23(4):519-23.
29. Balbinot AL, Kalil RAK, Prates PR, Sant'Anna JRM, Wender OC, Teixeira Filho GF, et al. Valvoplastia sem suporte para insuficiência mitral degenerativa: resultados a longo prazo. *Arq Bras Cardiol.* 2008;90(6):396-402.
30. Alves Junior L, Rodrigues AJ, Evora PRB, Basseto S, Scorzoni Filho A, Luciano PM, et al. Fatores de risco em septuagenários ou mais idosos submetidos a revascularização do miocárdio e ou operações valvares. *Rev Bras Cir Cardiovasc.* 2008;23(4):550-5.
31. Pontes JCDV, Silva GVR, Benafatti RA, Machado NP, Pontelli R, Pontes ERJC. Fatores de risco no desenvolvimento de insuficiência renal aguda após cirurgia de revascularização miocárdica com CEC. *Rev Bras Cir Cardiovasc.* 2007;22(4):484-90.
32. Lopes JB, Puig LB, Dallan LAO, Tossuniam CE, Gaiotto FA, Stolf NAG. Revascularização híbrida do miocárdio: uma alternativa para reduzir o tempo de perfusão. *Rev Bras Cir Cardiovasc.* 2008;23(2):240-4.
33. Halonen J, Halonen P, Järvinen O, Taskinen P, Auvinen T, Tarkka M, et al. Corticosteroids for the prevention of atrial fibrillation after cardiac surgery: a randomized controlled trial. *JAMA.* 2007;297(14):1562-7.
34. Fried LP, Kronmal RA, Newman AB, Bild DE, Mittelmark MB, Polak JF, et al. Risk factors for 5-years mortality in older adults: the Cardiovascular Health Study. *JAMA.* 1998;279(8):585-92.
35. Silva JP, Fonseca L, Baumgratz JF, Castro RM, Franchi SM, Sylos C, et al. Síndrome da hipoplasia do coração esquerdo: a influência da estratégia cirúrgica nos resultados. *Arq Bras Cardiol.* 2007;88(3):354-60.
36. Lobo Filho JG, Albuquerque JMAC, Gomes CBG, Siqueira RP, Landim RM, Oliveira RRM, et al. Revascularização cirúrgica das artérias posteriores do coração sem circulação extracorpórea. *Arq Bras Cardiol.* 1999;72(5):593-6.
37. Strabelli TMV, Stolf NAG, Uip DE. Uso profilático de um índice de risco de complicações após cirurgia cardíaca. *Arq Bras Cardiol.* 2008;91(5):342-7.
38. Reis C, Barbiero SM, Ribas L. O efeito do índice de massa corporal sobre as complicações no pós-operatório de cirurgia de revascularização do miocárdio em idosos. *Rev Bras Cir Cardiovasc.* 2008;23(4):524-9.
39. Amato VL, Timerman A, Paes AT, Baltar VT, Farsky PS, Farran JA, et al. Resultados imediatos da cirurgia de revascularização miocárdica: comparação entre homens e mulheres. *Arq Bras Cardiol.* 2004;83:14-20.
40. Oliveira MAB, Botelho PHH, Brandi AC, Santos CA, Soares MJF, Zaiantchick M, et al. Importância da troponina I no diagnóstico do infarto do miocárdio no pós-operatório de cirurgia de revascularização. *Rev Bras Cir Cardiovasc.* 2009;24(1):11-4.
41. Nina VJS, Assef MAS, Rodrigues RR, Mendes VGG, Lages JS, Amorim AMM, et al. Reconstrução da parede torácica com suporte metálico externo: técnica alternativa na mediastinite pós-esternotomia. *Rev Bras Cir Cardiovasc.* 2008;23(4):507-11.
42. Carrascal Y, Maroto L, Rey J, Arévalo A, Arroyo J, Echevarría JR, et al. Impact of preoperative anemia on cardiac surgery in octogenarians. *Interact Cardiovasc Thorac Surg.* 2009;10(2):249-55.