

ERRATUM / ERRATA

In the manuscript “Magnitude and variation of the burden of cancer mortality in Brazil and Federation Units, 1990 and 2015”, DOI: 10.1590/1980-5497201700050009, published in the Rev. bras. epidemiol. 2017; 20 (Suppl 1): 102-115.

Page 107, Where it reads:

Table 2. Standardized mortality rates/100,000 inhabitants (Brazil and FU) 2015 and percentage variation from 1990 to 2015 for five main types of cancers, men.

UF	MEN					
	All	Breast	Lung	Stomach	Colon/Rectum	Esophagus
BR	172.1 (-5.6)	33.4 (+12.2)	25.9 (-12.0)	19.5 (-37.3)	14.1 (+29.5)	10.5 (-14.1)
RO	149.9 (-13.3)	32.8 (+8.4)	21.9 (-7.5)	22.1 (-42.8)	8.1 (+15.2)	8.3 (-15.8)
AC	153.9 (+3.7)	32.5 (+25.8)	24.0 (+6.0)	23.0 (-22.2)	8.8 (+33.5)	5.6 (-0.3)
AM	184.5 (+1.0)	33.9 (+17.7)	33.5 (-8.1)	31.6 (-18.1)	9.7 (+31.7)	6.2 (-1.1)
RR	157.6 (-5.5)	37.4 (+14.0)	23.2 (-12.3)	23.5 (-36.0)	7.4 (+35.1)	5.7 (-8.0)
PA	148.3 (+0.2)	31.4 (+26.5)	23.2 (-8.5)	26.3 (-26.7)	8.7 (+32.9)	4.9 (-4.4)
AP	175.4 (+10.6)	34.0 (+36.6)	28.6 (+8.6)	38.6 (-17.1)	7.7 (+61.9)	5.8 (+19.5)
TO	141.1 (+17.9)	38.5 (+41.8)	18.2 (+20.5)	13.5 (-23.4)	9.0 (+63.1)	5.7 (+15.0)
MA	136.3 (-16.2)	32.8 (-0.1)	18.6 (-18.9)	18.0 (-42.8)	8.0 (-0.7)	4.1 (-18.0)
PI	138.2 (+16.8)	33.3 (+26.0)	19.6 (+18.1)	13.0 (-13.0)	8.8 (+51.0)	6.4 (+22.7)
CE	185.7 (+34.6)	36.8 (+41.5)	26.4 (+55.2)	28.6 (-1.4)	11.1 (+84.9)	11.6 (+55.6)
RN	164.5 (+28.0)	34.9 (+40.9)	21.1 (+36.1)	22.8 (-2.5)	10.2 (+53.4)	8.7 (+40.8)
PB	162.1 (+40.9)	35.0 (+45.0)	19.5 (+54.1)	21.3 (+14.0)	9.5 (+66.5)	9.4 (+47.3)
PE	152.1 (+17.1)	34.8 (+26.8)	20.8 (+16.7)	15.8 (-4.5)	8.8 (+48.3)	7.5 (+21.8)
AL	125.6 (+6.9)	29.3 (+22.4)	15.8 (-3.2)	13.6 (-18.5)	8.2 (+40.0)	5.9 (+7.6)
SE	152.0 (+15.0)	38.4 (+38.5)	21.1 (+6.0)	14.5 (-12.2)	10.0 (+47.1)	6.1 (+11.8)
BA	162.6 (+19.7)	38.7 (+40.9)	19.1 (+12.6)	18.7 (-13.8)	11.2 (+44.2)	10.0 (+27.3)
MG	167.0 (-8.0)	32.3 (+9.8)	21.8 (-8.7)	20.4 (-43.9)	12.6 (+31.8)	13.4 (-14.5)
ES	168.1 (-8.5)	31.9 (+14.0)	23.9 (-11.6)	20.5 (-46.5)	11.6 (+27.3)	14.4 (-9.1)
RJ	176.0 (-16.6)	34.2 (+2.9)	28.5 (-31.2)	17.8 (-44.2)	17.5 (+17.0)	9.0 (-23.3)
SP	173.4 (-17.4)	31.1 (-0.9)	25.1 (-24.5)	18.6 (-50.0)	17.9 (+22.5)	9.8 (-27.1)
PR	190.2 (-7.4)	34.0 (+14.6)	27.2 (-4.9)	21.7 (-44.7)	16.4 (+30.1)	13.9 (-25.5)
SC	195.0 (-13.2)	31.0 (+2.8)	36.7 (-10.8)	21.4 (-43.7)	14.0 (+13.7)	14.2 (-30.6)
RS	216.1 (-18.2)	34.4 (-5.1)	46.2 (-26.8)	17.2 (-44.7)	18.3 (+6.2)	16.7 (-32.0)
MS	167.2 (+3.3)	34.8 (+19.5)	24.8 (+3.0)	18.7 (-36.3)	13.7 (+50.1)	10.6 (+3.1)
MT	164.9 (-1.4)	36.9 (+16.4)	25.1 (+3.4)	19.5 (-42.6)	11.3 (+33.0)	9.6 (+0.5)
GO	155.6 (-4.9)	33.3 (+9.3)	24.0 (-3.9)	15.7 (-40.5)	12.0 (+25.0)	8.1 (-12.7)
DF	156.6 (-12.2)	33.2 (+0.6)	23.4 (-16.1)	14.9 (-43.5)	14.0 (+25.9)	7.5 (-18.4)

Note: percentage variation 1990 and 2015 in parenthesis; BR: Brazil; FU in acronyms.

It should read:

Table 2. Standardized mortality rates/100,000 inhabitants (Brazil and FU) 2015 and percentage variation from 1990 to 2015 for five main types of cancers, men.

UF	MEN					
	All	Prostate	Lung	Stomach	Colon/Rectum	Esophagus
BR	172.1 (-5.6)	33.4 (+12.2)	25.9 (-12.0)	19.5 (-37.3)	14.1 (+29.5)	10.5 (-14.1)
RO	149.9 (-13.3)	32.8 (+8.4)	21.9 (-7.5)	22.1 (-42.8)	8.1 (+15.2)	8.3 (-15.8)
AC	153.9 (+3.7)	32.5 (+25.8)	24.0 (+6.0)	23.0 (-22.2)	8.8 (+33.5)	5.6 (-0.3)
AM	184.5 (+1.0)	33.9 (+17.7)	33.5 (-8.1)	31.6 (-18.1)	9.7 (+31.7)	6.2 (-1.1)
RR	157.6 (-5.5)	37.4 (+14.0)	23.2 (-12.3)	23.5 (-36.0)	7.4 (+35.1)	5.7 (-8.0)
PA	148.3 (+0.2)	31.4 (+26.5)	23.2 (-8.5)	26.3 (-26.7)	8.7 (+32.9)	4.9 (-4.4)
AP	175.4 (+10.6)	34.0 (+36.6)	28.6 (+8.6)	38.6 (-17.1)	7.7 (+61.9)	5.8 (+19.5)
TO	141.1 (+17.9)	38.5 (+41.8)	18.2 (+20.5)	13.5 (-23.4)	9.0 (+63.1)	5.7 (+15.0)
MA	136.3 (-16.2)	32.8 (-0.1)	18.6 (-18.9)	18.0 (-42.8)	8.0 (-0.7)	4.1 (-18.0)
PI	138.2 (+16.8)	33.3 (+26.0)	19.6 (+18.1)	13.0 (-13.0)	8.8 (+51.0)	6.4 (+22.7)
CE	185.7 (+34.6)	36.8 (+41.5)	26.4 (+55.2)	28.6 (-1.4)	11.1 (+84.9)	11.6 (+55.6)
RN	164.5 (+28.0)	34.9 (+40.9)	21.1 (+36.1)	22.8 (-2.5)	10.2 (+53.4)	8.7 (+40.8)
PB	162.1 (+40.9)	35.0 (+45.0)	19.5 (+54.1)	21.3 (+14.0)	9.5 (+66.5)	9.4 (+47.3)
PE	152.1 (+17.1)	34.8 (+26.8)	20.8 (+16.7)	15.8 (-4.5)	8.8 (+48.3)	7.5 (+21.8)
AL	125.6 (+6.9)	29.3 (+22.4)	15.8 (-3.2)	13.6 (-18.5)	8.2 (+40.0)	5.9 (+7.6)
SE	152.0 (+15.0)	38.4 (+38.5)	21.1 (+6.0)	14.5 (-12.2)	10.0 (+47.1)	6.1 (+11.8)
BA	162.6 (+19.7)	38.7 (+40.9)	19.1 (+12.6)	18.7 (-13.8)	11.2 (+44.2)	10.0 (+27.3)
MG	167.0 (-8.0)	32.3 (+9.8)	21.8 (-8.7)	20.4 (-43.9)	12.6 (+31.8)	13.4 (-14.5)
ES	168.1 (-8.5)	31.9 (+14.0)	23.9 (-11.6)	20.5 (-46.5)	11.6 (+27.3)	14.4 (-9.1)
RJ	176.0 (-16.6)	34.2 (+2.9)	28.5 (-31.2)	17.8 (-44.2)	17.5 (+17.0)	9.0 (-23.3)
SP	173.4 (-17.4)	31.1 (-0.9)	25.1 (-24.5)	18.6 (-50.0)	17.9 (+22.5)	9.8 (-27.1)
PR	190.2 (-7.4)	34.0 (+14.6)	27.2 (-4.9)	21.7 (-44.7)	16.4 (+30.1)	13.9 (-25.5)
SC	195.0 (-13.2)	31.0 (+2.8)	36.7 (-10.8)	21.4 (-43.7)	14.0 (+13.7)	14.2 (-30.6)
RS	216.1 (-18.2)	34.4 (-5.1)	46.2 (-26.8)	17.2 (-44.7)	18.3 (+6.2)	16.7 (-32.0)
MS	167.2 (+3.3)	34.8 (+19.5)	24.8 (+3.0)	18.7 (-36.3)	13.7 (+50.1)	10.6 (+3.1)
MT	164.9 (-1.4)	36.9 (+16.4)	25.1 (+3.4)	19.5 (-42.6)	11.3 (+33.0)	9.6 (+0.5)
GO	155.6 (-4.9)	33.3 (+9.3)	24.0 (-3.9)	15.7 (-40.5)	12.0 (+25.0)	8.1 (-12.7)
DF	156.6 (-12.2)	33.2 (+0.6)	23.4 (-16.1)	14.9 (-43.5)	14.0 (+25.9)	7.5 (-18.4)

Note: percentage variation 1990 and 2015 in parenthesis; BR: Brazil; FU in acronyms.

Page 110, Where it reads:




Figure 2. Standardized rates of cancer mortality in women: (A) overall; (B) prostate; and (C) lung. States of Brazil, 1990–2015.

It should read:




Figure 2. Standardized rates of cancer mortality in men: (A) overall; (B) prostate; and (C) lung. States of Brazil, 1990–2015.