



Original Article

Plants from the Brazilian Traditional Medicine: species from the books of the Polish physician Piotr Czerniewicz (Pedro Luiz Napoleão Chernoviz, 1812–1881)



Letícia M. Ricardo^{a,c}, Juliana de Paula-Souza^d, Aretha Andrade^{b,c}, Maria G.L. Brandão^{b,c,*}

^a Ministério da Saúde, Departamento de Assistência Farmacêutica e Insumos Estratégicos, Esplanada dos Ministérios, Brasília, DF, Brazil

^b Centro Especializado em Plantas Aromáticas, Medicinais e Tóxicas, Museu de História Natural e Jardim Botânico, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

^c Faculdade de Farmácia, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

^d Universidade Federal de São João del Rei, Campus Sete Lagoas, Sete Lagoas, MG, Brazil

ARTICLE INFO

Article history:

Received 1 October 2016

Accepted 10 January 2017

Available online 2 March 2017

Keywords:

Historical records

Useful plants

Medicinal plants

19 century

ABSTRACT

The Brazilian flora is very rich in medicinal plants, and much information about the traditional use of the Brazilian plants is only available from early literature and we are facing a rapid process of loss of biodiversity. To retrieve data about useful plants registered in the books of the Polish physician P.L.N. Chernoviz, who lived in Brazil for 15 years in the 19th century. The aim is to improve our knowledge about Brazilian plants, and to ensure the benefits of sharing it with potential users. Data about Brazilian plants were obtained from six editions of the book *Formulary and Medical Guide* (Formulário e Guia Médico), published in 1864, 1874, 1888, 1892, 1897 and 1920. All this information was then organized in boxes, and correlated with pharmacological studies from PubMed and Lilacs. A total of 238 species were recorded and 117 different traditional uses were registered for the plants. The most common uses were to treat general skin diseases and intermittent fevers, as purgative, diuretics and antidiarrheal. From the total, only 61 species (25.6%) have been to date subject to pharmacological studies and had their activities confirmed. Chernoviz books represent an important source of data about plants used in traditional medicine of Brazil. Their records were collected at a time when Brazil's native vegetation was still largely intact, and traditional medicine was practiced based on native plants. We argue that these plants must be prioritized in development of medicines, submitting them to clinical studies or by considering their traditionality, as established by WHO. Strategies for the protection of the traditional knowledge are also necessary.

© 2017 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Since 2002, the World Health Organization (WHO) has been stimulating the rescue of data from plants used in ancient medical practice, since they are considered potentially useful in the development of new drugs (WHO, 2002). The American plants were widely used long before the arrival of the Europeans in the continent in the 15th century, and besides being one of the richest countries in biodiversity, Brazil is also one of the most diverse in terms of Amerindian culture (Neves, 2006; Forzza et al., 2012). However, both plants and traditional knowledge are under heavy threat due to the continued destruction of the native ecosystems,

as a consequence of the successive economic cycles along the five centuries of increasing human occupancy. In fact, today only 7% of the Atlantic Forest survives, and the Cerrado and Caatinga are also in a rapid process of destruction (Giulietti et al., 2005). This situation highlights the necessity to promote the appropriate use and conservation of useful species from the Brazilian biodiversity, as suggested by the WHO.

Historical research can play an important role in recovering valuable ethnopharmacological data regarding the use of plants (Heinrich et al., 2006; Medeiros, 2008). Over the last ten years, our research group has concentrated efforts to recover informations about useful Brazilian plants in books and other documents from the past centuries. Most registers about Brazilian plants have been recovered from works of Europeans that travelled or lived in the country in the 19th century. Hundreds of species of medicinal plants used in Brazil were registered by these scientists (Brandão

* Corresponding author.

E-mail: mbrandao@ufmg.br (M.G. Brandão).

et al., 2008; Breitbach et al., 2013; Fagg et al., 2015). Among the Europeans that lived in Brazil in this period is the Polish physicist Piotr Czerniewicz (1812–1881) or Pedro Luiz Napoleão Chernoviz, as he was known in Brazil.

Chernoviz was born in Lukov, but was forced to leave the country while still a medical student at the University of Warsaw in 1831, due to his involvement in an uprising against the Russians. In Montpellier, he continued his studies and in 1837 he was awarded by the French government with the medal of merit, due to his courageous participation in a cholera epidemic (Guimarães, 2005). Chernoviz arrived in Rio de Janeiro in the early 1840s to work as a medical doctor. He returned to France in 1855 and died in 1881.

A year after his arrival in Brazil, Chernoviz wrote the "Formulary and Medical Guide" (*Formulário e Guia Médico*) directed to physicians, a book that achieved 19 editions during his life, being published by his son afterwards. In 1842 he published the "Dictionary of Popular Medicine" (*Dicionário de Medicina Popular*), which had a more accessible language to be used by the population. Both books were a striking success in Brazil, especially the "Dictionary" that was widely used in the inner parts of the country, where professional medical assistance was virtually nonexistent. Both books present a good description about the medical techniques at the time but more importantly, a rich register of the traditional uses of Brazilian plants. The aim of this study was to retrieve data about useful Brazilian plants in six editions of the book Formulary and Medical Guide, and discuss about the possibility of their current use.

Methods

Survey of Chernoviz books

Data on the use of native plants were carefully collected from the following editions of the book *Formulary and Medical Guide*: 6th edition (1864), 9th edition (1874), 13th (1888), 15th (1892), 16th (1897) and 19th (1920). The editions of 1864, 1874 and 1892 were studied in the Libraries of Medicine and Pharmacy School of University of Paris. The edition of 1888 was consulted in the Library of the Museum of Natural History, also in Paris. The others (1897 and 1920) were consulted in the Natural History Museum and Botanical Gardens and the Pharmacy Faculty of the Federal University of Minas Gerais, in Belo Horizonte, Brazil.

Only the species referred as "plant from Brazil" by the author were considered. Box 1 shows the species cited by Chernoviz that were submitted to pharmacological studies correlated with the traditional uses. In Box 2 are listed the other species. Different popular names were attributed by the author for the plants in the subsequent editions, but in this study we have considered only those names from the last edition, published in 1920, considering the most complete. Only the current scientific names, obtained by consulting the website www.floradobrasil.jbrj.gov.br, from the Rio de Janeiro Botanical Garden, were included in the Boxes.

The confirmation of the geographic origin of the plants was also observed in this same website. Some discrepancies about the occurrence of the plants cited by Chernoviz and our current knowledge of the Brazilian flora were observed. For instance, *Cestrum nocturnum* and *Coccus nucifera* are currently not considered natives in the country's flora however common these plants might be. In other cases, this is just a matter of misapplied names – *Drymis winteri* does not occur in Brazil, the correct identification of the species found in this country being *Drimys granatensis* instead. Moreover, *Chiococca* spp. and *Senebiera pinnatifida*, cited by the author as exotic plants, are currently known to be natives.

Many species, that only the popular name were cited by the author, were not included in the Boxes as "amapá",

"muirapuama", "pajurá", "cabeça-de-negro", "cabeça-de-moleque" and "tamaquaré". Other scientific names cited by the author were also excluded from the Boxes since they were not confirmed as valid names, namely "cipó-guyra" (*Bignonia guyra*), "rabo-de-tatu" or "sumaré" (*Cypripedium brasiliensis*), "caámembeça" (*Polygala paraensis*), "cabacinho" or "buchá-dos-caçadores" (*Momordica bucha*) and "jaborandi" (*Monnieria trifolia*). "Chique-chique" was also excluded from this survey, since the scientific name assigned to it by Chernoviz (*Cereus jamacaru*) does not match the species to which this popular name has always been assigned (*Pilosocereus gounellei*), which could lead to ambiguous interpretations. Likewise, the species assigned to "caferana" or "jacaré-açu" (*Tachia guianensis* Aubl.) does not occur in Brazil, and although this is probably a case of misidentification, this plant was excluded.

Survey on other historical references about Brazilian plants

We performed searches for data about the plants cited by Chernoviz in other historical bibliographical sources. The oldest one was *Triaga Brasilica*, a recipe from the 16th century (Pereira et al., 1996). The book *Historiae Naturalis & Medicae* published by Piso in 1648 was also studied, but since only the popular names of the plants were cited. Data registered in the 1840s manuscripts of George Gardner (Fagg et al., 2015), the field books (Brandão et al., 2012) and the book *Plantes usuelles des Brasiiliens* by Auguste de Saint-Hilaire (Saint-Hilaire, 1824), the book *Materia Medica* by the german Karl von Martius (Martius, 1843) and the 1st Edition of Brazilian official *Pharmacopoeia* were also obtained (Brandão et al., 2009).

Survey of laboratory studies correlated with traditional uses

Searches in PubMed and Lilacs for information about pharmacological studies correlated with the traditional use were also performed, and the results are listed in Box 1.

Results and discussion

In the present study, we have recovered data about Brazilian useful plants from six editions of the *Formulary and Medical Guide*, published by Chernoviz (followed by his son) from 1864 to 1920. This book reached 19 different editions, being the first published already in 1841, just one year after the author arrived in Brazil. The number of Brazilian plants was dramatically increased along the new editions. As a physician, Chernoviz wrote detailed information about the efficacy and preparation of the remedies from each plant. As he describes in the introduction of each edition, the data were collected from literature available at that time and also from his medical experience during the 15 years he lived in Rio de Janeiro.

Data from 238 useful plants could be retrieved from the books (Boxes 1 and 2). The species are distributed among 83 families, Asteraceae and Fabaceae being the most represented with nineteen (8%) and seventeen (7%), respectively. The other most represented families were Rubiaceae twelve (5%), Euphorbiaceae, Lauraceae and Malvaceae (ten species, 4%), Apocynaceae (nine species, 4%), Solanaceae (eight species, 3%), Cucurbitaceae (seven species, 3%), Bignoniaceae and Rutaceae (six species, 3%), Meliaceae and Verbenaceae (five species), Araceae, Lamiaceae, Lecythidaceae and Myrtaceae, (four species) and Boraginaceae, Gentianaceae, Hypericaceae, Loganiaceae, Menispermaceae, Moraceae, Phyllantaceae, Sapindaceae and Sapotaceae (three species). The remaining families are represented by one or two species.

A total of 117 different traditional uses were recorded for the plants in the six studied editions, and the two most common were as purgative and to treat intermittent fevers (30 and 26 species, respectively). Other well documented uses were as diuretic (21), antidiarrheal and tonic (18), stimulant (17), as astringent, febrifuge

Box 1: Plants from the Formulary and Medical Guide, their uses, first edition of publication and recent studies correlated with the traditional uses

Family/scientific name	Popular name	Part/indication	Edition	Recent studies correlated with use	
Amaranthaceae <i>Chenopodium ambrosioides</i> L. ^{a,f}	herba-de-santa-maria	Vermifuge	1864	Amoebicidal (Avila-Blanco et al., 2014), acaricidal (Kouam et al., 2015), antischistosomal (Kamel et al., 2011)	
Anacardiaceae <i>Anacardium occidentale</i> L. ^{a,e,f}	cajueiro, acaju	Lupus, acne, eczema, ulcers, psoriasis, leprosy	1874	Anti-inflammatory, wound healing (Olajide et al., 2004; Pawar et al., 2004)	
	<i>Schinus terebinthifolius</i> Raddi ^{a,c,d,e,f}	aroeira, corneiba	Leaves to treat ulcers, bark astringent, to treat leg oedemas	1864	Wound healing (Estevão et al., 2013), gastric ulcer (Santos et al., 2013)
Apocynaceae <i>Geissospermum vellosii</i> Allemão	camará-de-bilro, camará-do-mato, pao-pereira, forquilha, pau-de-pente	Bark to treat intermittent fevers and tonic	1864	Antiplasmodial (Mbeunkui et al., 2012)	
Aquifoliaceae <i>Ilex paraguariensis</i> A. St.-Hil. ^{c,d,e,f}	mate, congonha	Stimulant, sudorific, digestive ^a	1874	Stimulant (Santos et al., 2015)	
Aristolochiaceae <i>Aristolochia trilobata</i> L. ^b	milhomens, jarrinha	Lack of appetite, fevers, skin ulcers	1864	Topical antiinflammatory (Sosa et al., 2002); antibacterial (Camporese et al., 2003)	
Asteraceae <i>Acanthospermum australe</i> (Loefl.) Kuntze ^e	picão-da-praia	Tonic, diuretic, intermittent fevers	1864	Antimalarial (Carvalho et al., 1991; Carvalho and Krettli, 1991)	
<i>Ageratum conyzoides</i> L. ^{d,e}	herba-de-são-joão, mentrasto	Stimulant, antidiarrheal, baths	1864	Antimicrobial (Singh et al., 2013; Adetutu et al., 2012)	
<i>Ayapana triplinervis</i> (Vahl) R.M.King & H.Rob. ^{e,f}	japana, ayapana	Sudorific, wound healing	1864	Antibacterial (Gupta et al., 2002)	
<i>Baccharis genistelloides</i> (Lam.) Pers.; <i>Baccharis crispa</i> Spreng. ^{c,e,f}	carqueja-amargosa	Tonic, febrifuge	1864	Anti-inflammatory (Nogueira et al., 2011; Paul et al., 2009)	
<i>Bidens pilosa</i> L. ^e	picão, cuambú, guambú	Jaundice, ulcers	1874	Hepatoprotective (Kwiecinski et al., 2011; Suzigan et al., 2009)	
<i>Chaptalia nutans</i> (L.) Pol.	língua-de-vaca	Tonic, bronchitis, skin ulcers	1874	Antibacterial (Truiti et al., 2003); Antiinflammatory (Badilla et al., 1999)	
<i>P. sagittalis</i> (Lam.) Cabrera ^a	quitoco, caculage	Excitant, digestive	1864	Gastroprotective (Figueiredo et al., 2011)	
<i>Sonchus oleraceus</i> L.	serralhinha	Aperient, diuretic, jaundice, nephritis	1874	Anti-inflammatory (Vilela et al., 2010)	
Bignoniacae <i>Zeyheria montana</i> Mart.	mandioquinha-do-campo, bolsa-de-pastor	Roots skin disease	1874	Antiproliferative (Seito et al., 2011), antiinflammatory, antinociceptive (Guenka et al., 2008)	
Bixaceae <i>Bixa orellana</i> L. ^{a,b,e}	urucu, orucu	Expectorant, bronchitis	1864	Antimicrobial (Fleischer et al., 2003; Castello et al., 2002), antihistamine (Yong et al., 2013)	
Boraginaceae <i>Heliotropium indicum</i> L. ^e	aguará-ciunhá-açú, jacuá-acanga	Skin ulcers and burns	1874	Wound healing (Dash and Murthy, 2011; Reddy et al., 2002)	
Convolvulaceae <i>Operculina</i> sp. ^{e,f}	batata-de-purga, ipú, jalapa, purga-de-amaro-leite, de cayapó, gentio, de joão-paes	Roots and seeds purgative	1874	Laxative (Paganotte et al., 2016; Michelin and Salgado, 2004)	
Crassulaceae <i>Kalanchoe crenata</i> (Andrews) Haw. ^c	sayão	Leaves as wound healing	1864	Antiinflammatory (Costa et al., 2006; Ibrahim et al., 2002)	
Dilleniaceae <i>Curatella americana</i> L. ^{c,d,e}	sambaiba	Barks astringent, skin ulcers	1874	Anti-ulcerogenic (Hiruma-Lima et al., 2009)	
Euphorbiaceae <i>Joannesia princeps</i> Vell. ^{a,e}	anda-açu, purga-de-gentio, paulistas, coco-de-purga, fruta-de-arara	Purgative	1864	Episodes of diarrhoea (Nishioka and Escalante, 1997)	
<i>Croton antisiphiliticus</i> Mart. ^{c,e}	herba-mular, alcaphoreira, curraleira, pé-de-perdiz	Leaves sudorific, skin ulcers, stimulant	1874	Antiinflammatory (Reis et al., 2014)	
Fabaceae <i>Bowdichia virgilioides</i> Kunth ^{e,f}	sebipira, sucopira	Bark tonic and diaphoretic, to treat skin diseases	1874	Sport resistance (Santos et al., 2014), skin disease (Agra et al., 2013); antiinflammatory (Barros et al., 2010; Thomazzi et al., 2010)	
<i>Copaifera</i> spp. ^{a,d,e,f}	copahiba	Astringent, blennorrhagy and other venereal diseases	1864	Wound healing (Paiva et al., 2002); Antiinflammatory (Vargas et al., 2015; Gelmini et al., 2013); Antimicrobial (Souza et al., 2011; Santos et al., 2008)	

<i>Dahlstedtia pinnata</i> (Benth.) Malme	timbó boticário	Narcotic, liver disorders	1888	Sedative effect (Della Loggia et al., 1981)
<i>Erythrina</i> spp. ^f	mulungú, crista-de-galo	Bark sedative, seeds are poison	1874	Anxiolytic (Santos Rosa et al., 2012; Raupp et al., 2008)
<i>Mimosa tenuiflora</i> (Willd.) Poir. ^e	jurema	Narcotic, skin ulcers	1874	Wound healing (Zippel et al., 2009; Rivera-Arce et al., 2007)
<i>Parapiptadenia rigida</i> (Benth.) Brenan ^e	angico	Bronchitis, liver disorders	1874	Antimicrobial, analgesic, anti-inflammatory (Araujo et al., 2014)
<i>Stryphnodendron adstringens</i> (Mart.) Coville ^{c,e,f}	barbatimão	Barks astringent, venereal disease, wound healing	1874	Wound healing (Pinto et al., 2015; Coelho et al., 2010), candidiasis (Morey et al., 2015; Luiz et al., 2015; Ishida et al., 2006)
Hypericaceae				
<i>Hypericum brasiliense</i> Choisy ^{b,f}	alecrim-bravo	Snake bites, aromatic baths	1920	Snake bites (Dal Belo et al., 2013)
<i>Hypericum connatum</i> Lam. ^e	orelha-de-gato	Astringent, throat inflammation	1874	Oral lesions (Fritz et al., 2007), antibacterial (Fratianni et al., 2013)
Lauraceae				
<i>Licaria puchury-major</i> (Mart.) Kosterm. ^e	puchury, pichurim	Seeds stimulant, tonic, dyspepsy, leucorrhea	1874	Psychopharmacological (Carlini et al., 1983)
<i>Ocotea odorifera</i> (Vell.) Rohwer ^e	pereiorá, casca-preciosa	Nervous system, leucorrhea, oedema, antidiarrheal	1874	Antifungal (Yamaguchi et al., 2011)
Lecythidaceae				
<i>Bertholletia excelsa</i> Bonpl. ^e	castanha-do-maranhão, do pará	Barks jaundice, intermittent fever, renal calculi	1892	Antiplasmodial (Oliveira et al., 2015)
Loganiaceae				
<i>Spigelia laurina</i> Cham. & Schltl. ^e	espigelia, arapabaca	Vermifuge	1892	Anthelmintic (Ademola et al., 2007; Jegede et al., 2006)
<i>Strychnos pseudoquina</i> A.St.-Hil. ^{c,d,e,f}	quina-do-campo	Bark intermittent fevers	1874	Antimalarial (Andrade-Neto et al., 2003)
Loranthaceae				
<i>Struthanthus marginatus</i> (Desr.) Blume ^f	herba-de-passarinho, enxerto-de-passarinho	Leaves to treat lung diseases	1888	Antimicobacterial (Leitão et al., 2013)
Malvaceae				
<i>Gossypium</i> spp. ^{e,f}	algodoeiro	Wounds, erysipelas	1864	Wound healing (Annan and Houghton, 2008)
Meliaceae				
<i>Cabralea canjerana</i> (Vell.) Mart. ^{e,f}	canjerana	Oil prevent insect bites, bark purgative and vomitive	1874	Insect repellent (Magrini et al., 2014)
<i>Carapa guanensis</i> Aubl. ^e	andiroba, carapa	Bark febrifuge and vermifuge, skin ulcers, insect repellent	1874	Antiplasmodial (Pereira et al., 2014; Miranda Júnior et al., 2012); vermicudge (Sarria et al., 2011); wound healing (Nayak et al., 2010, 2011); antimicrobial (Meccia et al., 2013); antiinflammatory (Henriques and Penido, 2014); insect repellent (Miot et al., 2004).
<i>Guarea guidonia</i> (L.) Sleumer ^e	marinheiro, gitó	Bark purgative and vermicudge	1874	Antiprotozoal (Weniger et al., 2001)
<i>Trichilia</i> sp. ^e	marinheiro-de-folha-miúda	Roots to treat intermittent fevers	1874	Antimalarial (Castro et al., 1996)
Menispermaceae				
<i>Cissampelos ovalifolia</i> DC. ^{b,c,d,e}	orelha-de-onça	Roots to treat snake bites and intermittent fevers	1874	Antimalarial (Fischer et al., 2004)
Myrtaceae				
<i>Psidium pomiferum</i> L., <i>P. guajava</i> L. ^{a,e,f}	goiabeira	Leaves and barks antidiarrheal, leg oedema	1874	Antidiarrheal (Birdi et al., 2010, 2014; Ojewole et al., 2008); anti-inflammatory (Araujo et al., 2014; Jang et al., 2014)
Passiflorace				
<i>Passiflora maliformis</i> L. ^{c,e}	maracujá	Convalescences	1920	Nutritive (Devi Ramaiya et al., 2013)
Phyllanthaceae				
<i>Phyllanthus microphyllus</i> Kunth, <i>P. niruri</i> L. ^e	herba-pombinha	Diuretic	1864	Kidney troubles (Giribabu et al., 2014; Micali et al., 2006)
Phytolaccaceae				
<i>Petiveria alliacea</i> L., <i>Seguieria floribunda</i> Benth. ^e	pau-de-alho, cipó, guararema, ybirarema.	Baths for rheumatic pain, hemorrhoid and dropsy	1874	Antinociceptive (Gomes et al., 2005), anti-inflammatory (Lopes-Martins et al., 2002)
Piperaceae				
<i>Piper mikianum</i> (Kunth) Steud., <i>P. peletatum</i> L., <i>P. umbellatum</i> L. ^{a,b,c,e}	periparoba, aguaxima, caapeba	Leaves rheumatism ^a and to treat ulcers ^a ; roots as stomachic, sudorific, intermittent fevers	1864	Anti-inflammatory (Iwamoto et al., 2015; Roersch, 2010). Antimalarial (Rocha e Silva et al., 2015)
Plantaginaceae				
<i>Scoparia dulcis</i> L. ^c	vassorinha, tupeiçava	Bronchitis	1874	Antiinflammatory (Tsai et al., 2011)
Rubiaceae				
<i>Carapichea ipecacuanha</i> (Brot.) L.Andersson ^{a,b,c,d,e}	ipecacuanha, poaya, ipéca	Roots vomitive ^a , tonic, expectorant, antidiarrheal	1864	Several studies
<i>Palicourea rigida</i> Kunth ^{c,d,e}	herba-de-rato, don-bernardo, douradinha-de-campo, gritadeira	Poison	1874	Toxic (Cook et al., 2014)

<i>Remijia ferruginea</i> (A.St.-Hil.) DC. ^{c,e,f}	quina-da-serra, remijio, do campo	Barks bitter, to treat intermittent fevers	1864	Antimalarial (Andrade-Neto et al., 2003)
Rutaceae <i>Zanthoxylum fagara</i> (L.) Sarg. ^{c,e}	coentrilho	Ear ache	1920	Antinociceptive, antiinflammatory (Villalba et al., 2007)
Sapindaceae <i>Paulinia cupana</i> Kunth ^{e,f}	guaraná	Tonic, antidiarrheal	1864	Improves fatigue (Oliveira Campos et al., 2011)
Smilacaceae <i>Smilax salsaparrilha</i> L., <i>S. syphilitica</i> Humb. & Bonpl. ex Willd. ^{e,f}	salsaparrilha, japecanga, salsaparrilha-do-brasil	Antisyphilitic, skin disease, rheumatism and gout	1864	Several studies with <i>Smilax</i> spp.
Simaroubaceae <i>Simaba ferruginea</i> A.St.-Hil. ^e	calunga	Dyspepsy, intermittent fevers	1874	Antiulcer (Souza-Almeida et al., 2011; Noldin et al., 2005)
<i>Simarouba amara</i> Aubl. ^{e,f}	simaruba	Bitter, febrifuge, antidiarrheal	1864	Hepatoprotective (Maranhão et al., 2014); malaria (Franssen et al., 1997); gastrointestinal disorders (Caceres et al., 1990)
Solanaceae <i>Cestrum parqui</i> L'Hér. ^e	coerana	Leaves stimulant, carminative	1874	Psychosis (Siskind et al., 2014)
<i>Physalis angulata</i> L. ^{c,e}	camapú, juápoca	Ear ache, tonic, liver disorders	1874	Antinociceptive (Bastos et al., 2006; Choi and Hwang, 2003)
<i>Solanum paniculatum</i> L. ^{a,b,c,e,f}	jurubeba, juripeba, jupeba	Roots liver disorders, leaves as wound healing	1874	Digestive (Mesia-Vela et al., 2002)
Verbenaceae <i>Lantana camara</i> L. ^{e,f}	camará, cambará	Baths for rheumatism	1874	Anti-inflammatory (Ghosh et al., 2010)
Other historical references.				
^a Cited by Piso (1648) (based in popular name).				
^b Cited in Triaga Brasilica (Pereira et al., 1996).				
^c Cited by Saint-Hilaire in his field book (Brandão et al., 2012).				
^d Cited by Gardner in his manuscripts (Fagg et al., 2015).				
^e Cited by Von Martius in Materiae Medicinae (Martius, 1843).				
^f Cited in 1st edition of Brazilian Official Pharmacopoeia (Brandão et al., 2009).				

and to treat skin ulcers (13), to treat liver disorders, skin diseases and as vermifuge (11), to treat bronchitis, snake bites and as sudorific (10), as antisyphilitic, emollient and to treat rheumatism (9), as bitter, to treat dropsy and wound healing (8), as emetic and to treat jaundice (7), and as emmenagogue and stomachic (6). Other 90 indications were attributed to five or less plants.

Most plants cited by Chernoviz have been already used in Brazil in the previous century, confirming their importance in traditional secular medicine. Informations on some species were already been compiled by the Portuguese priests in the early colonization of the continent. The *Triaga* (*Teriac*), for example, was a Roman ancient remedy used to treat fevers and poisoning, and had part of the components of the original European recipe gradually substituted with plants native to Brazil in 16th century (Pereira et al., 1996). Fourteen plants from the *Triaga Brasilica* were cited in his book: *Cissampelos* sp. and *Chondodendron platiphyllum*, Menispermaceae; *Aristolochia* sp., Aristolochiaceae; *Piper peltatum* and *Piper anisum*, Piperaceae; *Solanum paniculatum*, Solanaceae; *Senna occidentalis* and *Myroxylon balsamum*, Fabaceae; *Carapichea ipecacuanha*, Rubiaceae; *Dorstenia* sp., Moraceae; *Dicyellum caryophyllum*, Lauraceae; *Pradosia lactescens*, Sapotaceae; and *Bixa orellana*, Bixaceae. In the 17th century, the Dutch G. Piso lived for eight years in the northeastern coast and recorded his observations in the book *Historiae Naturalis & Medicae*, published in 1648 (Pisonis, 1648). This work describes the plants used by the Amerindians at the time and was the only source of information on the biodiversity of South America until the late 18th century. Plants registered by Guilherme Piso in the 17th century were also identified in the Chernoviz books (by popular names): *Anacardium occidentale* and *Schinus terebinthifolius*, Anacardiaceae; *Hancornia speciosa*, Apocynaceae; *Bixa orellana*, Bixaceae; *Protium icicariba*, Burseraceae; *Joannesia princeps*, Euphorbiaceae; *Andira fraxinifolia*, Copaifera spp. and *Hymenaea* spp., Fabaceae; *Sassafras albidum*, Lauraceae; *Eugenia ayacuchae* and *Psidium guajava*, Myrtaceae; *Piper* sp., Piperaceae; *Pilocarpus pennatifolius*, Rutaceae; *Solanum paniculatum*, Solanaceae; and

Cecropia palmate, Urticaceae. In 19th century, Portuguese crown opened the borders to the scientific expeditions of naturalists from other European countries. The pioneer was the German Alexander von Humboldt, who explored the area now occupied by Venezuela, Guyana, and the Brazilian states of Amapá and Roraima, from 1799 to 1804. In the 19th century, dozens of other scientists from different parts of Europe travelled to South America, especially Brazil, describing the flora, fauna, mineral wealth and local customs. The contribution of these naturalists to the knowledge of South American biodiversity is incalculable – thousands of plants were brought to the light of science and thousands of new genera and species have been described, based on the materials they collected. From the works of the 19th century's European naturalists, 192 species (80.1%) were also cited by von Martius, 59 species (24.8%) by A. de Saint-Hilaire, and 24 (10%) by G. Gardner. The same traditional was observed for 65% of the species cited by von Martius and 43% of the species cited by Saint-Hilaire. Eleven species cited by Chernoviz were cited by all the three European naturalists: *Gomphrena arborescens*, Amaranthaceae; *Schinus terebinthifolius*, Anacardiaceae; *Hancornia speciosa*, Apocynaceae; *Ilex paraguariensis*, Aquifoliaceae; *Hymenaea courbaril*; Fabaceae; *Deianira* sp., Gentianaceae; *Strychnos pseudoquina*, Loganiaceae; *Cissampelos ovalifolia*, Menispermaceae; *Chiococca alba*, *Palicourea rigida*, Rubiaceae; and *Simarouba versicolor*, Simaroubaceae. This result shows the relevance of the use of such species in the Brazilian traditional medicine on the past. In addition, 51 species were included in the 1st Edition of the Brazilian Official Pharmacopoeia showing their importance also in conventional medicine (Boxes 1 and 2).

It is interesting to note that, on the other side, 34 plants cited by Chernoviz in his work had not been published previously in the consulted bibliography, and these data were probably collected by his own experience as medical doctor in Rio de Janeiro. In fact, Chernoviz's interest in Brazilian medicinal plants was high, since he has incorporated several species along the subsequent editions. While sixty-four plant species (26.9% of the total) were referred as "plants

Box 2: Plants from the Formulary and Medical Guide, their uses and first editions of publication.

Box 2: Plants from the Formulary and Medical Guide, their uses and first editions of publication.				
Family/scientific name	Popular name	Part/indication	Edition	
Adoxaceae <i>Sambucus australis</i> Cham. & Schleidl. ^{e,f}	sabugueiro-do-brasil	Flowers sudorific, roots purgative	1874	
Amaranthaceae <i>Gomphrena arborescens</i> L.f. ^{c,d,e}	paratudo	Bitter, aromatic, debility, diarrhoea, intermittent fevers, snake bites	1874	
Apiaceae <i>Eryngium pristis</i> Cham. & Schleidl. ^e	língua-de-tucano	Diuretic, throat inflammation	1874	
Apocynaceae <i>Allamanda cathartica</i> L. ^e	allamanda cathartica	Cathartic	1920	
<i>Hancornia speciosa</i> Gomes ^{a,c,d,e}	mangabeira	Antituberculosis	1888	
<i>Himatanthus drasticus</i> (Mart.) Plumel ^{c,e}	tiborna, raivosa	Juice drastic, intermittent fevers, jaundice	1874	
<i>Himatanthus phagedaenicus</i> (Mart.) Woodson ^e	sucuúba, sebuu-uva	Juice vermifuge, skin ulcers	1874	
<i>Mandevilla alexicaca</i> (Mart. ex Stadelm.) M.F.Sales ^e	purga-do-campo	Roots purgative, jaundice, oedema of abdominal viscera	1874	
<i>Mandevilla longiflora</i> (Desf.) Pichon ^e	flor-de-babado, flor-de-babeiro	Roots purgative, used in veterinary	1874	
<i>Mandevilla tenuifolia</i> (J.C.Mikan) Woodson ^{d,e}	purga-de-pastor	Roots purgative	1874	
<i>Odontadenia puncticulosa</i> (Rich.) Pulle ^e	cipó-cururú	Purgative	1874	
Araceae <i>Caladium bicolor</i> (Aiton) Vent. ^e	tinhorão, papagaio, bezerro	Leaves for treat tonsilitis	1864	
<i>Monstera adansonii</i> Schott ^e	pau-de-cobra	Stimulant, narcotic, diuretic, purgative, emmenagogue	1897	
<i>Philodendron imbe</i> Schott ex Kunth ^e	imbé, tracuans	Skin ulcers, roots purgative	1874	
<i>Pistia stratiotes</i> L. ^e	flor-d'água, lentilha-d'água	Leaves to treat abcess and other skin diseases	1874	
Araliaceae <i>Hydrocotyle bonariensis</i> Lam. ^e	erva-capitão, erva-do-capitão, acariçoba	Diuretic, liver disorders	1874	
Arecaceae <i>Copernicia prunifera</i> (Mill.) H.E.Moore ^{d,f}	carnaúba	Roots diuretic, syphilis, rheumatism	1892	
Asparagaceae <i>Herreria salsaparrilha</i> Mart. ^e	salsaparrilha, japecanga	Syphilis	1864	
Asteraceae <i>Acmella oleracea</i> (L.) R.K.Jansen ^e	agrião-do-pará, jambú, nhambú, jambú-açú	Excitant, antiescorbutic	1864	
<i>Baccharis articulata</i> (Lam.) Pers. ^e	carqueja-doce	Tonic, febrifuge and for aromatic baths	1874	
<i>Baccharis ochracea</i> Spreng. ^e	erva-santa	Stomachic	1874	
<i>Baccharis</i> spp.	alecrim-selvagem, alecrim-do-mato	Stimulant, aromatic baths	1864	
<i>Conoclinium prasiiifolia</i> (DC.) R.M.King & H.Rob. ^e	maria-preta	Stimulant	1874	
<i>Eclipta erecta</i> L. ^e	tangaracá	Antidiarrheal	1874	
<i>Elephantopus mollis</i> Kunth ^{c,e}	erva-collegio, fumo-bravo, erva-grossa, suçuáya	Leaves emollient, roots to treat bronchitis and fevers	1864	
<i>Mikania cordifolia</i> (L.f.) Willd. ^e	erva-de-cobra	Snake bites	1874	
<i>Mikania guaco</i> Kunth	guaco	Snake bites, cholera, respiratory diseases	1874	
<i>Mikania officinalis</i> Mart. ^e	coração-de-jesú	Tonic, stimulant and intermittent fevers	1874	
<i>Sonchus levis</i> Vell.	serralha	Desobstruent and depurative	1874	
Begoniaceae <i>Begonia hirtella</i> Link ^e	surucura	Astringent, antidiarrheal	1864	
<i>Begonia</i> spp. ^e	azedinha-do-brejo, do-sapo	Astringent, antidiarrheal	1874	
Bignoniaceae <i>Anemopaegma arvense</i> (Vell.) Stellfeld ex de Souza	catuaba	Aphrodisiac	1920	
<i>Crescentia cujete</i> L. ^e	cabacinha	Barks antidiarrheal, fruits laxative and expectorant	1897	
<i>Handroanthus serratifolius</i> (Vahl) S.Grose ^e	ipeuva, cinco-folhas	Diuretic, depurative, to treat syphilis and rheumatism	1897	
<i>Jacaranda caroba</i> (Vell.) DC. ^{c,f}	caroba	Leaves against syphilis	1874	
<i>Jacaranda procera</i> (Willd.) Spreng. ^c	caroba	Antisyphilitic	1864	
Boraginaceae <i>Echium plantagineum</i> L. ^e	borracha-chimaroná	Leaves emollient	1874	
<i>Heliotropium elongatum</i> (Lehm.) I.M.Johnst. ^e	crista-de-galo	Respiratory disease	1874	
Brassicaceae <i>Coronopus didymus</i> (L.) Sm. ^e	mastroço	Antiescorbutic	1864	
Bromeliaceae <i>Tillandsia usneoides</i> (L.) L. ^d	barba-de-velho	Antihemorrhoidal	1888	
Burseraceae <i>Protium icicaria</i> (DC.) Marchand ^{a,e,f}	elemi	Skin diseases	1864	

Cactaceae <i>Hylocereus setaceus</i> (Salm-Dyck) R.Bauer ^e	jamacurú, cumbéba, mandacarú, urumbéba	Fruits antiescorbutic, fevers, gastritis, pulmonary and skin disease	1874
Calophyllaceae <i>Kielmeyera speciosa</i> A.St.-Hil. ^{c,e}	malva-do-campo, folha-santa, pinhão	Leaves emollient	1874
Cannaceae <i>Canna glauca</i> L. ^e	imberi, albará, erva-dos-feridos	Rheumatic pain, wound healing and skin ulcers	1874
Caricaceae <i>Jacaratia spinosa</i> (Aubl.) A.DC.	jaracatiá	Juice of fruit is vermifuge	1874
Celastraceae <i>Hippocratea volubilis</i> L.	amendoeira-do-mato	Febrifuge, snake bites, expectorant	1920
Clusiaceae <i>Clusia insignis</i> Mart. ^e	quapoy, aqui	Flowers for dry skin	1888
Combretaceae <i>Terminalia argentea</i> Mart. ^e	caxaporra-do-gentio	Resin purgative	1874
Commelinaceae <i>Commelina erecta</i> L. ^{c,e} <i>Tripogandra diuretica</i> (Mart.) Handlos ^{e,f}	trapoeraba-rana, marianinha trapoeraba	Diuretics, dropsy, baths for rheumatism Diuretics, dropsy	1874 1864
Convolvulaceae <i>Cuscuta umbellata</i> Kunth ^{c,e,f}	cipó-de-chumbo	Juice as wound healing	1864
Costaceae <i>Costus spiralis</i> (Jacq.) Roscoe	canna-do-brejo, macaco, matto, pacocaatinga, uacayá	Leucorrhea	1874
Cucurbitaceae <i>Cayaponia espelina</i> (Silva Manso) Cogn. ^f	cerejeira-de-purga, espelina, tomba	Tonic, stomachic, purgative, emetic, liver disorders	1874
<i>Cayaponia pilosa</i> (Vell.) Cogn. ^e	cayapó, anna-pinta	Seeds purgative, depurative, skin disease, emmenagogue	1874
<i>Cayaponia martiana</i> (Cogn.) Cogn. ^e	tayuyá, abóbora-do-mato,	Fruits dropsy, purgative, amenorrhea, leaves to treat ulcers	1864
<i>Fevillea cordifolia</i> L. ^e	nhandiroba	stomachic, purgative, emetic, liver disorders	1874
<i>Luffa operculata</i> (L.) Cogn. ^e	bucha-dos-paulistas, purga-de-joão-paes, buchinha	Purgative	1874
Cyatheaceae <i>Cyathea microdonta</i> (Desv.) Domin ^e	rabo-de-bugio	Bronchitis	1874
Cyperaceae <i>Kyllinga odorata</i> Vahl ^e	capim-cheiroso, cidreira, marinho, acapé, jarapé	Carminative, antispasmodic, sudorific, diuretic	1874
Dilleniaceae <i>Davilla rugosa</i> Poir. ^{c,e,f}	sambaibinha, cipó-de-carijó, cipó-de-caboclo	Leaves to treat oedema of legs and testicles	1874
Droseraceae <i>Drosera</i> spp. ^c	drosera	Antitussigen	1920
Euphorbiaceae <i>Cnidoscolus urens</i> (L.) Arthur var. <i>urens</i> ^b	queimadeira, pinha	Impingen	1888
<i>Croton campestris</i> A.St.-Hil., <i>C. fulvus</i> Mart. ^{c,e}	velame-do-campo	Roots purgative	1874
<i>Euphorbia hyssopifolia</i> L. ^{c,e}	erva-de-santa-luzia	Eyes disorders, skin ulcers	1864
<i>Euphorbia papillosa</i> A.St.-Hil. ^e	maleiteira, leiteira	Purgative	1874
<i>Euphorbia potentilloides</i> Boiss. ^e	andorinha, erva-de-passarinho	Skin and eyes ulcers	1874
<i>Hevea guianensis</i> Aubl. ^e	borracha, cauchuc, goma elástica, seringa	Tuberculosis	1874
<i>Hura crepitans</i> L. ^e	assacú	Leprosy	1864
<i>Jatropha elliptica</i> (Pohl) Oken ^{c,e}	jalapão, tiú, raiz-de-lagarto	Roots purgative, jaundice, dropsy	1874
Fabaceae <i>Andira fraxinifolia</i> Benth. ^{a,c,e}	angelim	Vermifuge	1864
<i>Andira inermis</i> (W.Wright) DC.	andira inermis	Bark vermiculite and purgative	1897
<i>Arachis hypogaea</i> L. ^{e,f}	mandobi, amendoim	Aphrodisiac	1874
<i>Dipteryx odorata</i> (Aubl.) Willd. ^e	cumarí, fava-tonka	Tonic	1874
<i>Hymenaea stigonocarpa</i> Mart., <i>Hymenaea martiana</i> Hayne, <i>Hymenaea courbaril</i> L. ^{a,c,d,e}	jatahy, jatahyba, jetehy, jatobá	Resin hemoptysis	1874
<i>Leptolobium elegans</i> Vogel	perobinha-do-campo	To treat epilepsy, antisiphilitic	1920
<i>Myroxylon balsamum</i> (L.) Harms ^{b,f}	bálsamo-de-tolú	Bronchitis	1864
<i>Pachyrhizus erosus</i> (L.) Urb.	jacatupé	Antidiarrheal, antihemorrhoidal	1888
<i>Senna occidentalis</i> (L.) Link ^{b,e,f}	fedegoso, pajámariobá, pájomarioba	Roots diuretic, tonic, dropsy	1864
<i>Senna rugosa</i> (G.Don) H.S.Irwin & Barneby ^e	amendoirana, alcaçuz bravo, bico-de-corvo, boi gordo, paratudo	Vermifuge	1874
Gentianaceae <i>Calolisianthus pendulus</i> (Mart.) Gilg,	genciana-brasileira	Roots febrifuge	1874
<i>Calolisianthus amplissimus</i> (Mart.) Gilg ^e			

<i>Deianira</i> sp. ^{c,d,e}	centáurea-menor-do-brasil	Bitter flowers sell as centaurea in Europe	1892
<i>Potalia resinifera</i> Mart. ^e	anabi	Leaves astringent, to treat ophthalmia	1874
Humiriaceae			
<i>Endopleura uchi</i> (Huber) Cuatrec.	uxi	Seeds hemostatic	1874
Hypericaceae			
<i>Vismia guianensis</i> (Aubl.) Choisy ^e	caaopiá, pau-de-lacre	Purgative	1874
Iridaceae			
<i>Trimezia galaxioides</i> (Gomes) Ravenna ^e	baririçó, maririçó	Roots purgative	1864
<i>Trimezia juncifolia</i> (Klatt) Benth. & Hook.f. ^{d,e}	ruibarbo-do-campo, pireto	Roots purgative	1920
Krameriaceae			
<i>Krameria argentea</i> Mart. ex Spreng. ^{e,f}	ratânia-do-brasil, da-terra	Astringent	1888
Lamiaceae			
<i>Hyptis radicans</i> (Pohl) Harley & J.F.B.Pastore ^{e,f}	paracary, hortelâ-brava, mentrasto, meladinha	Asthma and snake bites	1874
<i>Leonotis nepetifolia</i> (L.) R.Br. ^{e,f}	cordão-de-frade, cordão-de-são-francisco, leonuro	Excitant, baths for children	1864
<i>Ocimum americanum</i> L. ^e	alfavaca-do-campo, remedio-do-vaqueiro	Excitant, sudorific, constipation	1864
<i>Vitex megapotamica</i> (Spreng.) Moldenke ^e	tarumá	Baths for rheumatism	1874
Lauraceae			
<i>Aniba puchury-minor</i> (Mart.) Mez ^e	puchury-miri	Same as Puchury	1874
<i>Cinnamomum verum</i> J.Presl ^e	canela	Dyspepsy, antidiarrheal	1920
<i>Dicyellum caryophyllum</i> (Mart.) Nees ^{b,e,f}	pau-de-cravo, cravo-maranhão, imyráquiynhá	Aromatic	1874
<i>Ocotea cujumary</i> Mart. ^e	cujumary	Seeds for dyspepsy	1874
<i>Ocotea cymbarum</i> Kunth ^e	sassafrás-do-brasil, pau-sassafrás	Barks tonic	1874
<i>Ocotea longifolia</i> Kunth ^e	canela-de-cheiro	Oil for rheumatic pain and spasm of muscle	1874
<i>Nectandra reticulata</i> (Ruiz & Pav.) Mez ^e		Aromatic, emmenagogue and stomachic	1874
<i>Sassafras albidum</i> (Nutt.) Nees ^a	sassafrás	Syphilis, wound healing, rheumatism	1864
Lecythidaceae			
<i>Cariniana legalis</i> (Mart.) Kuntze	jequitibá	Astringent, antidiarrheal	1864
<i>Eschweilera grandiflora</i> (Aubl.) Sandwith ^e	sapucaieira	Diuretic, jaundice, hepatites, intermittent fevers	1874
<i>Gustavia hexapetala</i> (Aubl.) Sm. ^e	janiparindiba	Leaves for liver disorders	1874
Linderniaceae			
<i>Lindernia difusa</i> (L.) Wettst. ^e	caa-ataya, erva-de-ferro, mata cana, oreilha de rato, purga de joão paes	Emetic	1874
Loganiaceae			
<i>Strychnos toxifera</i> R.H.Schomb. ex Benth. ^e	curare, ervadura, uiráy, ticúna, woorara	Muscle relaxant	1874
Malvaceae			
<i>Guazuma ulmifolia</i> Lam. ^{c,e}	mutamba, guazuma	Astringent, skin/scalp disease	1897
<i>Helicteres sacarolha</i> A. St.-Hil. ^{c,e}	sacarrolha, rosea-para-mulas	Venereal disease, emollient	1874
<i>Pavonia sidifolia</i> Kunth ^e	malva diurética	Emollient, diuretic, dysuria	1888
<i>Sida planicaulis</i> Cav. ^e	vassoura	Emollient	1864
<i>Sphaeralcea bonariensis</i> (Cav.) Griseb ^e	malvaíscio	Emollient against cough	1888
<i>Triumfetta eriocarpa</i> A.St.-Hil., <i>T. semitriloba</i> Jacq., <i>T. sepium</i> A.St.-Hil., Juss. & Cambess. ^e	carrapicho de calçada	Blennorrhea	1874
<i>Urena lobata</i> L. ^{c,e,f}	guaxima, malvaíscio	Emollient	1864
<i>Urena sinuata</i> L. ^e	carrapicho	Flowers emollient against cough	1874
<i>Waltheria communis</i> A.St.-Hil. ^{c,e,f}	douradinha	Emollient against cough	1874
Melastomaceae			
<i>Miconia cinnamomifolia</i> (DC.) Naudin	erva dutra	Leaves antidiarrheal	1874
Meliaceae			
<i>Guarea macrophylla</i> ssp. <i>spicaeflora</i> (A.Juss.) T.D.Penn. ^e	marinheiro de folha larga, tuaiussú, utuapoca	Bark and roots purgative	1874
Menispermaceae			
<i>Chondrodendron platiphyllum</i> (A.St.-Hil.) Miers	abutua, butua, parreira brava	Roots for intermittent fevers	1864
<i>Cissampelos glaberrima</i> A.St.-Hil. ^{b,e}	cipó de cobra, caapeba, erva-de-nossa-senhora	Snake bites	1874
Moraceae			
<i>Dorstenia arifolia</i> Lam., <i>D. brasiliensis</i> Lam. ^{b,c,e,f}	contraerva, caapiá, carapiá, caxapiá, cayapiá	Roots excitant, fevers, emmenagogue	1874
<i>Ficus adhatodifolia</i> Schott ex Spreng. ^e	coajinguva, cuáxinguba	Vermifuge	1874
<i>Ficus gomelleira</i> Kunth & C.D.Bouché ^e	gameleira, figueira-branca	Vermifuge, purgative	1888
Myristicaceae			
<i>Virola bicuhyba</i> (Schott ex Spreng.) Warb. ^e	bicuiba, bucuúba, moscadeira-do-brasil	Seeds to treat rheumatic pain	1874
<i>Virola sebifera</i> Aubl. ^e	ucuuba	Rheumatism	1874

Myrtaceae				
<i>Blepharocalyx salicifolius</i> (Kunth) O.Berg,	guabiroba	Leaves antidiarrheal, leucorrhea	1874	
<i>Campomanesia pubescens</i> (Mart. ex DC.) O.Berg ^{c,e}				
<i>Eugenia ayacuchae</i> Steyermark ^{a,d,f}	jaboticaba	Astringent, fruit for treat tonsilitis	1864	
<i>Eugenia pyriformis</i> Cambess. ^e	uvalheira	Refreshing	1874	
Nyctaginaceae				
<i>Boerhavia coccinea</i> Mill. ^{c,f}	erva-tostão, tangaraca	Diuretic, liver disorders	1864	
Ochnaceae				
<i>Ouratea</i> sp. ^c	batiputá	Oil from seeds to treat rheumatism	1888	
Orchidaceae				
<i>Vanilla planifolia</i> Jacq. ex Andrews ^{e,f}	baunilha	Stimulant, aphrodisiac, emmenagogue, diuretic	1864	
Phyllanthaceae				
<i>Phyllanthus brasiliensis</i> (Aubl.) Poir. ^b	conabi	Leaves emetic	1874	
Phytolaccaceae				
<i>Petiveria alliacea</i> L. ^{e,f}	pipi, raiz-de-guiné	Stimulant	1864	
Piperaceae				
<i>Piper anisum</i> (Spreng.) Angely ^{b,c,e}	jaborandi	Aromatic, sialagogue, odontalgic	1874	
Plumbaginaceae				
<i>Limonium brasiliense</i> (Boiss.) Kuntze ^f	guaycuru, baycuru	Roots astringent	1897	
<i>Plumbago scandens</i> L. ^{d,e}	queimadeira, loco, caa-pomonga	Vesicant, ear ache	1888	
Polygonaceae				
<i>Acantocladius brasiliensis</i> Klotzsch ex Hassk.	laranjeirinha-do-mato	Cholics	1874	
Poaceae				
<i>Andropogon bicornis</i> L. ^b	sapé	Diuretic	1864	
<i>Stenotaphrum secundatum</i> (Walter) Kuntze ^e	grama-da-praia	Roots emollient, diuretic, to treat inflammation	1874	
Polygonaceae				
<i>Polygonum punctatum</i> Elliott, <i>Muehlenbeckia sagittifolia</i> (Ortega) Meisn. ^{c,e}	erva-de-bicho, acataya, capeticova, cataya, persicaria, pimenta-d'água	Stimulant, diuretic, emmenagogue, antihemorrhoidal	1864	
Rosaceae				
<i>Prunus myrtifolia</i> (L.) Urb. ^e	cerejeira, gingeira-brava	Stimulant, nevralgy, liver disorders, antiemetic	1874	
Rubiaceae				
<i>Chiococca alba</i> (L.) Hitchc. ^{c,d,e}	cainca, cainana, cipó-cruz, cruzeirinha, fedorenta, raiz-de-frade, preta	Roots diuretic, purgative, dropsy, snake bites	1864	
<i>Coutarea hexandra</i> (Jacq.) K.Schum. ^{d,e}	quina-do-brasil, quina-de-pernambuco	Bark intermittent fevers	1864	
<i>Ladenbergia cujabensis</i> Klotzsch	quina-de-cuiabá	Barks bitter, to treat intermittent fevers	1864	
<i>Ladenbergia hexandra</i> (Pohl) Klotzsch	quina-do-rio-de-janeiro	Bark intermittent fevers	1874	
<i>Ladenbergia lambertiana</i> (Mart.) Klotzsch ^e	quina-do-brasil	Barks bitter, to treat intermittent fevers	1864	
<i>Remijia firmula</i> (Mart.) Wedd. ^e	quina-do-brasil	Barks bitter, to treat intermittent fevers	1864	
<i>Remijia macrocnemias</i> (Mart.) Wedd. ^e	quina-do-brasil	Barks bitter, to treat intermittent fevers	1864	
<i>Rustia formosa</i> (Cham. & Schldl. ex DC.) Klotzsch ^e	quina-do-brasil	Barks bitter, to treat intermittent fevers	1864	
<i>Schizocalyx cuspidatus</i> (A.St.-Hil.) Kainul. & B.Bremer ^e	quina-do-brasil, quina-do-mato	Bark intermittent fevers	1864	
Rutaceae				
<i>Angostura trifoliata</i> (Willd.) T.S.Elias	angustura	Tonic, antidiarrheal, diaphoretic, sudorific, bronchitis	1874	
<i>Esenbeckia febrifuga</i> (A.St.-Hil.) A.Juss. ex Mart. ^{c,e}	três-folhas-vermelhas, laranjeira-do-mato, quina	Barks as febrifuge	1874	
<i>Galipea jasminiflora</i> (A.St.-Hil.) Engl. ^{c,e}	três-folhas-brancas, quina-falsa	Barks for treat intermittent fevers	1874	
<i>Pilocarpus pennatifolius</i> Lem. ^a	jaborandi	Sudorific, sialagogue, bronchitis, hydropsy, fevers	1874	
<i>Zanthoxylum tingoassuiba</i> A.St.-Hil. ^{c,f}	tinguaciba	Intermittent fevers	1888	
Sapindaceae				
<i>Cupania vernalis</i> A.St.-Hil.	caboatan-de-capoeira	Barks for asthma and cough	1888	
<i>Paullinia pinnata</i> L. ^e	timbó, cururu-apé	Narcotic, sedative	1864	
Sapotaceae				
<i>Manilkara excelsa</i> (Ducke) Standl.	massaranduba	Respiratory disorders	1874	
<i>Pouteria caitito</i> (Ruiz & Pav.) Radlk. ^e	abiaba	Antidiarrheal, intermittent fevers	1897	
<i>Pradosia lactescens</i> (Vell.) Radlk. ^{b,f}	monésia, buranhém, guaranhém	Baths to treat erysipela	1864	
Scrophulariaceae				
<i>Buddleja stachyoides</i> Cham. & Schldl. ^{c,e,f}	barbasco, verbasco, calção-de-velho	Leaves and flowers emollient	1874	
Simaroubaceae				
<i>Simarouba versicolor</i> A.St.-Hil. ^{c,d,e}	parahyba	Bark against head lice	1874	
Siparunaceae				
<i>Siparuna brasiliensis</i> (Spreng.) A.DC.	limoeiro-bravo	Leaves for contusion, respiratory disease	1874	

Solanaceae			
<i>Brunfelsia uniflora</i> (Pohl) D.Don ^{d,e}	manacá, camgabá, geratacaca, jeretaca	Roots purgative, antisyphilitic	1874
<i>Solanum aculeatissimum</i> Jacq.	arrebanta-cavallo	Digestive disorders, skin diseases	1888
<i>Solanum americanum</i> Mill. ^{c,e}	giquirioba, jiquirioba	Skin disease	1892
<i>Solanum cernuum</i> Vell. ^{c,e}	velame-do-mato, braço-de-preguiça	Leaves liver disorders	1874
<i>Solanum pseudoquina</i> A. St.-Hil. ^{c,e}	quina-de-são-paulo	Bark intermittent fevers	1874
Styracaceae			
<i>Styrax maninul</i> B.Walln., <i>S. ferrugineus</i> Nees & Mart. ^e	estoraque-do-brasil	Stimulant	1897
Urticaceae			
<i>Bohemeria caudata</i> Sw. ^e	asa-peixe	Baths antihemorrhoidal and diuretic	1874
<i>Cecropia palmata</i> Willd. ^{a,c,e}	ambayba, imbayba, umbuba, árvore-da-preguiça	Skin disease, venereal disease	1864
Verbenaceae			
<i>Glandularia microphylla</i> (Kunth) Cabrera ^e	alecrim-do-campo	Aromatic baths	1892
<i>Lippia brasiliensis</i> (Link) T.R.S.Silva ^e	camará, erva-sagrada	Febrifuge	1897
<i>Lippia pseudothea</i> (A.St.-Hil.) Schauer ^{c,e,f}	chá-de-pedestre, chá-de-frade	Stimulant	1874
<i>Stachytarpheta jamaicensis</i> (L.) Vahl ^e	gervao, orgiâo, urgevão	Leaves sudorific and stimulant	1864
Violaceae			
<i>Anchieta pyrifolia</i> (Mart.) G.Don ^{c,e,f}	cipó-sumá, piragaia	Roots purgative	1874
Winteraceae			
<i>Drimys brasiliensis</i> Miers ^{c,e}	casca-de-anta, paratudo	Antispasmodic	1874
Zingiberaceae			
<i>Renealmia aromatica</i> (Aubl.) Griseb., <i>Renealmia alpinia</i> (Rottb.) Maas ^e	pacová, cuité-açú, paco-seroca	Roots stomachic, to treat skin ulcers	1874

^a Cited by Piso (1648) (based in popular name).

^b Cited in Triaga Brasílica (Pereira et al., 1996).

^c Cited by Saint-Hilaire in his field book (Brandão et al., 2012).

^d Cited by Gardner in his manuscripts (Fagg et al., 2015).

^e Cited by Von Martius in Materiae Medicae (Martius, 1843).

^f Cited in 1st edition of Brazilian Official Pharmacopoeia (Brandão et al., 2009).

from Brazil" in the 6th edition from 1864, higher numbers were observed in subsequent editions, reaching a total of 238 in the last edition, from 1920.

Among the traditional uses, purgative plants were the most cited by the author (30 species), being 21 new species included in the edition of 1874. Purgatives were widely used in the 19th century due to the belief that diseases were caused by a "dirty body" and it was necessary to "purge", for cleansing. Purgatives from plants are no longer used in the medical practice, and this fact can explain the existence of few data about them in the medical literature – only a report for *J. princeps* was found, describing episodes of diarrhoea after its use (Nishioka and Escalante, 1997), and two studies showing the laxative effect of *Operculina macrocarpa* in mice (Michelin and Salgado, 2004; Paganotte et al., 2016). On the other side, plants used to treat diarrhoea were also very important in the 19th century and the most known and used preparation was made of ipecacuanha root, *Carapichea ipecacuanha*, Rubiaceae. The beneficial effect of this plant described as early as the 16th century is due to the presence of the alkaloids emetine and cephaline. These substances act specifically on *Entamoeba* parasites that cause diarrhoea, which were prevalent and often fatal in Chernoviz's time. Although eighteen Brazilian plants were cited as antidiarrheal, besides *C. ipecacuanha*, only *Psidium guajava* had this property evaluated (Ojewole et al., 2008; Birdi et al., 2010, 2014). *Cacalia mentrasto* and *Simarouba amara* were evaluated as antimicrobial (Caceres et al., 1990; Adetutu et al., 2012; Singh et al., 2013), which can be related to an antidiarrheal function.

A total of 55 species were cited as useful to treat general skin diseases in the six studied editions, including lupus, acne, eczema, psoriasis, leprosy, astringent, emollient, as wound healing, to treat skin ulcers, abscess, burns, dry skin, erysipela and impingen. From these, eighteen studies were performed to date in order to verify the efficacy of these plants (Box 1). Other 21 plants were registered as diuretics, nine being cited already in the first edition (*Tripogandra diuretica*, *Senna occidentalis*, *Boerhavia hirsuta*, *Phyllanthus* spp.,

Andropogon bicornis and *Polygonum* spp., *Acanthospermum australe*, *Chiococca alba* and *Vanilla planifolia*). Despite the higher number of species used and their old use, only *Phyllanthus niruri* has been submitted to pharmacological studies, in order to verify its diuretic activity (Micali et al., 2006; Giribabu et al., 2014). Fever was a very common health problem at that time, but the most feared was the intermittent fever, showing typical symptoms of malaria. The medical cure for malaria originated in the 17th century when the Spaniards learned, from the Native Americans living in Peru, the use of the bark from *Cinchona* spp., especially *Cinchona succirubra* Pav. ex Klotzsch (Rubiaceae), known as "china" or "quina". In the 19th century, the Portuguese stimulated the search for substitutes for *Cinchona* and many species were named "quina" (Cosenza et al., 2013). Among the 39 plants registered in the books to treat fevers, 26 species were pointed by Chernoviz as useful to treat intermittent fevers, among them seven species of Brazilian quinas (*Strychnos pseudoquina*, *Ladenbergia hexandra*, *Remijia ferruginea*, *Schizocalyx cuspidatus*, *Esenbeckia febrifuga*, *Galipea jasminiflora*, and *Solanum pseudoquina*). From these, only *Strychnos pseudoquina* and *R. ferruginea* were evaluated to date as antimalarial and they showed a moderated activity (Andrade-Neto et al., 2003).

Until the first years of the 20th century, the Pharmacopoeia General from Portugal published in 1794, was the official medical reference used in Brazil. In this Pharmacopoeia, few Brazilian species were officialized, among them abutua (*Chondrodendron platiphyllum*, Menispermaceae), copahyba balsam (*Copaifera* spp., Fabaceae), contraherva (*Dorstenia* spp., Moraceae), Elemi or Almacega from Brazil (*Protium icicaria*, Proteaceae), ipecacuanha (*Carapichea ipecacuanha*, Rubiaceae), ipecacuanha branca (*Pombalia calceolaria*, Violaceae) and *Smilax salsaparrilha*, Smilacaceae. The Portuguese Pharmacopoeia was replaced by the Chernoviz's *Formulary and Medical Guide* and this book was then required in all Pharmacies in Brazil. This occurred until the publication of the first edition of the Brazilian Pharmacopoeia (FBRAS), in 1926. Fifty-one native plant species previously described by Chernoviz were then

formalized in the conventional medicine by their inclusion in the first edition of FBRAS (Brandão et al., 2009). The Brazilian plants, cited by Chernoviz, and included in the 1st edition of FBRAS, are examples of American medicinal plants used both in traditional and conventional medicine.

It is unquestionable that the rational use of medicinal plants should be supported by proper laboratorial investigations and clinical trials (Cravotto et al., 2010; Atanasov et al., 2015). Sixty-one species cited by Chernoviz (25.6%) have been submitted to *in vitro* and pre-clinical studies and their traditional uses were confirmed. We argue that these plants must be prioritized in development of medicines, submitting them to clinical studies or by considering their traditionality, as established by WHO. It is also very important to define strategies for the protection of the biodiversity and the Native American traditional heritage.

Authors' contributions

All authors contributed to the revision and writing of the manuscript. JPS reviewed and updated the botanical names. The results presented here are part of the PhD thesis or LMR.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We thank CNPq (563563/2010 REFLORA) for financial support and all the Libraries for making their collections available to our studies. We are also grateful to Dr. Maria Fernanda Calió (Universidade Federal de São Carlos, São Carlos, Brazil), Dr. Rubens Queiroz (Universidade Federal da Paraíba, João Pessoa, Brazil), Dr. Pedro Luís Rodrigues de Moraes (Universidade Estadual Paulista, Rio Claro, Brazil), Dr. Alexandre Zuntini (Universidade de São Paulo, São Paulo, Brazil), Dr. Camila Dellanheze Inácio and Dr. Lilian Eggers (Universidade Federal do Rio Grande do Sul), and Dr. Juliana Gastaldello Rando (Universidade do Oeste da Bahia) for their valuable nomenclatural aid. We are also grateful to Tamara Castro for the help in revision of the books.

References

- Ademola, I.O., Fagbemi, B.O., Idowu, S.O., 2007. *Anthelmintic activity of Spigelia anthelmia extract against gastrointestinal nematodes of sheep*. Parasitol. Res. 101, 63–69.
- Adetutu, A., Morgan, W.A., Corcoran, O., Chimezie, F., 2012. *Antibacterial activity and in vitro cytotoxicity of extracts and fractions of Parkia biglobosa (Jacq.) Benth. stem bark and Ageratum conyzoides Linn. leaves*. Environ. Toxicol. Pharmacol. 34, 478–483.
- Agra, I.K., Pires, L.L., Carvalho, O.S., Silva-Filho, E.A., Smanniotti, S., Barreto, E., 2013. *Evaluation of wound healing and antimicrobial properties of aqueous extract from Bowdichia virgilioides stem barks in mice*. An. Acad. Bras. Cienc. 85, 945–954.
- Andrade-Neto, V.F., Brandão, M.G., Stehmann, J.R., Oliveira, L.A., Krettli, A.U., 2003. *Antimalarial activity of Cinchona-like plants used to treat fever and malaria in Brazil*. J. Ethnopharmacol. 87, 253–256.
- Annan, K., Houghton, P.J., 2008. *Antibacterial, antioxidant and fibroblast growth stimulation of aqueous extracts of Ficus asperifolia Miq. and Gossypium arboreum L. wound-healing plants of Ghana*. J. Ethnopharmacol. 119, 141–144.
- Araujo, A.A., Soares, L.A., Assunção Ferreira, M.R., Souza Neto, M.A., Silva, G.R., Araújo Jr., R.F., Guerra, G.C., Melo, M.C., 2014. *Quantification of polyphenols and evaluation of antimicrobial, analgesic and anti-inflammatory activities of aqueous and acetone-water extracts of Libidibia ferrea, Parapiptadenia rigida and Psidium guajava*. J. Ethnopharmacol. 156, 88–96.
- Atanasov, A.G., Waltenberger, B., Pferschy-Wenzig, E.M., Linder, T., Wawrosch, C., Uhrin, P., Temml, V., Wang, L., Schwaiger, S., Heiss, E.H., Rollinger, J.M., Schuster, D., Breuss, J.M., Bochkov, V., Mihovilovic, M.D., Kopp, B., Bauer, R., Dirsch, V.M., Stuppner, H., 2015. *Discovery and resupply of pharmacologically active plant-derived natural product: a review*. Biotechnol. Adv. 33, 1582–1614.
- Avila-Blanco, M.E., Rodríguez, M.G., Moreno, D.J.L., Muñoz-Ortega, M., Ventura-Juárez, J., 2014. *Amoebicidal activity of essential oil of Dysphania ambrosioides* (L.) Mosyakin & Clemants in an amoebic liver abscess hamster model. Evid. Based Complement. Alternat. Med., <http://dx.doi.org/10.1155/2014/930208>.
- Badilla, B., Mora, G., Poveda, L.J., 1999. *Anti-inflammatory activity of aqueous extracts of five Costa Rican medicinal plants in Sprague-Dawley rats*. Rev. Biol. Trop. 47, 723–727.
- Barros, W.M., Rao, V.S., Silva, R.M., Lima, J.C., Martins, D.T., 2010. *Anti-inflammatory effect of the ethanolic extract from Bowdichia virgilioides H.B.K. stem bark*. An. Acad. Bras. Cienc. 82, 609–616.
- Bastos, G.N., Santos, A.R., Ferreira, V.M., Costa, A.M., Bispo, C.I., Silveira, A.J., Nascimento, J.L., 2006. *Antinociceptive effect of the aqueous extract obtained from roots of Physalis angulata L. on mice*. J. Ethnopharmacol. 103, 241–245.
- Birdi, T.J., Daswani, P., Brijesh, S., Tetali, P., Natu, A., Antia, N., 2010. *Newer insights into the mechanism of action of Psidium guajava L. leaves in infectious diarrhoea*. BMC Complement. Altern. Med., <http://dx.doi.org/10.1186/1472-6882-10-33>.
- Birdi, T.J., Brijesh, S., Daswani, P.G., 2014. *Bactericidal effect of selected antidiarrhoeal medicinal plants on intracellular heat-stable enterotoxin-producing Escherichia coli*. Indian J. Pharm. Sci. 76, 229–235.
- Brandão, M.G.L., Zanetti, N.N., Oliveira, P., Grael, C.F., Santos, A.C., Monte-Mór, R.L., 2008. *Brazilian medicinal plants described by 19th century European naturalists and in the Official Pharmacopoeia*. J. Ethnopharmacol. 120, 141–148.
- Brandão, M.G.L., Cosenza, G.P., Grael, C.F.F., Netto Junior, N.L., Monte-Mór, R.L.M., 2009. *Traditional uses of American plant species from the 1st edition of Brazilian Official Pharmacopoeia*. Rev. Bras. Farmacogn. 19, 478–487.
- Brandão, M.G.L., Pignal, M., Romaníu, S., Grael, C.F.F., Fagg, C.W., 2012. *Useful Brazilian plants listed in the field books of the French naturalist Auguste de Saint-Hilaire (1779–1853)*. J. Ethnopharmacol. 143, 488–500.
- Breitbach, U.B., Niehues, M., Lopes, N.P., Faria, J.Q., Brandão, M.G.L., 2013. *Amazonian Brazilian medicinal plants described by C.F.P. von Martius in the 19th century*. J. Ethnopharmacol. 147, 180–189.
- Caceres, A., Cano, O., Samayoa, B., Aguilar, L., 1990. *Plants used in Guatemala for the treatment of gastrointestinal disorders. 1. Screening of 84 plants against enterobacteria*. J. Ethnopharmacol. 30, 55–73.
- Camporese, A., Balick, M.J., Arvigo, R., Esposito, R.G., Morsellino, N., Simone, F., Tubaro, A., 2003. *Screening of anti-bacterial activity of medicinal plants from Belize (Central America)*. J. Ethnopharmacol. 87, 103–107.
- Carlini, E.A., Oliveira, A.B., Oliveira, G.G., 1983. *Psychopharmacological effects of the essential oil fraction and of the hydrolate obtained from the seeds of Licaria puchury-major*. J. Ethnopharmacol. 8, 225–236.
- Carvalho, L.H., Krettli, A.U., 1991. *Antimalarial chemotherapy with natural products and chemically defined molecules*. Mem. I. Oswaldo Cruz 86, 181–184.
- Carvalho, L.H., Brandão, M.G.L., Santos-Filho, D., Lopes, J.L., Krettli, A.U., 1991. *Antimalarial activity of crude extracts from Brazilian plants studied in vivo in Plasmodium berghei-infected mice and in vitro against Plasmodium falciparum in culture*. Braz. J. Med. Biol. Res. 24, 1113–1123.
- Castello, M.C., Phatak, A., Chandra, N., Sharon, M., 2002. *Antimicrobial activity of crude extracts from plant parts and corresponding calli of Bixa orellana L*. Indian J. Exp. Biol. 40, 1378–1381.
- Castro, O., Barrios, M., Chinchilla, M., Guerrero, O., 1996. *Biological and chemical evaluation of the effect of plant extracts on Plasmodium berghei*. Rev. Biol. Trop. 44, 361–367.
- Chernoviz, P.L.N., 1864. *Formulário e Guia Médica*, 6th ed. Em casa do Autor, Pariz.
- Chernoviz, P.L.N., 1874. *Formulário e Guia Médico*, 9th ed. Em casa do Autor, Pariz.
- Chernoviz, P.L.N., 1888. *Formulário e Guia Médico*, 13th ed. Livraria de A. Roger e F. Chernoviz, Pariz.
- Chernoviz, P.L.N., 1892. *Formulário e Guia Médico*, 14th ed. Livraria de A. Roger e F. Chernoviz, Pariz.
- Chernoviz, P.L.N., 1897. *Formulário e Guia Médico*, 16th ed. Livraria de A. Roger e F. Chernoviz, Pariz.
- Chernoviz, P.L.N., 1996 [1920]. *A Grande Farmacopeia Brasileira. Formulário e Guia Médico*, vol. 2., 19th ed. Itatiaia, Belo Horizonte. Rio de Janeiro.
- Choi, E.M., Hwang, J.K., 2003. *Investigations of anti-inflammatory and antinociceptive activities of Piper cubeba, Physalis angulata and Rosa hybrida*. J. Ethnopharmacol. 89, 171–175.
- Coelho, J.M., Antonioli, A.B., Nunes e Silva, D., Carvalho, T.M., Pontes, E.R., Odashiro, A.N., 2010. *Effects of silver sulfadiazine, ipê roxo (Tabebuia avellaneda) extract and barbatimão (Stryphnodendron adstringens) extract on cutaneous wound healing in rats*. Rev. Col. Bras. Cir. 37, 45–51.
- Cook, D., Lee, S.T., Taylor, C.M., Bassinner, B., Riet-Correa, F., Pfister, J.A., Gardner, D.R., 2014. *Detection of toxic monofluoroacetate in Palicourea species*. Toxicon 80, 9–16.
- Cosenza, G.P., Somavilla, N.S., Fagg, C.W., Brandão, M.G.L., 2013. *Bitter plants used as substitute of Cinchona spp. (quina) in Brazilian traditional medicine*. J. Ethnopharmacol. 149, 790–796.
- Costa, S.S., Souza, M.L., Ibrahim, T., Melo, G.O., Almeida, A.P., Guette, C., Férezou, J.P., Koatz, V.L., 2006. *Kalanchoine dimala, an anti-inflammatory salt from Kalanchoe brasiliensis*. J. Nat. Prod. 69, 815–818.
- Cravotto, G., Boffa, L., Genzini, L., Garella, D., 2010. *Phytotherapeutics: an evaluation of the potential of 1000 plants*. J. Clin. Pharm. Ther. 35, 11–48.
- Dal Belo, C.A., Lucho, A.P., Vinadé, L., Rocha, L., Seibert França, H., Marangoni, S., Rodrigues-Simioni, L., 2013. *In vitro antiophidian mechanisms of Hypericum brasiliense choisy standardized extract: quercetin-dependent neuroprotection*. Biomed. Res. Int., <http://dx.doi.org/10.1155/2013/943520>.
- Dash, G.K., Murthy, P.N., 2011. *Studies on wound healing activity of Heliotropium indicum Linn. leaves on rats*. ISRN Pharmacol., <http://dx.doi.org/10.5402/2011/847980>.

- Della Loggia, R., Tubaro, A., Redaelli, C., 1981. Evaluation of the activity on the mouse CNS of several plant extracts and a combination of them. *Rev. Neurol.* 51, 297–310.
- Devi Ramaiya, S., Bujang, J.S., Zakaria, M.H., King, W.S., Shaffiq Sahrir, M.A., 2013. Sugars, ascorbic acid, total phenolic content and total antioxidant activity in passion fruit (*Passiflora*) cultivars. *J. Sci. Food Agric.* 93, 1198–1205.
- Estevão, L.R., Mendonça, F.S., Baratella-Evêncio, L., Simões, R.S., Barros, M.E., Arantes, R.M., Rachid, M.A., Evêncio-Neto, J., 2013. Effects of aroeira (*Schinus terebinthifolius* Radlk) oil on cutaneous wound healing in rats. *Acta Cir. Bras.* 28, 202–209.
- Fagg, C.W., Lughadha, E.N., Milliken, W., Hind, D.N., Brandão, M.G.L., 2015. Useful Brazilian plants listed in the manuscripts and publications of the Scottish medic and naturalist George Gardner (1810–1849). *J. Ethnopharmacol.* 161, 18–29.
- Figueiredo, S.M., Nascimento, F.P., Freitas, C.S., Baggio, C.H., Soldi, C., Pizzolatti, M.G., Ibarrola Mdel, C., Arrua, R.L., Santos, A.R., 2011. Antinociceptive and gastro-protective actions of ethanolic extract from *Pluchea sagittalis* (Lam.) Cabrera. *J. Ethnopharmacol.* 135, 603–609.
- Fischer, D.C., Amorim Gualda, N.C., Bachiega, D., Carvalho, C.S., Lupo, F.N., Bonotto, S.V., Alves, M.O., Yogi, A., Santi, S.M., Avila, P.E., Kirchgatter, K., Moreno, P.R., 2004. In vitro screening for antiplasmodial activity of isoquinoline alkaloids from Brazilian plant species. *Acta Trop.* 92, 261–266.
- Fleischer, T.C., Ameade, E.P., Mensah, M.L., Sawyer, I.K., 2003. Antimicrobial activity of the leaves and seeds of *Bixa orellana*. *Fitoterapia* 74, 136–138.
- Forzza, R.C., Baumgratz, J.F., Bicudo, C.E.M., Canhos, D.A.L.C., Anibal, A.C., Marcus, A.N., Costa, A.F.C., Hopkins, D.P., Leitman, M.G., Lohmann, L.G., Lughadha, E.N., Maia, L.C.M., Menezes, G., Morim, M., Peixoto, M.P., Pirani, A.L., Prado, J.R., Queiroz, J., Souza, L.P., Souza, S., Castro, V., Stehmann, J.R., Sylvestre, L., Walter, S., Bruno, M.T., Zappi, D.C., 2012. New Brazilian floristic list highlights conservation challenges. *Bioscience* 62, 39–45.
- Franssen, F.F., Smeijsters, L.J., Berger, I., Medinilla Aldana, B.E., 1997. In vivo and in vitro antimalarial activities of some plants traditionally used in Guatemala against malaria. *Antimicrob. Agents Chemother.* 41, 1500–1503.
- Fratanni, F., Nazzaro, F., Marandino, A., Fusco, M.R., Coppola, R., Feo, V., Martino, L., 2013. Biochemical composition, antimicrobial activities, and anti-quorum-sensing activities of ethanol and ethyl acetate extracts from *Hypericum connatum* Lam. (*Guttiferae*). *J. Med. Food* 16, 454–459.
- Fritz, D., Venturi, C.R., Carginin, S., Schripsema, J., Roehe, P.M., Montanha, J.A., Poser, G.L., 2007. Herpes virus inhibitory substances from *Hypericum connatum* Lam., a plant used in southern Brazil to treat oral lesions. *J. Ethnopharmacol.* 113, 517–520.
- Gelmini, F., Beretta, G., Anselmi, C., Centini, M., Magni, P., Ruscica, M., Cavalchini, A., Maffei Facino, R., 2013. GC-MS profiling of the phytochemical constituents of the oleoresins from *Copaifera langsdorffii* Desf. and a preliminary in vivo evaluation of its antipsoriatic effect. *Int. J. Pharm.* 440, 170–178.
- Ghosh, S., Sarma, M., Patra, A., Hazra, B., 2010. Anti-inflammatory and anticancer compounds isolated from *Ventilago madraspatana* Gaertn., *Rubia cordifolia* Linn. and *Lantana camara* Linn. *J. Pharm. Pharmacol.* 62, 1158–1166.
- Giribabu, N., Rao, P.V., Kumar, K.P., Muniandy, S., Swapna Rekha, S., Salleh, N., 2014. Aqueous extract of *Phyllanthus niruri* leaves displays in vitro antioxidant activity and prevents the elevation of oxidative stress in the kidney of streptozotocin-induced diabetic male rats. *Evid. Based Complement. Alternat. Med.*, <http://dx.doi.org/10.1155/2014/834815>.
- Giulietti, A.M., Raymond, M.H., Queiroz, L.P., Wanderley, M.G.L., Berg, V., 2005. Biodiversity and conservation of plants in Brazil. *Conserv. Biol.* 19, 632–639.
- Gomes, P.B., Oliveira, M.M., Nogueira, C.R., Noronha, E.C., Carneiro, L.M., Bezerra, J.N., Neto, M.A., Vasconcelos, S.M., Fonteles, M.M., Viana, G.S., Sousa, F.C., 2005. Study of antinociceptive effect of isolated fractions from *Petiveria alliacea* L. (tipi) in mice. *Biol. Pharm. Bull.* 28, 42–46.
- Guenka, L.C., Gomes, R.C., Melo, V.L., Kitaniishi, C.R., Pereira, P.S., França, S.C., Couto, L.B., Beleboni, R.O., 2008. Anti-inflammatory and anti-nociceptive effects of *Zeyheria montana* (Bignoniacae) ethanol extract. *Mem. I. Oswaldo Cruz* 103, 768–772.
- Guimarães, M.R.C., 2005. Chernoviz e os manuais de medicina popular no Império. *Hist. Cienc. Saude Man.* 12, 501–514.
- Gupta, M., Mazumder, U.K., Chaudhuri, I., Chaudhuri, R.K., Bose, P., Bhattacharya, S., Manikandan, L., Patra, S., 2002. Antimicrobial activity of *Eupatorium ayapana*. *Fitoterapia* 73, 168–170.
- Heinrich, M., Kufer, J., Leonti, M., Pardo-de-Santayanna, M., 2006. Ethnobotany and ethnopharmacology – interdisciplinary links with the historical sciences. *J. Ethnopharmacol.* 107, 157–160.
- Henriques, M., Penido, C., 2014. The therapeutic properties of *Carapa guianensis*. *Curr. Pharm. Des.* 20, 850–856.
- Hiruma-Lima, C.A., Rodrigues, C.M., Kushima, H., Moraes, T.M., Lolis, S.F., Feitosa, S.B., Magri, L.P., Soares, F.R., Cola, M.M., Andrade, F.D., Vilegas, W., Souza Brito, A.R., 2009. The anti-ulcerogenic effects of *Curatella americana* L. *J. Ethnopharmacol.* 121, 425–432.
- Ibrahim, T., Cunha, J.M., Madi, K., Fonseca, L.M., Costa, S.S., Gonçalves Koatz, V.L., 2002. Immunomodulatory and anti-inflammatory effects of *Kalanchoe brasiliensis*. *Int. Immunopharmacol.* 2, 875–883.
- Ishida, K., Mello, J.C., Cortez, D.A., Filho, B.P., Ueda-Nakamura, T., Nakamura, C.V., 2006. Influence of tannins from *Stryphnodendron adstringens* on growth and virulence factors of *Candida albicans*. *J. Antimicrob. Chemother.* 58, 942–949.
- Iwamoto, L.H., Vendramini-Costa, D.B., Monteiro, P.A., Ruiz, A.L., Sousa, I.M., Foglio, M.A., Carvalho, J.E., Rodrigues, R.A., 2015. Anticancer and anti-inflammatory activities of a standardized dichloromethane extract from *Piper umbellatum* L. leaves. *Evid. Based. Complement. Alternat. Med.*, <http://dx.doi.org/10.1155/2015/948737>.
- Jang, M., Jeong, S.W., Cho, S.K., Ahn, K.S., Lee, J.H., Yang, D.C., Kim, J.C., 2014. Anti-inflammatory effects of an ethanolic extract of guava (*Psidium guajava* L.) leaves in vitro and in vivo. *J. Med. Food* 17, 678–685.
- Jegede, O.C., Ajanusi, J.O., Adaudi, A.O., Agbede, R.I., 2006. Anthelmintic efficacy of extracts of *Spigelia anthelmia* Linn on experimental *Nippostrongylus brasiliensis* in rats. *Jpn. J. Vet. Sci.* 7, 229–232.
- Kamel, E.G., El-Emam, M.A., Mahmoud, S.S., Fouada, F.M., Bayaumy, F.E., 2011. Parasitological and biochemical parameters in *Schistosoma mansoni*-infected mice treated with methanol extract from the plants *Chenopodium ambrosioides*, *Conyza dioscoridis* and *Sesbania sesban*. *Parasitol. Int.* 60, 388–392.
- Kouam, M.K., Payne, V.K., Miégué, E., Tendonkeng, F., Lemoufouet, J., Kana, J.R., Boukila, B., Pamo, E.T., Mnm, B., 2015. Evaluation of in vivo acaricidal effect of soap containing essential oil of *Chenopodium ambrosioides* leaves on *Rhipicephalus lunulatus* in the western highland of Cameroon. *J. Pathog.*, <http://dx.doi.org/10.1155/2015/516869>.
- Kwieciński, M.R., Felipe, K.B., Correia, J.F., Ferreira, E.A., Rossi, M.H., Moura Gatti, F., Filho, D.W., Pedrosa, R.C., 2011. Brazilian *Bidens pilosa* Linné yields fraction containing quercetin-derived flavonoid with free radical scavenger activity and hepatoprotective effects. *Libyan J. Med.* 6, 1–8.
- Leitão, F., Leitão, S.G., Almeida, M.Z., Cantos, J., Coelho, T., Silva, P.E., 2013. Medicinal plants from open-air markets in the State of Rio de Janeiro, Brazil as a potential source of new antimycobacterial agents. *J. Ethnopharmacol.* 149, 513–521.
- Lopes-Martins, R.A., Pegoraro, D.H., Woisky, R., Penna, S.C., Sertié, J.A., 2002. The anti-inflammatory and analgesic effects of a crude extract of *Petiveria alliacea* L. (Phytolaccaceae). *Phytomedicine* 9, 245–248.
- Luiz, R.L., Vila, T.V., Mello, J.C., Nakamura, C.V., Rozental, S., Ishida, K., 2015. Proanthocyanidins polymeric tannin from *Stryphnodendron adstringens* are active against *Candida albicans* biofilms. *BMC Complement. Altern. Med.*, <http://dx.doi.org/10.1186/s12906-015-0597-4>.
- Magrini, F.E., Specht, A., Gaio, J., Girelli, C.P., Migues, I., Heinzen, H., Sartori, V.C., Cesio, V., 2014. Viability of *Cabralia canjerana* extracts to control the South American fruit fly, *Anastrepha fraterculus*. *J. Insect Sci.*, <http://dx.doi.org/10.1093/jis/14.1.47>.
- Maranhão, H.M., Vasconcelos, C.F., Rolim, L.A., Neto, P.J., Neto, J.C., Filho, R.C., Fernandes, M.P., Costa-Silva, J.H., Araújo, A.V., Wanderley, A.C., 2014. Hepatoprotective effect of the aqueous extract of *Simarouba amara* Aublet (Simaroubaceae) stem bark against carbon tetrachloride (CCl₄)-induced hepatic damage in rats. *Molecules* 19, 17735–17746.
- Martius, C.F.P., 1843. *Systema Materiae Medicinae Vegetabilis Brasilensis, Lipsiae*.
- Mbeunkui, F., Grace, M.H., Lategan, C., Smith, P.J., Raskin, I., Lila, M.A., 2012. In vitro antiparasitic activity of indole alkaloids from the stem bark of *Geissospermum velosii*. *J. Ethnopharmacol.* 139, 471–477.
- Meccia, G., Quintero, P., Rojas, L.B., Usobilaga, A., Velasco, J., Diaz, T., Diaz, C., Velásquez, J., Toro, M., 2013. Chemical composition of the essential oil from the leaves of *Carapa guianensis* collected from Venezuelan Guayana and the antimicrobial activity of the oil and crude extracts. *Nat. Prod. Commun.* 8, 1641–1642.
- Medeiros, M.F.T., 2008. Historical Ethnobotany: an approach through historical documents and their implications now days. In: Albuquerque, U.P., Hanzaki, N. (Eds.), Recent Development and Case Studies in Ethnobotany. Brazilian Society of Ethnobiology and Ethnoecology, Recife, pp. 127–142.
- Mesia-Vela, S., Santos, M.T., Soucar, C., Lima-Landman, M.T., Lapa, A.J., 2002. *Solanum paniculatum* L. (Juruábeba): potent inhibitor of gastric acid secretion in mice. *Phytomedicine* 9, 508–514.
- Micali, S., Sighinolfi, M.C., Celia, A., Stefani, S., Grande, M., Cicero, A.F., Bianchi, G., 2006. Can *Phyllanthus niruri* affect the efficacy of extracorporeal shock wave lithotripsy for renal stones? A randomized, prospective, long-term study. *J. Urol.* 176, 1020–1022.
- Michelin, D.C., Salgado, H.R.N., 2004. Evaluation of the laxative activity of *Operculina macrocarpa* L. Urban (Convolvulaceae). *Rev. Bras. Farmacogn.* 14, 105–109.
- Miot, H.A., Batistella, R.F., Batista, K.A., Volpato, D.E., Augusto, L.S., Madeira, N.G., Haddad Jr., V., Miot, L.D., 2004. Comparative study of the topical effectiveness of the andiroba oil (*Carapa guianensis*) and DEET 50% as repellent for *Aedes sp.* *Rev. I. Med. Trop.* 46, 253–256.
- Miranda Júnior, R.N., Dolabela, M.F., Silva, M.N., Póvoa, M.M., Maia, J.G., 2012. Antiplasmodial activity of the andiroba (*Carapa guianensis* Aubl., Meliaceae) oil and its limonoid-rich fraction. *J. Ethnopharmacol.* 142, 679–683.
- Morey, A.T., Souza, F.C., Santos, J.P., Pereira, C.A., Cardoso, J.D., Almeida, R.S., Costa, M.A., Mello, J.C., Nakamura, C.V., Pingue-Filho, P., Yamauchi, L.M., Yamada-Ogata, S.F., 2015. Antifungal activity of condensed tannins from *Stryphnodendron adstringens*: effect on *Candida tropicalis* growth and adhesion properties. *Curr. Pharm. Biotechnol.* 17, 365–375.
- Nayak, B.S., Kanhai, J., Milne, D.M., Swanston, W.H., Mayers, S., Eversley, M., Rao, A.V., 2010. Investigation of the wound healing activity of *Carapa guianensis* L. (Meliaceae) bark extract in rats using excision, incision, and dead space wound models. *J. Med. Food* 13, 1141–1146.
- Nayak, B.S., Kanhai, J., Milne, D.M., Pinto Pereira, L., Swanston, W.H., 2011. Experimental evaluation of ethanolic extract of *Carapa guianensis* L. leaf for its wound healing activity using three wound models. *Evid. Based. Complement. Alternat. Med.*, <http://dx.doi.org/10.1093/ecam/nep160>.
- Neves, E.G., 2006. *Arqueologia da Amazônia*. Jorge Zahar Editor, Rio de Janeiro.
- Nishioka, S.E.A., Escalante, R.D., 1997. Poisoning by the ingestion of seeds of the fruit of the "cotieira" (*Joannesia princeps*). *São Paulo Med. J.* 115, 1366–1367.
- Nogueira, N.P., Reis, P.A., Laranja, G.A., Pinto, A.C., Aiub, C.A., Felzenszwalb, I., Paes, M.C., Bastos, F.F., Bastos, V.L., Sabino, K.C., Coelho, M.G., 2011. In vitro and in vivo

- toxicological evaluation of extract and fractions from *Baccharis trimera* with anti-inflammatory activity.** *J. Ethnopharmacol.* 138, 513–522.
- Noldin, V.F., Oliveira Martins, D.T., Marcello, C.M., Silva Lima, J.C., Delle Monache, F., Cechinel Filho, V., 2005. Phytochemical and antiulcerogenic properties of rhizomes from *Simaba ferruginea* St. Hill. (Simaroubaceae). *Z. Naturforsch.* 60, 701–706.
- Ojewole, J.A., Awe, E.O., Chiwororo, W.D., 2008. Antidiarrhoeal activity of *Psidium guajava* Linn. (Myrtaceae) leaf aqueous extract in rodents. *J. Smooth Muscle Res.* 44, 195–207.
- Olajide, O.A., Aderogba, M.A., Adedapo, A.D., Makinde, J.M., 2004. Effects of *Anacardium occidentale* stem bark extract on in vivo inflammatory models. *J. Ethnopharmacol.* 95, 139–142.
- Oliveira Campos, M.P., Riechelmann, R., Martins, L.C., Hassan, B.J., Casa, F.B., Del Giglio, A., 2011. Guarana (*Paullinia cupana*) improves fatigue in breast cancer patients undergoing systemic chemotherapy. *J. Altern. Complement. Med.* 17, 505–512.
- Oliveira, D.R., Krettli, A.U., Aguiar, A.C., Leitão, G.G., Vieira, M.N., Martins, K.S., Leitão, S.G., 2015. Ethnopharmacological evaluation of medicinal plants used against malaria by Quilombola Communities from Oriximiná, Brazil. *J. Ethnopharmacol.* 173, 424–434.
- Paganotte, D.M., Sannomiya, M., Rinaldo, D., Vilegas, W., Salgado, H.R.N., 2016. *Operculina macrocarpa*: chemical and intestinal motility effect in mice. *Rev. Bras. Farmacogn.* 26, 427–432.
- Paiva, L.A., Alencar Cunha, K.M., Santos, F.A., Gramosa, N.V., Silveira, E.R., Rao, V.S., 2002. Investigation on the wound healing activity of oleo-resin from *Copaifera langsdorffii* in rats. *Phytother. Res.* 16, 737–739.
- Paul, E.L., Lunardelli, A., Caberlon, E., Oliveira, C.B., Santos, R.C., Biolchi, V., Bastos, C.M., Moreira, K.B., Nunes, F.B., Gosmann, G., Oliveira, J.R., 2009. Anti-inflammatory and immunomodulatory effects of *Baccharis trimera* aqueous extract on induced pleurisy in rats and lymphoproliferation in vitro. *Inflammation* 32, 419–425.
- Pawar, S.P., Sathwane, P.N., Metkar, B.R., Pal, S.C., Kasture, V.S., Kasture, S.B., 2004. Anti-inflammatory and analgesic activity of *Anacardium Occidentale* leaf extracts. *Anc. Sci. Life.* 19, 169–173.
- Pereira, N.A., Jaccoud, R.J.S., Mors, W.B., 1996. Triaga Brasiliaca: renewed interest in a seventeenth-century panacea. *Toxicon* 34, 511–516.
- Pereira, T.B., Rocha e Silva, L.F., Amorim, R.C., Melo, M.R., Zaccardi de Souza, R.C., Eberlin, M.N., Lima, E.S., Vasconcellos, M.C., Pohlit, A.M., 2014. In vitro and in vivo anti-malarial activity of limonoids isolated from the residual seed biomass from *Carapa guianensis* (andiroba) oil production. *Malaria J.* 13, 317.
- Pinto, S.C., Bueno, F.G., Panizzi, G.P., Morais, G., Santos, P.V., Baesso, M.L., Leite-Mello, E.V., Mello, J.C., 2013. *Stryphnodendron adstringens*: clarifying wound healing in Streptozotocin-Induced diabetic rats. *Planta Med.* 81, 1090–1096.
- Pisonis, G., 1648. Historiae Naturalis & Medicinae. De Arboribus, Fruticibus & Herbis Medicinalibus, ac Alimentois in Brasilia. Elzevier, Amsterdam.
- Raupp, I.M., Sereniki, A., Virtuoso, S., Ghislandi, C., Cavalcanti e Silva, E.L., Trebien, H.A., Miguel, O.G., Andreatini, R., 2008. Anxiolytic-like effect of chronic treatment with *Erythrina velutina* extract in the elevated plus-maze test. *J. Ethnopharmacol.* 118, 295–299.
- Reddy, J.S., Rao, P.R., Reddy, M.S., 2002. Wound healing effects of *Heliotropium indicum*, *Plumbago zeylanicum* and *Acalypha indica* in rats. *J. Ethnopharmacol.* 79, 249–251.
- Reis, G.O., Vicente, G., Carvalho, F.K., Heller, M., Micke, G.A., Pizzolatti, M.G., Fröde, T.S., 2014. *Croton antisyphiliticus* Mart. attenuates the inflammatory response to carageenan-induced pleurisy in mice. *Inflammopharmacology* 22, 115–126.
- Rivera-Arce, E., Chávez-Soto, M.A., Herrera-Arellano, A., Arzate, S., Agüero, J., Feria-Romero, I.A., Cruz-Guzmán, A., Lozoya, X., 2007. Therapeutic effectiveness of a *Mimosa tenuiflora* cortex extract in venous leg ulceration treatment. *J. Ethnopharmacol.* 109, 523–528.
- Rocha e Silva, L.F., Nogueira, K.L., Pinto, A.C., Katzin, A.M., Sussmann, R.A., Muniz, M.P., Andrade Neto, V.F., Chaves, F.C., Coutinho, J.P., Lima, E.S., Krettli, A.U., Tadei, W.P., Pohlit, A.M., 2015. In vivo antimalarial activity and mechanisms of action of 4-nerolidylcatechol derivatives. *Antimicrob. Agents Chemother.* 59, 3271–3280.
- Roersch, C.M., 2010. *Piper umbellatum* L.: a comparative cross-cultural analysis of its medicinal uses and an ethnopharmacological evaluation. *J. Ethnopharmacol.* 131, 522–537.
- Santos, A.O., Ueda-Nakamura, T., Dias Filho, B.P., Veiga Junior, V.F., Pinto, A.C., Nakamura, C.V., 2008. Antimicrobial activity of Brazilian copaiba oils obtained from different species of the *Copaifera* genus. *Mem. I. Oswaldo Cruz* 103, 277–281.
- Santos, O.J., Malafaia, O., Ribas-Filho, J.M., Czeczk, N.G., Santos, R.H., Santos, R.A., 2013. Influence of *Schinus terebinthifolius* Radji (aroeira) and *Carapa guianensis* Aublet (andiroba) in the healing process of gastrorrhaphies. *Arq. Bras. Cir. Dig.* 26, 84–91.
- Santos, J.L., Dantas, R.E., Lima, C.A., Araújo, S.S., Almeida, E.C., Marçal, A.C., Estevam, C.S., 2014. Protective effect of a hydroethanolic extract from *Bowdichia virgilioides* on muscular damage and oxidative stress caused by strenuous resistance training in rats. *J. Int. Soc. Sports Nutr.* 11, 58.
- Santos, E.C., Bicca, M.A., Blum-Silva, C.H., Costa, A.P., Dos Santos, A.A., Schenkel, E.P., Farina, M., Reginatto, F.H., de Lima, T.C., 2015. Anxiolytic-like, stimulant and neuroprotective effects of *Ilex paraguariensis* extracts in mice. *Neuroscience* 292, 13–21.
- Santos Rosa, D., Faggion, S.A., Gavin, A.S., Anderson de Souza, M., Fachim, H.A., Ferreira dos Santos, W., Soares Pereira, A.M., Cunha, A.O., Beleboni, R.O., 2012. Erysothrine, an alkaloid extracted from flowers of *Erythrina mulungu* Mart. ex Benth: evaluating its anticonvulsant and anxiolytic potential. *Epilepsy Behav.* 23, 205–212.
- Sarria, A.L., Soares, M.S., Matos, A.P., Fernandes, J.B., Vieira, P.C., Silva, M.F., 2011. Effect of triterpenoids and limonoids isolated from *Cabralea canjerana* and *Carapa guianensis* (Meliaceae) against *Spodoptera frugiperda* (J. E. Smith). *Z. Naturforsch.* 66, 245–250.
- Seito, L.N., Ruiz, A.L., Vendramini-Costa, D., Tinti, S.V., Carvalho, J.E., Bastos, J.K., Di Stasi, L.C., 2011. Antiproliferative activity of three methoxylated flavonoids isolated from *Zeyheria montana* Mart. (Bignoniaceae) leaves. *Phytother. Res.* 25, 1447–1450.
- Singh, B.R., Singh, V., Ebibeni, N., Singh, R.K., 2013. Antimicrobial and herbal drug resistance in enteric bacteria isolated from faecal droppings of common house lizard/Gecko (*Hemidactylus frenatus*). *Int. J. Microbiol.*, <http://dx.doi.org/10.1155/2013/340848>.
- Siskind, D., Robert, C., Crawford, S., 2014. *Cestrum parqui* and psychosis. *Australas. Psychiatry* 22, 305–306.
- Sosa, S., Balick, M.J., Arvigo, R., Esposito, R.G., Pizza, C., Altinier, G., Tubaro, A., 2002. Screening of the topical anti-inflammatory activity of some Central American plants. *J. Ethnopharmacol.* 81, 211–215.
- Souza-Almeida, E.S., Filho, V.C., Niero, R., Clasen, B.K., Balogun, S.O., Oliveira Martins, D.T., 2011. Pharmacological mechanisms underlying the anti-ulcer activity of methanol extract and canthin-6-one of *Simaba ferruginea* A. St-Hil. in animal models. *J. Ethnopharmacol.* 134, 630–636.
- Souza, A.B., Martins, C.H., Souza, M.G., Furtado, N.A., Heleno, V.C., de Sousa, J.P., Rocha, E.M., Bastos, J.K., Cunha, W.R., Veneziani, R.C., Ambrósio, S.R., 2011. Antimicrobial activity of terpenoids from *Copaifera langsdorffii* Desf. against cariogenic bacteria. *Phytother. Res.* 25, 215–220.
- Suzigan, M.I., Battachio, A.P., Coelho, K.L., Coelho, C.A., 2009. An aqueous extract of *Bidens pilosa* L. protects liver from cholestatic disease: experimental study in young rats. *Acta Cir. Bras.* 24, 347–352.
- Thomazzi, S.M., Silva, C.B., Silveira, D.C., Vasconcellos, C.L., Lira, A.F., Cambui, E.V., Estevam, C.S., Antoniolli, A.R., 2010. Antinociceptive and anti-inflammatory activities of *Bowdichia virgilioides* (sucupira). *J. Ethnopharmacol.* 127, 451–456.
- Truiti, M.C., Sarragiotto, M.H., Abreu Filho, B.A., Nakamura, C.V., Dias Filho, B.P., 2003. In vitro antibacterial activity of a 7-O-beta-D-glucopyranosyl-nutanolcoumarin from *Chaptalia nutans* (Asteraceae). *Mem. I. Oswaldo Cruz* 98, 283–286.
- Tsai, J.C., Peng, W.H., Chiu, T.H., Lai, S.C., Lee, C.Y., 2011. Anti-inflammatory effects of *Scoparia dulcis* L. and betulinic acid. *Am. J. Chin. Med.* 39, 943–956.
- Vargas, F.S., Almeida, D.O.P., Aranha, E.S., Boleti, A.P.A., Newton, P., Vasconcellos, M.C., Junior, V.F., Lima, E.S., 2015. Biological activities and cytotoxicity of diterpenes from *Copaifera* spp. Oleoresins. *Molecules* 20, 6194–6210.
- Villalba, M.A., Carmo, M.I., Leite, M.N., Sousa, O.V., 2007. Atividades farmacológicas dos extratos de *Zanthoxylum chiloperone* (Rutaceae). *Rev. Bras. Farmacogn.* 17, 236–241.
- Vilela, F.C., Bitencourt, A.D., Cabral, L.D., Franqui, L.S., Soncini, R., Giusti-Paiva, A., 2010. Anti-inflammatory and antipyretic effects of *Sonchus*. *J. Ethnopharmacol.* 127, 737–741.
- Weniger, B., Robledo, S., Arango, G.J., Deharo, E., Aragón, R., Muñoz, V., Callapa, J., Lobstein, A., Anton, R., 2001. Antiprotozoal activities of Colombian plants. *J. Ethnopharmacol.* 78, 193–200.
- WHO, 2002. Traditional Medicine Strategy 2002–2005. World Health Organization, Geneva.
- Yamaguchi, M.U., Garcia, F.P., Cortez, D.A., Ueda-Nakamura, T., Filho, B.P., Nakamura, C.V., 2011. Antifungal effects of ellagittannin isolated from leaves of *Ocotea odorifera* (Lauraceae). *Antonie Van Leeuwenhoek* 99, 507–514.
- Yong, Y.K., Zakaria, Z.A., Kadir, A.A., Somchit, M.N., Ee Cheng Lian, G., Ahmad, Z., 2013. Chemical constituents and antihistamine activity of *Bixa orellana* leaf extract. *BMC Complement. Altern. Med.*, <http://dx.doi.org/10.1186/1472-6882-13-32>.
- Zippel, J., Deters, A., Hensel, A., 2009. Arabinogalactans from *Mimosa tenuiflora* (Willd.) Poiret bark as active principles for wound-healing properties: specific enhancement of dermal fibroblast activity and minor influence on HaCaT keratinocytes. *J. Ethnopharmacol.* 124, 391–396.