

# INTERRELATIONSHIP AND ENSO RELATIONSHIP OF FORTALEZA RAINFALL IN DIFFERENT SEASONS

Rajaram P. Kane

Received March 10, 2000 / Accepted December 25, 2001

The rainfall series at Fortaleza, Ceara, NE Brazil for the four seasons DJF, MAM (main rainy season), JJA, SON were poorly intercorrelated (correlation 0.45 or less, common variance only ~20%). The relationship with El Niños was poor, with only about half of the El Niños associated with droughts. Among the El Niños, Unambiguous ENSOW had a better affinity for droughts, more so for DJF. But some droughts occurred in the absence of El Niños and even during La Niña years, and some excess rains occurred during El Niño years. A large interference from other factors (mostly Atlantic SST etc.) unrelated to the ENSO phenomenon is obvious. The media seems to be giving undue importance to El Niño and La Niña only, which needs to be diluted.

**Key words:** El Niño; Fortaleza; Rainfall.

**INTERCORRELAÇÕES E RELAÇÕES COM A ENSO DA PRECIPITAÇÃO EM FORTALEZA EM DIFERENTES ESTAÇÕES** - As séries de precipitação em Fortaleza, Ceará, nordeste do Brasil para as quatro estações DJF, MAM (principal estação chuvosa), JJA, SON estão pobremente intercorrelacionadas (correlação de 0,45 ou menos, variância comum de apenas ~20%). A relação com os El Niños foi pobre, com apenas cerca da metade dos El Niños associados com secas. Entre os El Niños, ENSOs não ambíguos têm melhor afinidade para secas, mais para DJF. Mas algumas secas ocorreram na ausência de El Niños e mesmo durante anos de La Niña, e algum excesso de chuvas ocorreu durante anos de El Niño. Uma larga interferência de outros fatores (principalmente SST Atlântico, etc.) não correlacionados ao fenômeno ENSO é óbvia. Parece que a mídia está dando importância indevida ao El Niño e à La Niña apenas, que precisam ser diluídos.

**Palavras-chave:** El Niño; Fortaleza; Precipitação.

Instituto Nacional de Pesquisas Espaciais - INPE  
Caixa Postal 515, 12201-970 - São José dos Campos, SP, Brazil  
kane@laser.inpe.br

## INTRODUCTION

The main rainy season at Fortaleza, Ceara in NE Brazil is MAM (March, April, May). Fig. 1 (a) shows the average rainfalls (climatology) in various months, while Fig. 1 (b) shows the seasonal details for seasons MAM, DJF (December, January, February), JJA (June, July, August), SON (September, October, November), which are also given in Table 1.

The percentage standard deviations are 38% for MAM, 61% for DJF, 68% for JJA, and 72% for SON, indicating that smaller average rainfalls have larger percentage deviations, similar to the larger percentage deviations for geographical locations with smaller rainfalls and vice versa (Kousky and Chu, 1978; Kousky, 1979). All these rainfalls have large year-to-year fluctuations, some probably related to El Niño events. The purpose of the present paper is to




Figure 1 - a) Average monthly variation of Fortaleza rainfall for 1849-1999; b) Characteristics of the 4 seasons DJF, MAM, JJA, SON.

Season	Average rainfall		Standard deviation	
	mm	Percentage of annual rainfall	mm	Percentage
DJF	314	22	± 191	± 61
MAM	879	61	± 337	± 38
JJA	206	14	± 140	± 68
SON	47	3	± 34	± 72
Annual	1446	100	± 519	± 36

Table 1 - Characteristics of the Fortaleza rainfall, average for 1849-1999.

check whether all seasonal rainfalls vary in a similar way and have similar El Niño associations.

**DATA**

Rainfall data for Fortaleza (3° 42' S, 38° 31' W) Ceara, north-east Brazil were obtained from FUNCEME (Fundação Cearense de Meteorologia e Recursos Hídricos). El Niño years were obtained from the list in Quinn et al. (1987). SST (sea surface temperature) anomaly data for Puerto Chicama (Peruvian coast, 8° S, 80° W) were obtained from

Deser and Wallace (1987) and from private communications from Dr. Todd Mitchell and Dr. Don Garrett (also available at [http://tao.atmos.washington.edu/data\\_sets/chicama\\_sst/](http://tao.atmos.washington.edu/data_sets/chicama_sst/)). Data for SST anomalies for the Pacific regions, Nino 1+2 (0° -10° S, 90° W-80° W), Nino 3 (5° N-5° S, 150° W-90° W), Nino 3.4 (5° N-5° S, 170° W-120° W) and Nino 4 (5° N-5° S, 160° E-150° W) were obtained from the Web site of the Climate Prediction Center (CPC) of the National Oceanic and Atmospheric Administration (NOAA), Washington D. C. (<http://www.cpc.ncep.noaa.gov/data/indices>).

**PROCESSING OF RAINFALL DATA AT FORTALEZA: INTERCORRELATIONS**

All data of any particular season were expressed in normalized units (deviations from mean of the series of that season, divided by the standard deviation of the series of that season). It was noticed that the deviations were generally not similar for all the seasons. An intercorrelation analysis gave results shown in Table 2. Since Puerto Chicama SST anomalies were available only from 1925 onwards, the correlations were obtained separately for 1849-1924 (Table 2 a) and for 1925-1999 (Table 2b). As can be seen, the correlations are low, the maximum being  $0.45 \pm 0.09$  between MAM and JJA during 1849-1924, explaining only ~20% (square of 0.45) of variance. Thus, the rainfalls in the various seasons have anomalies almost independent of each other. In their prediction schemes for NE Brazil seasonal (March-June) rainfall, Hastenrath and Greischar (1993) used five potential predictors, one of which was the pre-season October-January rainfall, for which the training period (1921-1957) correlation was +0.55, significant at a 1% level. In the case of Fortaleza, the value of the correlation is lower, reducing its prediction potential.

**SIMILARITIES AND DISSIMILARITIES OF THE RAINFALL EXTREMES**

Extremes of the rainfall anomalies (magnitudes exceeding  $1.0 \sigma$ ) in DJF and/or the succeeding MAM fell into four categories : Similar variations, either (i) DJF and MAM positive or (ii) DJF and MAM negative, and dissimilar variations, (iii) DJF positive, MAM negative or (iv) DJF negative, MAM positive. Table 3 gives the years and rainfall anomaly values in these four categories, for DJF, MAM and JJA seasons. The rainfalls in SON are very small and are not considered here.

There are many more events in category (a) when DJF and MAM were similar and these would yield a positive correlation, but the events in category (b) when DJF and MAM are dissimilar, would reduce the correlation. Hence the overall value of the correlation was only  $\sim +0.45$  or less. The JJA values were sometimes like MAM and sometimes like DJF, but in events of category (ii) when DJF and MAM were both negative, JJA also was mostly negative, indicating long-lasting droughts in these years. Some of these are El Niño years (shown bold), but El Niño years appear in other categories also (i), (iii), (iv). Thus, El Niño effects are complicated, as discussed further.

**RELATIONSHIP WITH EL NIÑO EVENTS**

**(A) Relationship with a finer classification of El Niños**

El Niños are popularly believed to be associated with droughts in many parts of the world, notably Indonesia, India and northeast Brazil. El Niños (positive temperature anomalies) generally start near the Peru-Ecuador coast (for example, at Puerto Chicama, 8° S, 80° W) and spread westward in the Pacific within a month or two. During the El Niño years, the Southern Oscillation Index SOI (represented by Tahiti minus Darwin atmospheric pressure difference T-D) has a minimum. However, not all El Niños seem to be effective. Recently, Kane (1997a,b; 1998 a,b,c; 1999 a,b) attempted a finer classification in which *Unambiguous ENSOW* type events were found to be overwhelmingly associated with droughts in India, southeastern Australia and some other regions. These were El Niño (EN) years (Quinn et al., 1987 list), during which the *12-month running means* of the Southern Oscillation Index SOI (represented by Tahiti minus Darwin atmospheric pressure difference T-D) had a minimum (SO) and the equatorial eastern Pacific sea surface temperatures

(a) 1849-1924	DJF	MAM	JJA	SON	(b) 1925-1999	DJF	MAM	JJA	SON
DJF	1.00				DJF	1.00			
MAM	0.40	1.00			MAM	0.31	1.00		
JJA	0.15	0.45	1.00		JJA	0.25	0.24	1.00	
SON	0.13	0.24	0.17	1.00	SON	0.35	0.31	0.25	1.00

**Table 2** - Correlations between the normalized rainfall deviations in different seasons for (a) 1849-1924 and (b) 1925-1999.

(a) DJF, MAM similar

(i) DJF, MAM positive			(ii) DJF, MAM negative				
Year	DJF	MAM	JJA	Year	DJF	MAM	JJA
1860	0.13	3.27	0.55	<b>1877</b>	-1.43	-1.92	-0.25
1872	2.24	0.66	1.58	<b>1878</b>	-1.03	-1.56	-1.27
<b>1873</b>	1.92	0.77	-0.33	1879	-1.06	-1.48	-1.20
1894	2.24	2.53	0.59	1898	-1.24	-1.50	-1.20
<b>1899</b>	2.21	2.16	1.47	<b>1900</b>	-0.98	-2.37	-0.31
1910	1.60	1.10	-0.73	<b>1951</b>	-1.49	-1.34	0.03
<b>1912</b>	1.61	2.13	1.26	<b>1958</b>	-1.29	-1.80	-0.27
<b>1917</b>	2.77	0.21	-0.20				
1921	1.71	0.86	-0.12				
<b>1939</b>	1.64	0.30	-0.34				
1964	3.60	1.11	0.41				
1974	1.00	1.88	0.49				
1986	1.90	0.95	0.79				
<b>1995</b>	0.52	1.76	0.24				

(b) DJF, MAM dissimilar

(iii) DJF negative, MAM positive			(iv) DJF positive, MAM negative				
Year	DJF	MAM	JJA	Year	DJF	MAM	JJA
1866	-0.82	3.14	0.60	1893	2.25	-0.54	-0.90
1895	-0.58	1.81	2.74	<b>1932</b>	0.14	-1.64	-0.19
<b>1896</b>	-0.90	1.56	1.09	1935	1.86	-0.01	-0.30
				1961	1.80	-0.18	-0.12
				1980	1.64	-1.76	-0.38

**Table 3** - Years when rainfall anomalies in DJF and/or MAM seasons were extremes (both DJF and MAM exceeding  $1.0 \sigma$ , or any one exceeding  $1.5\sigma$ ) and **a**) when DJF and MAM were similar and **b**) when DJF and MAM were dissimilar. Years shown bold are El Niño years.

SST had a maximum (*W*) in the middle of the calendar year. If the maxima were not in the middle of the calendar year but in the beginning or end, the events were called *Ambiguous ENSOW*. El Niño events not falling into these two categories were called *Other EN*. However, there were years without El Niños (i.e., not in the Quinn et al., 1987 list) and these were of the types (i) only SO or only W, (ii) SOW, (iii) C (colder SST, anti-El Niño, also called La Niña), (iv) SOC (SO in the earlier part of the year, followed by C in later part) and (v) Non-events. Table 4 shows the rainfall status of the DJF, MAM, JJA, SON seasons at Fortaleza for El Niño years of the types (a) Unambiguous ENSOW, (b) Ambiguous ENSOW, (c) Other El Niños, (d) All El Niños. The All India summer monsoon rainfall status (IMR) is also given,

for comparison. Instead of giving the actual rainfall normalised deviations, data are presented by 6 symbols representing broad groups. Thus, +, f, F represent positive deviations in the ranges 0 to  $0.5 \sigma$ ;  $0.5$  to  $1.0 \sigma$  and, exceeding  $1.0 \sigma$  respectively, while -, d, D represent negative deviations in the ranges 0 to  $-0.5 \sigma$ ,  $-0.5$  to  $-1.0 \sigma$  and, exceeding  $-1.0 \sigma$  respectively. The symbols S (strong), M (Moderate), W (Weak) indicate the strength of the El Niño as given by Quinn et al. (1987), while I and II indicate the first and second years of double events (El Niños in two consecutive years, 1957-1958 etc.).

- (1) As can be seen in Table 4 (a) for the 16 events of type Unambiguous ENSOW, all had a negative deviation for Indian rainfall (IMR), indicating

(a) Unambiguous ENSOW		Fortaleza		Fortaleza		Fortaleza		Fortaleza	
Year	IMR	DJF	MAM	JJA	SON				
S 1877 I	D	D	D	-	d				
M 1888 I	d	d	D	-	D				
M 1895 I	-	d	F	F	f				
S 1899 I	D	F	F	F	d				
M 1902	d	d	D	-	-				
M 1905	D	D	+	d	F				
S 1911 I	D	f	-	f	d				
S 1918 I	D	+	f	-	f				
M 1930 I	d	-	d	-	d				
S 1941 II	D	-	d	d	D				
M 1951	D	D	D	+	F				
S 1957 I	d	d	+	D	+				
M 1965	D	d	f	F	d				
S 1972 I	D	d	d	F	d				
S 1982 I	D	-	d	-	f				
M 1987	D	-	d	f	d				
16 events									
F		0	1	2	4	2			
f		0	1	2	2	3			
+		0	1	2	1	1			
-		1	4	1	6	1			
d		4	6	5	2	7			
D		11	3	4	1	2			
(F,f,+)(-,d,D)	0/16	3/13	6/10	7/9	6/10				
Ratio (Negative/Total)	1.00	0.81	0.62	0.56	0.62				

(c) Other EN		Fortaleza		Fortaleza		Fortaleza		Fortaleza	
Year	IMR	DJF	MAM	JJA	SON				
ENSO									
1871	-	D	-	f	f				
1873	D	F	f	-	F				
1880	-	D	f	+	+				
1891	d	d	d	d	-				
1900	+	d	D	-	D				
1912	d	F	F	F	+				
ENW									
1884	f	D	-	d	+				
EN									
1897	+	f	f	F	f				
1929	-	-	d	-	d				
1932	d	+	D	-	-				
1939	d	F	+	-	F				
1943	+	D	-	d	-				
ENC									
1874	F	+	-	f	-				
1887	f	-	+	D	D				
1889	f	d	D	d	-				
1907	d	d	D	-	-				
1917	F	F	+	-	f				
1973	f	F	f	F	f				
18 events									
F		2	5	1	3	2			
f		4	1	4	2	4			
+		3	2	3	1	3			
-		3	2	4	7	6			
d		5	4	2	4	1			
D		1	4	4	1	2			
(F,f,+)(-,d,D)	9/9	8/10	8/10	6/12	9/9				
Ratio (Negative/Total)	0.5	0.56	0.56	0.67	0.5				

(b) Ambiguous ENSOW		Fortaleza		Fortaleza		Fortaleza		Fortaleza	
Year	IMR	DJF	MAM	JJA	SON				
S 1878 II	F	D	D	D	D				
M 1914	f	f	+	F	d				
M 1919 II	+	d	D	d	F				
M 1923	-	+	+	d	d				
S 1925 I	d	+	-	d	+				
S 1926 II	f	+	+	d	d				
M 1931 II	+	f	d	d	d				
S 1940 I	-	-	+	+	-				
W 1948	+	-	+	+	-				
M 1953	f	d	-	d	-				
S 1958 II	+	D	D	-	d				
W 1963	+	f	F	D	-				
W 1969	-	-	f	F	-				
M 1976	+	f	-	d	-				
S 1983 II	F	-	D	d	d				
M 1992 I	d	+	d	-	d				
M 1993 II	+	d	d	f	-				
S 1997 I	+	D	+	D	-				
S 1998 II	+	-	d	d	d				
19 events									
F		2	0	1	2	1			
f		3	4	1	1	0			
+		9	4	6	2	1			
-		3	5	3	2	8			
d		2	3	4	9	8			
D		0	3	4	3	1			
(F,f,+)(-,d,D)	14/5	8/11	8/11	5/14	2/17				
Ratio (Negative/Total)	0.26	0.58	0.58	0.74	0.89				

(d) All 53 events									
F		4	6	4	9	5			
f		7	6	7	5	7			
+		12	7	11	4	5			
-		7	11	8	15	15			
d		11	13	11	15	16			
D		12	10	12	5	5			
(F,f,+)(-,d,D)		23/30	19/34	22/31	18/35	17/36			
Ratio (Negative/Total)		0.57	0.64	0.58	0.66	0.68			

**Table 4** - Rainfall status of the Indian rainfall (IMR) and the DJF, MAM, JJA, SON seasons at Fortaleza for El Niño years of the types **a)** Unambiguous ENSOW, **b)** Ambiguous ENSOW, **c)** Other El Niños, **d)** All El Niños. Symbols +, f, F represent positive deviations in the ranges 0 to 0.5 σ, 0.5 to 1.0 σ and, exceeding 1.0 σ, while -, d, D represent negative deviations in the ranges 0 to -0.5 σ, -0.5 to -1.0 σ and, exceeding -1.0 σ. S (strong), M (Moderate), W (Weak) indicate the strength of the El Niño as given by Quinn et al. (1987), while I and II indicate the first and second years of double events (El Niños in two consecutive years, 1957-1958 etc.).

an overwhelming affinity for negative deviations (droughts). The ratio of (Negative/Total) was 1.00. For DJF (Fortaleza) also, the ratio was quite high (0.81, 13 out of 16 events negative), but for MAM (Fortaleza), the ratio was smaller (0.62). Thus, the relationship of Unambiguous ENSOW was much better with pre-season DJF rainfall than with main-season MEM rainfall. Considering that there would be

roughly equal numbers of positive and negative deviations by random chance (Ratio Negative/Total=0.50), MAM and SON (ratios 0.62) had only a slight bias for droughts while JJA (ratio 0.56) had a very poor relationship.  
 (2) For the 19 events of type Ambiguous ENSOW in Table 4 (b), the Indian rainfall IMR had a ratio 0.26, indicating a slight bias towards excess rains rather than droughts. There were only 2 events of

mild droughts, 1925 and 1992. Many of the events were II year events, thus indicating that *for India, the II year events would not be associated with droughts and may even have excess rains*. This was dramatically confirmed in the recent 1997-1998 El Niño, which was strongest in known history, had large SST anomalies during the Indian monsoon period June-September, but the anomalies reached *maximum only by 1997 end* and hence, both 1997 and 1998 qualified as Ambiguous ENSOW. *Therefore, no droughts were expected in India in 1997 and 1998, and none occurred*. For DJF and MAM at Fortaleza, ratios were 0.58, indicating poor relationship, but the II year events mostly had droughts. Thus, *for Fortaleza, the II year events were favourable for droughts in pre- and main-rainy season*. This was confirmed in 1997-1998, when 1998 had severe droughts. Incidentally, for the JJA and SON seasons, the ratios were high (0.74 and 0.89), indicating that the post-season rainfalls were likely to be deficit during Ambiguous ENSOW, including during the II years.

(3) Table 4 (c) for Other EN consists other El Niños of four types:

- (i) ENSOs, when SO existed but Pacific temperatures were normal (no W). There were 6 events and Indian rainfall was mostly deficit, as if these were Unambiguous ENSOW. For Fortaleza, results were mixed, droughts in some events and excess rains in others.
- (ii) ENW, when Pacific was warmer, but SOI was normal (not negative, SO did not exist). There was only one event, with mild excess rains in India and droughts in DJF, MAM, JJA of Fortaleza.
- (iii) EN, when an El Niño was listed in Quinn et al. (1987) list, but SO and W did not exist. IMR and Fortaleza showed mixed results, droughts in some events and excess rains in others.
- (iv) ENC. There were 6 events. Here, it does not mean that EN (El Niño) and C (La Niña) occurred simultaneously. (This is impossible). What happened was that El Niño existed in the early part of the calendar year and a La

Niña followed later. For rainfalls occurring in the early part of the year, the expected effect would be of EN (droughts), while for rainfalls occurring in the latter part of the year, the expected effect would be of C (excess rains). In India, there were mostly excess rains (effect of C), while at Fortaleza, effects were mixed (droughts in 1889, 1907, excess rains in 1917, 1973) in DJF and MAM. The overall ratios were near 0.50, indicating poor relationships, but some events had severe floods, while others had deficit rains. Thus, only Unambiguous ENSOW had significant relationships with droughts. In other cases, results were mixed.

- (v) Table 4 (d) shows the results for all El Niños (total 53 events). The ratios are near 0.50, indicating poor relationship if all events are considered.

### (B) Evolution of El Niños in the Pacific

The El Niños generally appear first in the Peru-Ecuador coast and later spread into the Pacific. Fig 2 (a) shows a plot of SST anomalies at Puerto Chicama (Peru coast) at the top, followed by anomalies in regions Niño 1+2, Niño 3, Niño 3.4 and Niño 4, for the El Niño of 1957-1958, for which data were available only from 1950 onwards. As can be seen, the anomaly magnitudes are largest near the coast (80° W), decrease to about half in the Niño 1+2 region (90° W-80° W), still lesser in the Niño 3.4 region (170° W-120° W) and still lesser in the Niño 4 region (160° E-150° W). The commencement (threshold 0.4° C as suggested by Trenberth, 1998) was in February 1957 at Puerto Chicama but later by a month or two westward in the Pacific. Similar plots were made for all the major El Niño events since 1950 and all showed the above characteristics, namely, decrease of magnitude and commencement later by about 2 months, away from the coast. However, the event of 1982-1983 was a glaring exception. As shown in Fig. 2 (b), the magnitudes decreased away from the coast, but the event seems to have started earlier in western Pacific, by May 1982 in Niño 3, 3.4, 4 regions, by July 1982 in Niño 1+2 region, and by October 1982 at Puerto Chicama on the Peru coast.

Fig. 2 (c) shows the recentmost giant event of 1997-1998, which shows characteristics similar to those of the 1957-1958 event.

**(C) Evolution of Fortaleza rainfall anomalies in individual El Niño events**

Stastical results have their own value, but individual events are more illustrative, particularly because El Niños may start in different months (Deser and Wallace, 1987) and may have different implications for rainfalls in different seasons. Many El Niños in the past had started during January or February or March and some statistical results are presented in Kane (2000). Fig. 3 shows the rainfall evolution at Fortaleza during several individual events, one frame of two rows for each event. Since SST anomalies at Puerto Chicama give a reasonably good representation of the El Niño evolution in the Pacific, only SST anomalies at Puerto Chicama are plotted in the first row, and rainfall normalized deviations at Fortleza in the second row. Positive anomalies are painted black and negative anomalies are shown hatched, for both SST and rainfall. Thus, black SST is expected to match with hatched rainfall, and vice versa. The following may be noted in Fig. 3:-


(1) Event of 1925-1926: Data for SST are not available for 1924, but the event started probably

in the latter half of 1924. Fortaleza had excess rains in the beginning of 1924, probably because of La Niña conditions at that time. The numbers 1, 2, 3, 4, in succession indicate rainfall deviations for 1 (DJF), 2 (circled, MAM), 3 (JJA), 4 (SON). During the first half of 1925, the El Niño was strong and Fortaleza 2, 3, 4 were negative, as expected. The El Niño weakened and disappeared by February 1926, but droughts prevailed in 1926 and 1927 in all seasons except MAM (not expected).


(2) Events of 1929-1932: There was a mild El Niño in the first half of 1929, a strong El Niño in 1930, La Niña in 1931 and a weak El Niño in the first half of 1932. The rainfalls were largely deficit in all the 4 years in all seasons except DJF of 1931 and 1932. The deficits in 1931 were not expected, as it was a La Niña year.

(3) Event of 1940-1941: Year 1939 was normal, but the rainfalls were highly excess (not expected). The El Niño commenced late in 1940, terminated by middle of 1941, but erupted again mildly and lasted upto 1941 end. But large deficit rains were seen in the whole of 1941 and 1942, in all seasons. The 1942 deficits were unexpected.

(4) Events of 1951 and 1953: These events were preceded by La Niñas in 1950 and 1952. But, except for SON in 1951 and MAM in 1952, the



**Figure 2** - Plots of SST anomalies at Puerto Chicama (Peru coast) and in regions Nino 1+2, Nino3, Nino3.4, Nino 4 in the Pacific for the El Niño events of a) 1957-1958, b) 1982-1983, c) 1997-1998. Positive anomalies are blackened. Full circles indicate commencements of the SST anomalies.



**Figure 3** - Plots for four consecutive years having El Niño events in any one or two years. In each frame, the upper plot is for Puerto Chicama SST anomalies and the lower plot for Fortaleza normalized rainfall for seasons 1 (DJF), 2 (circled) (MAM), 3 (JJA), 4 (SON). Positive anomalies are blackened and negative anomalies are hatched.



- whole period had heavy rain deficits in all seasons (not expected), with poor matching with El Niños.
- (5) Event of 1957-1958: The event started in February of 1957 and lasted upto April of 1958. But the whole period 1956-1959 had large deficit rains. The deficit in 1956 was unexpected, as 1956 SST was normal.
  - (6) Event of 1965: This was a moderate event, starting in March 1965 and ending in February 1966. But the whole period was characterised by large excess rains except in JJA of 1963 and DJF of 1966 when large deficits occurred. Years 1964 and 1966 had La Niña conditions, but the overall matching was not good.
  - (7) Event of 1972: The event started in February of 1972, lasted for about 12 months and was followed by La Niña conditions in 1973-1974. By and large, SST positive anomalies were associated with rainfall deficits, and negative SST anomalies with excess rains, in all seasons. JJA rainfall anomalies showed exceptions.
  - (8) Event of 1976: This was a moderate event during 1976 only and showed matching moderate rain deficit in 1976. Year 1975 was a La Niña year and showed rainfall excesses. However, 1977 and 1978 were normal SST years but showed abnormal excess rains in JJA and SON (not expected).
  - (9) Event of 1982-1983: This was a strong event, starting in October 1982 at Puerto Chicama ( but earlier in the Pacific, as shown by the dashed line) and lasting upto late 1983. But there were large rain deficits in 1981 and early 1982 (not expected). Year 1984 was a mild La Niña. The deficit in 1983 and excess in 1984 was as expected.
  - (10) Event of 1987: This was a moderate event confined to the first half of 1987 and had a matching rain deficit. Years 1986, 1988 and 1989 had La Niña conditions and had matching rain excesses.
  - (11) Event of 1991-1993: There was a mild El Niño in the first few months of 1991, a strong event in the end of 1991 lasting upto the middle of 1992, and again a small event in the first half of 1993. The rainfalls were matching, except in the beginning of 1991 when droughts were expected but did

not occur, and in 1990 when SST was normal, but large rainfall deficits followed by excesses occurred.

- (12) Event of 1997-1998: By far the largest event, it commenced in February-March 1997 and ended in June 1998. There were matching rainfall deficits, but there were deficits in late 1996 also, before the El Niño commencements (not expected).

Thus, whereas the rainfall deficits were generally associated with positive SST anomalies and rainfall excesses with negative SST anomalies, *there were often rainfall anomalies with no corresponding SST anomalies*. Obviously other factors were intervening in some cases. .

#### **(D) Characteristics of years of rainfall extremes**

Table 5 lists the years of severe droughts and severe excess rains (deviations exceeding  $1.0 \sigma$ ) in DJF and MAM seasons at Fortaleza, and the corresponding characteristics of these years in our finer classification. Unambiguous ENSOW are denoted as U, Ambiguous ENSOW as A and Non-events as Non..

An examination of Table 5 reveals that from the 15 severe droughts in DJF, 10 occurred as (3U, 3A, 1ENSO, 1ENW, 1EN, 1SO), all El Niño-like events, but 5 occurred as (4C, La Niñas, 1 Non). From the 21 severe excess rain years in DJF, 6 occurred as (1U, 1ENSO, 1EN, 1SOW, 1SO, 1W), all El Niño-like events, not supposed to cause floods. Similar discrepancies are seen for MAM also. Thus, severe droughts can occur not only in El Niño-like years, but even in C (La Niñas) and Non-events, and severe excess rains can occur not only in C (La Niña) years but even in El Niño-like years . Under these circumstances, expecting a good relationship with El Niños in general is obviously futile and frustrating. It is gratifying to note that the relationship with Unambiguous ENSOW is fairly good, but obviously, that is not the only source. If severe and mild events are considered, there were 47 droughts in DJF and 41 droughts in MAM, only 23 of which were associated with El Niños (of all types, see Table 4 d). Factors unrelated to the ENSO (El Niño/Southern Oscillation) must be playing a considerable role.

(b) Severe excess rains F (exceeding 1.0  $\sigma$ )

Excess rains in DJF season			Excess rains in MAM season		
Year	Characteristic	Rainfall ( $\sigma$ )	Year	Characteristic	Rainfall ( $\sigma$ )
1872	C	2.24	<b>1894</b>	C	2.53
1873	C	1.92	1895	Non	1.81
1893	C	2.25	1896	U	1.56
<b>1894</b>	C	2.24	<b>1899</b>	U	2.16
<b>1899</b>	U	2.21	<b>1910</b>	C	1.10
1901	Non	1.14	<b>1912</b>	ENSO	2.13
<b>1910</b>	C	1.10	1924	C	1.16
<b>1912</b>	ENSO	2.13	1947	Non	1.00
1913	SOW	1.20	1949	SOC	1.69
1917	ENC	2.77	1963	A	1.31
1921	C	1.71	<b>1964</b>	C	1.11
1935	SOC	1.86	<b>1974</b>	SO	1.88
1939	EN	1.64	1995	A	1.76
1945	Non	1.20			
1961	C	1.80			
<b>1964</b>	C	3.60			
1973	ENC	1.48			
<b>1974</b>	SO	1.00			
1980	Non	1.64			
1985	Non	1.04			
1986	W	1.90			
Total	Severe F	21 events	Total	Severe F	13 events
	Mild f	13 events		Mild f	26 events
	F and f	34 events		F and f	39 events

(a) Severe droughts D (exceeding -1.0  $\sigma$ )

Droughts in DJF season			Droughts in MAM season		
Year	Characteristic	Rainfall ( $\sigma$ )	Year	Characteristic	Rainfall ( $\sigma$ )
<b>1877</b>	U	-1.43	<b>1877</b>	U	-1.92
<b>1878</b>	A	-1.03	<b>1878</b>	A	-1.56
<b>1879</b>	C	-1.06	<b>1879</b>	C	-1.48
1880	ENSO	-1.36	1888	U	-1.47
1884	ENW	-1.31	1889	ENC	-1.28
1885	SO	-1.23	<b>1898</b>	C	-1.50
1892	C	-1.04	1900	ENSO	-2.37
<b>1898</b>	C	-1.24	1902	U	-1.05
1905	U	-1.01	1903	C	-1.34
1943	EN	-1.10	1907	ENC	-1.47
<b>1951</b>	U	-1.49	1908	C	-1.18
<b>1958</b>	A	-1.29	1915	Non	-1.84
1960	C	-1.44	1919	A	-1.89
1966	Non	-1.20	1932	EN	-1.64
1997	A	-1.26	1936	SOC	-1.34
			<b>1951</b>	U	-1.34
			1956	C	-1.21
			<b>1958</b>	A	-1.80
			1980	Non	-1.76
			1983	A	-1.28
Total	Severe D	15 events	Total	Severe D	20 events
	Mild d	32 events		Mild d	21 events
	D and d	47 events		D and d	41 events

**Table 5** - Years of **a**) severe droughts and **b**) severe excess rains, (magnitudes exceeding 1.0  $\sigma$ ) in DJF and MAM seasons at Fortaleza during 1872-1999, their normalized deviations, and their characteristics, where Unambiguous ENSOW are denoted by U, Ambiguous ENSOW by A, and Non-events by Non. Years common to DJF and MAM are shown bold. (At the bottom, the number of mild droughts and mild excess rains are also indicated).

The relationship of DJF droughts with some El Niños has an interesting aspect. The DJF attributed to any year is of December of the previous year and January, February of the current year. (For example, DJF attributed to 1950 is for December 1949 and January-February of 1950). As such, the DJF rainfall is centered in the January of the year mentioned and hence, *precedes the El Niños*, which commence in that year. How can rainfall react to El Niño which commenced later? Does it imply that DJF rainfall deficits can be *precursors* of El Niños? This needs further investigation. Hastenrath (1990) uses this fact in his regression analysis for rainfall predictions in NE Brazil.

## ROLE OF OTHER FACTORS USED FOR PREDICTIONS

Even though the mass media (newspapers, radio, television) refer only to El Niños and La Niñas as having major influences on rainfalls, many in the scientific community are aware of the important roles played by other factors *unrelated to ENSO*. In India, Himalayan Eurasian snow cover and many other factors are known to affect the rainfall regime and these factors are taken into account for prediction purposes (Thapliyal and Kulshrestha, 1992). Based on these formulations, IMD (India Meteorological Department) issues forecasts by May, applicable to the June-Sept. monsoon season. The forecasts are coming out fairly correct for the last 10 years or more. For NE Brazil, a considerable influence of tropical Atlantic SST was reported long ago (Markham and McLain, 1977). Other factors considered are, 700 mb circulation pattern over the North Atlantic (Namias, 1977), meridional displacement and strength of the Intertropical Convergence Zone (ITCZ) (Hastenrath and Heller, 1977), rainfall systems associated with tropical disturbances moving westward from the Atlantic towards northeast Brazil (Ramos, 1975; Yamazaki and Rao, 1977; Rao et al., 1993), and southern hemisphere cold fronts or their remains moving northward along the northeast coast of Brazil (Kousky and Chu, 1978; Kousky, 1979). There is a well defined large-scale atmospheric circulation pattern related to the sea surface temperature anomalies in the tropical Atlantic (Hastenrath and Heller, 1977;

Moura and Shukla, 1981). Hastenrath et al. (1984) and Hastenrath (1990) formulated prediction schemes involving zonal and meridional wind components over limited areas of the equatorial Atlantic, SST in tropical North and South Atlantic, SO index and pre-season rainfall itself in northeast Brazil, as predictors. Hastenrath and Greischar (1993) elaborated their list of predictors and Hastenrath and Druyan (1993) complemented this study by evaluating output of a 7-year run of the general circulation model (GCM) of the Goddard Institute of Space Studies (GISS). Earlier, Servain and Siva (1987) had investigated the relationship between tropical Atlantic SST, wind stress and regional precipitation indices and shown that for the seasonal time scale, the northward displacement of the ITCZ was accompanied by the strengthening of the southeast trades and/or relaxation of the northeast trades which is correlated with a decrease in NE Brazil rainfall. Recently, Wainer and Soares (1997) showed that for NNE Brazil, this was true on an interdecadal time scale also. Ward and Folland (1991) investigated the relationship between north northeast Brazil rainfall and SST in various parts. They calculated the covariance eigenvectors for SST in the Atlantic and Pacific Oceans. The Atlantic eigenvector 3 and Pacific eigenvector 1 had the best correlations, sufficient to provide a preliminary forecast of the March-April rainfall of northeast Brazil. Using two statistical techniques viz. MLR (Multiple Linear Regression) and LDA (Linear Discriminant Analysis), forecasts were made for 1987, 1988, 1989 and 1990 and were found to be good. Following the methodology outlined in Ward and Holland (1991), experimental forecasts of northeast Brazil rainfall at 1 and 0 month leads are issued using November-January and January-February predictor data. The worst forecast was for 1996 (predicted below normal, observed far above normal), probably due to a sharp change in Atlantic SST through the forecast season. These forecasts are *statistical* predictions. In the same group (UK Meteorological Office, Bracknell), Harrison makes a *dynamical* prediction of northeast Brazil rainfall using a version of the UKMO climate atmospheric general circulation model (AGCM). For 1996, the AGCM forecast was rainfall about 10% below average. Hence, both these forecasts turned out to be in *disagreement* with the observed rainfall

in 1996. Using an atmospheric GCM with persisted SST anomalies, Graham (1996) has been forecasting northeast Brazil rainfall since 1993. For 1996, his forecast proved erroneous. The forecast of Greischer and Hastenrath (1996), using the method of 5 predictors, also proved erroneous for 1996.

In summary, whereas forecasts for earlier years were reasonably correct, forecasts for 1996 were miserably poor in all these methods, probably because of the rapid changes in Atlantic SST during the forecast season, as mentioned by Colman et al. (1997b).


The forecasts for northeast Brazil rainfall for March-May 1997 were mostly of moderately dry conditions (Colman et al., 1997b; Harrison et al., 1997c; Graham, 1997; Greischar and Hastenrath, 1997b). The observed rainfall was a few percent below normal in some parts and normal in others, thus, basically conforming to the above predictions. The El Nino was still strong in January 1998 but was showing signs of weakening. Meanwhile, a dipole was developing in the Atlantic. A forecast given in the Experimental Long-Lead Forecast Bulletin (Colman et al., 1998) prepared in early March 1998 using February 1998 SST gave the best estimate for the most likely category as WET. However, it was also mentioned that, given the continuing strong El Nino, and the fact that the atmospheric model placed positive (wet) anomalies over the Nordeste, *the possibility of a dry or very dry season should also be considered*. In the same Bulletin, Greischar and Hastenrath (1998) also predicted wetter than normal conditions but pointed out that the equatorial Pacific SST and the field of the meridional wind component in the Atlantic sector favour drier conditions. Hence, the precipitation should be only *slightly above average*. In the same Bulletin, Cavalcanti et al. (1998a) reported predictions from CPTEC, INPE, BRAZIL, based on the CPTEC version of the COLA AGCM. Their prediction was of a *below average* precipitation over much of northeast Brazil, which came true, as the main rainy season in NE north Brazil (March, April, May 1998) suffered one of the *severest droughts in known history*. The media attributed it to the 1997-98 El Nino. Thus, in spite of the progress in understanding the mechanisms which affect rainfall in NE Brazil, predictions are still hazardous and could be widely in error. For the March-May 1999 rainfall

season of northeast Brazil, Cavalcanti et al. (1998b) and Colman and Davey (1998) predicted excess rainfall and the predictions came partly true. For the March-May 2000 rainfall, predictions are of excess rains (Cavalcanti et al., 1999; Colman and Davey, 1999).

The results presented here are for one location, Fortaleza. However, there is reason to believe that these are applicable to about 100 locations in NE Brazil, near about Fortaleza (Kane, 1999 a,b)

## PREDICTION BASED ON EXTRAPOLATION OF PERIODICITIES

Since Markham (1974) demonstrated that there were long-term periodicities (13 and 26 years) in the Fortaleza rainfall, attempts have been made to use the periodicities for predictions (Kane and Trivedi, 1986 and references therein). Maximum Entropy Spectral Analysis indicates significant periodicities at  $T=2-3, 3-4, 5-6, 13, 26$  and 50 years, but these are transient, with relative magnitudes different in different intervals. An update of the situation is given in Kane (1997c) where severe droughts are indicated during 2000-2010, but the prediction skill is not likely to be high. Probably, spectra from more recent data may give more reliable (or less uncertain) predictions, though the uncertainty about the magnitude and phase of the QBO and a QTO (Quasi-biennial and quasi-triennial oscillations) in any interval is a great stumbling block. Nevertheless, a spectral analysis was attempted using the seasonal values. Four-season (12-month) running means of Fortaleza rainfall were calculated for 1975-1997 (23 years, 92 seasonal values) and spectrally analysed. With a LPEF=30% of the data length, two significant peaks were obtained at  $T=14.24$  and 30.45 seasons, i. e., 3.6 and 7.6 years. (No peak was obtained at 13 or 26 years). Fig. 4 (a) shows a plot of the 92 seasonal values (thin line), while the thick line represents the values reconstructed by using the amplitudes and phases of the two periodicities. The matching is not very good for the dependent data 1975-1997 (correlation +0.42) and hence the predictions for the independent data (1998 onwards) may not be reliable; but increasing rainfall for 1998 onwards is indicated. Using LPEF=50% of data length, the plots were as in Fig. 4 (b). The thin line is



**Figure 4** - Plots of the 4-season (12-month) running means (thin lines) and the reconstructed values (thick lines) using periodicities **a)**  $T=3.6, 7.6$  years, **b)**  $T=2.69, 3.40, 4.97, 10.09$  years.

the plot of original values (4-season running means) while the thick line shows the reconstructed values, using the amplitudes and phases of four significant periodicities  $T=10.74, 13.60, 19.89, 40.37$  seasons i.e., 2.69, 3.40, 4.97, 10.09 years. (Notice that the 7.6-year periodicity obtained in LPEF=30% now split into 4.97 and 10.09 years for LPEF=50%). In Fig. 4 (b), the matching is very good for 1975-1997 (correlation +0.84) and the predictions are of above normal rainfalls for 1999 and 2000, but lower rainfalls for 2001 and 2002. However, it must be remembered that larger the number of periodicities considered, better is the fit for dependent data but worse for the independent data. Also, this method has severe limitations, mainly because of the transient nature of the QBO and QTO (Kane, 1998d), and, if an El Niño develops suddenly by 2000 end or early 2001, the situation may change, with a severe drought in 2001.

## CONCLUSIONS

Fortaleza ( $3^{\circ} 42' S, 38^{\circ} 31' W$ ) Ceara, north-east Brazil is prone to frequent severe droughts. The rainfalls in each of the seasons DJF (December, January, February), MAM (March, April, May), JJA (June, July, August), SON (September, October, November) for 1849-1999 were expressed in normalised units (deviations from the overall mean rainfall of the series of that season, divided by the standard deviation of the series) and the characteristics studied. The following was noted:-

- (1) The four series were poorly intercorrelated. A correlation of +0.45 or less was obtained, implying a common variance of only ~20%

(square of the correlation 0.45). Thus, rainfalls of the four seasons varied almost independently, though similar variations were sometimes seen between DJF and MAM.

- (2) Relationship with El Niños was poor, with only about half of the El Niños associated with droughts. Among the El Niños, Unambiguous ENSOW had a better relationship, more with DJF than with MAM. But many droughts occurred when there were no El Niños or even during La Niña conditions. Similarly, some La Niña years had excess rains.
- (3) Obviously, other factors unrelated to ENSO disrupt the ENSO relationship. Forecasts taking into consideration Atlantic parameters seem to give good results. As such, the undue importance given to El Niño and La Niña in the media needs to be pointed out and diluted.

Since El Niños may commence in different months, the differential behaviour of DJF and MAM rainfall is understandable. El Niños starting in say, March-April, may affect the MAM rainfall but not the DJF rainfall. On the other hand, predictions based on Atlantic parameters are issued from information available in December and hence, the predictions may turn out to be more true for DJF than for MAM. In December 1995, slightly below rainfall was predicted and the DJF rainfall was just normal (+0.03). But the Atlantic conditions changed rapidly in favour of excess rains and the MAM rainfall turned out to be far above normal (+0.96). These hazards in prediction need to be kept in mind. Relying on El Niño alone will certainly be imprudent.

## ACKNOWLEDGEMENTS

Thanks are due to Dr. Todd Mitchell and Dr. Don Garrett for supplying Puerto Chicama SST data, and to Dr. Don Garrett for El Niño and other data through the Web site of Climate Prediction Center, NOAA, NCEP, Washington. This work was partially supported by FNDCT, Brazil under contract FINEP-537/CT.

## REFERENCES

- Cavalcanti, I. F. A., Pezzi, L. P., Nobre, P., Sampaio, G., & Camargo Jr., H., 1998a.** Climate prediction of precipitation in Brazil for the nordeste rainy season, Center for Ocean-Land-Atmosphere Studies (COLA), Calverton, Maryland, Experimental Long-lead Forecast Bulletin 7, No. 1, March 1998, pp. 24-27.
- Cavalcanti, I. F. A., Pezzi, L. P., Marengo, J., Sampaio, G., & Sanches, M. B., 1998b.** Climate Prediction of precipitation over South America for DJF 1998/99 and MAM 1999, Center for Ocean-Land-Atmosphere Studies (COLA), Calverton, Maryland, Experimental Long-lead Forecast Bulletin 7, No. 4, December 1998, pp.
- Cavalcanti, I. F. A., Marengo, J. A., Castro, C., Sampaio, G., & Sanches, M. B., 1999.** Climate Prediction of precipitation over South America for DJF 1999/2000 and MAM 2000 using the CPTEC/COLA AGCM, Center for Ocean-Land-Atmosphere Studies (COLA), Calverton, Maryland, Experimental Long-lead Forecast Bulletin 8, No. 4, December 1999, pp. 43-46.
- Colman, A., & Davey, M., 1998.** Statistical Prediction of January-June 1999 Rainfall in Northeast Brazil using input from Multiple Regression and Discriminant Analysis, Center for Ocean-Land-Atmosphere Studies (COLA), Calverton, Maryland, Experimental Long-lead Forecast Bulletin 7, No. 4, December 1998, pp.
- Colman, A., & Davey, M., 1998.** Statistical Prediction of January-June 2000 Rainfall in Northeast Brazil using input from Multiple Regression and Discriminant Analysis, Center for Ocean-Land-Atmosphere Studies (COLA), Calverton, Maryland, Experimental Long-lead Forecast Bulletin 8, No. 4, December 1999, pp. 66-69.
- Colman, A., Davey, M., Harrison, M., Evans, T. & Evans, R., 1997.** Multiple Regression, Discriminant Analysis predictions of Mar-Apr-May 1997 rainfall in northeast Brazil, Climate Prediction Center (CPC), Experimental Long-lead Forecast Bulletin 6, No.1, 97, pp. 29-32, Natl. Cent. for Environ. Predic., Washington, D. C., March 1997.
- Colman, A., Davey, M., Harrison, M., & Evans, A., 1998.** Prediction of March-April-May 1998 rainfall in northeast Brazil using input from Multiple Regression, Discriminant Analysis and an Atmospheric Global Circulation Model, Center for Ocean-Land-Atmosphere Studies (COLA), Calverton, Maryland, Experimental Long-lead Forecast Bulletin 7, No. 1, March 1998, pp. 75-79.
- Deser, C. & Wallace, J. M., 1987.** El Niño events and their relation to the Southern Oscillation: 1925-1986, *J. Geophys. Res.* 92, 14189-14196.
- Graham, N., 1996.** Prediction of precipitation in northeast Brazil for boreal spring 1996 using an atmospheric GCM with persisted SST anomalies, Climate Prediction Center (CPC) Experimental Long-lead Forecast Bulletin 5, No. 1, pp. 5-6, Natl. Cent. for Environ. Predic., Washington, D. C., March 1996.
- Graham, N., 1997.** Prediction of precipitation in northeast Brazil for boreal spring 1997 using an atmospheric GCM with persisted SST anomalies, Climate Prediction Center (CPC), Experimental Long-lead Forecast Bulletin 6, No. 1, pp. 23-24, Natl. Cent. for Environ. Predic., Washington, D. C., March 1997.
- Greischar, L. & Hastenrath, S., 1997.** Multiple Regression and Discriminant Analysis to predict Mar-Apr-May-Jun 1997 rainfall in northeast Brazil, Climate Prediction Center (CPC), Experimental Long-lead Forecast Bulletin 6, No. 1, 33-34, Natl. Cent. for Environ. Predic., Washington, D. C., March 1997.
- Greischar, L. & Hastenrath, S., 1998.** Multiple Regression and Discriminant Analysis to predict Mar-Apr-May-Jun 1998 rainfall in northeast Brazil, Center for Ocean-Land-Atmosphere

Studies (COLA), Calverton, Maryland, Experimental Long-lead Forecast Bulletin 7, No. 1, March 1998, pp.80-81.

- Harrison, M., Evans, T., Evans, R., Davey, M. & Colman, A., 1997.** A dynamical one-month lead seasonal rainfall prediction for March to May 1997 for the north-eastern area of South America, Climate Prediction Center (CPC), Experimental Long-lead Forecast Bulletin 6, No. 1, pp. 25-28, Natl. Cent. for Environ. Predic., Washington, D. C., March 1997.
- Hastenrath, S., 1990.** Prediction of northeast Brazil rainfall anomalies. *J. Climate* **3**: 893-904.
- Hastenrath, S. & Druyan, L., 1993.** Circulation anomaly mechanisms in the tropical Atlantic sector during the Northeast Brazil rainy season: Results from the GISS General Circulation model. *J. Geophys. Res.*, **98**: 14917-14923.
- Hastenrath, S. & Greischar, L., 1993.** Circulation mechanisms related to Northeast Brazil rainfall anomalies. *J. Geophys. Res.*, **98**: 5093-5102.
- Hastenrath, S. & Heller, L., 1977.** Dynamics of climate Hazards in northeast Brazil. *Q. J. Roy. Meteorol. Soc.* **103**: 77-92.
- Hastenrath, S., Wu M-C & Chu, P-S., 1984.** Towards the monitoring and prediction of Northeast Brazil droughts. *Q. J. Roy. Meteorol. Soc.* **110**: 411-425.
- Kane, R. P., 1997a.** Relationship of El Nino/Southern oscillation and Pacific sea-surface temperature with rainfall in various regions of the globe, *Mon. Wea. Rev.*, **125**: 1792-1800.
- Kane, R. P., 1997b.** On the relationship of ENSO with rainfall over different parts of Australia, *Aust. Meteor. Mag.*, **46**: 39-49.
- Kane, R. P., 1997c.** Prediction of droughts in north-east Brazil: Role of ENSO and use of periodicities, *Int. J. Climatol.*, **17**: 655-665.
- Kane, R. P., 1998a.** Extremes of the ENSO phenomenon and the Indian summer monsoon rainfall, *Int. J. Climatol.*, **18**: 775-791.
- Kane, R. P., 1998b.** ENSO relationship to the rainfall of Sri Lanka, *Int. J. Climatol.*, **18**, 859-871.
- Kane R. P., 1998c.** El Niño, Southern Oscillation, equatorial eastern Pacific sea surface temperatures and summer monsoon rainfall in India. *Mausam*, **49**: 103-114.
- Kane, R. P., 1998d.** Quasi-biennial and quasi-triennial oscillations in the rainfall of northeast Brazil, *Rev. Brasileira Geofisica*, **16**: 37-52.
- Kane, R. P., 1999a.** Rainfall extremes in some selected parts of central and south America: ENSO and other relationships reexamined, *Int. J. Climatol.*, **19**, 423-455.
- Kane, R. P., 1999b.** El Niño timings and rainfall extremes in India, southeast Asia and China. *Int. J. Climatol.* **19**: 653-672.
- Kane, R. P., 2000.** Relationship between El Niño timings and rainfall extremes in NE Brazil, São Paulo city and south Brazil. *Rev. Bras. Meteorologia*, **15**, (No.1),
- Kane, R. P., & Trivedi, N. B., 1986.** Are droughts predictable? , *Climatic Change*, **8**, 209-223.
- Kousky, V. E., 1979.** Frontal influences on Northeast Brazil. *Mon. Wea. Rev.*, **107**: 1140-1153.
- Kousky, V. E. & Chu, P. S., 1978.** Fluctuations in annual rainfall for Northeast Brazil. *J. Meteorol. Soc. Japan*, **56**: 457-465.
- Markham, C. G., 1974.** Apparent periodicities in rainfall at Fortaleza, CE, Brazil, *J. Appl. Meteorol.* **13**: 176-179.
- Markham, C. G. & McLain, D., 1977.** Sea surface temperature related to rain in Ceara, north-eastern Brazil, *Nature* **265**: 320-323.
- Moura, A. D., & Shukla, J., 1981.** On the dynamics of droughts in Northeast Brazil: Observations, theory and numerical experiments with a General Circulation model. *J. Atmos. Sci.* **38**: 2653-2675.
- Namias, J., 1977.** Influences of northern hemisphere general circulation on drought in Northeast Brazil. *Tellus* **24**, 336-342.
- Quinn, W. H., Neal, V. T. & Antunes de Mayolo, S. E., 1987.** El Nino occurrences over the past four and a half centuries, *J. Geophys. Res.*, **92**: 14449-14461.
- Ramos, R. P. L., 1975.** Precipitation characteristics in the northeast Brazil dry region, *J. Geophys. Res.*, **80**: 1665-1678.
- Rao, V. B., de Lima, M. C. & Franchito, S. H., 1993.** Seasonal and interannual variations of rainfall over eastern Northeast Brazil. *J. Climate*, **6**: 1754-1763.

- Servain, J. and Seva, M., 1987.** On relationship between tropical Atlantic sea surface temperature, wind stress and regional precipitation indices: 1964-1984, *Ocean Air Interaction* **1**: 183-190.
- Thapliyal, V. & Kulshrestha, S. M., 1992.** Recent models for long-range forecasting of southwest monsoon rainfall over India, *Mausam* **43**: 239-248.
- Trenberth, K. E., 1998.** Development and forecasts of the 1997-98 El Nino: CLIVAR Scientific Issues, Exchanges, **3**: 4-14.
- Wainer, I. & Soares, J., 1997.** North northeast Brazil rainfall and its decadal-scale relationship to wind stress and sea surface temperature, *Geophys. Res. Lett.*, **24**: 277-280.
- Ward, M. N. & Folland, C. K., 1991.** Prediction of seasonal rainfall in the north nordeste of Brazil using eigenvectors of sea-surface temperature, *Int. J. Climatol.*, 711-743.
- Yamazaki, Y. & Rao, V. B., 1977.** Tropical cloudiness over the south Atlantic Ocean. *J. Meteorol. Soc. Japan*, **55**: 205-207.

#### NOTE ABOUT THE AUTHOR

Dr. Rajaram Purushottam Kane was born in India in 12th November 1926. His earlier education was in Gwalior, Madhya Pradesh, India. He received his B.Sc. in 1946 from Banaras Hindu University, India as a post-graduate student in 1948 and received his PhD. on "Time-variation of Cosmic Ray Intensity near Geomatic Equator" from Bombay University, India in 1953. During 1953-54, he was awarded a Fullbright & Smith-Mundt Fellowship to work with Prof. J. A.

Simpson at the Institute of Nuclear Studies, University of Chicago, Chicago, USA. During 1955-56, he was a Senior Fellow of the National Institute of Science of India at PRL and later, till 1978, he was on the PRL Faculty. Since May, 1978, he is at INPE (Instituto Nacional de Pesquisas Espaciais) at São José dos Campos, SP, Brazil, working in the fields of Geomagnetism and Aeronomy, Solar Physics, Rainfall studies, etc.

### **CURSO DE PÓS-GRADUAÇÃO EM GEOFÍSICA ESPACIAL INSTITUTO NACIONAL DE PESQUISAS ESPACIAIS - INPE**

#### **Linhas de Pesquisa:**

**Alta Atmosfera  
Eletricidade Atmosférica  
Geomagnetismo  
Ionosfera  
Luminescência Atmosférica  
Magnetosfera-Heliosfera  
Média e Baixa Atmosfera**