



Revista Brasileira de Hematologia e Hemoterapia Brazilian Journal of Hematology and Hemotherapy

www.rbhh.org



Original article

Translation into Portuguese and validation of the Blood Donation Reactions Inventory

Ana Carolina Garcia Braz, Rodrigo Guimarães dos Santos Almeida,
Edson Zangiacomi Martinez*

Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

ARTICLE INFO

Article history:

Received 5 November 2013

Accepted 17 December 2013

Keywords:

Blood donors

Validation studies

Translations

Hemotherapy service

Factor analysis, statistical

A B S T R A C T

Background: the Blood Donation Reactions Inventory (BDRI) scale was proposed as part of a study about the predictors of psychological reactions in volunteer blood donors, as uncomfortable reactions are associated with a lower probability to return for further donations.

Objective: to translate the Inventory into Brazilian Portuguese and evaluate its psychometric properties (validity and reliability). The inventory has 11 items, but the literature suggests that shorter inventories, of four or six items, should be used.

Methods: this study was carried out at the blood center of Franca, Brazil. Three people with knowledge of English and familiarity with medical terms translated the Blood Donation Reactions Inventory into Brazilian Portuguese. Aiming to evaluate the objectivity and relevance of the items of the translated instrument, its content was independently evaluated by a panel of eight assessors. After this, data on 1,001 blood donors was collected. Internal consistency was assessed by Cronbach's alpha coefficient. An exploratory factor analysis with varimax rotation was used to analyze the measure for construct validity.

Results: the sample consisted of 65.8% men, and 27.3% first time donors. Internal consistency determined by Cronbach's alpha coefficient was satisfactory for the 11, 6 and 4-item scales. Considering the factor analysis, the 11-item scale seems to measure more than one construct as three factors were identified with eigenvalues greater than 1. These factors correspond to 'vasovagal adverse reactions', 'fear' and 'anxiety/excitation'.

Conclusion: the Portuguese version of the Blood Donation Reactions Inventory is a valid and reliable instrument for collecting information regarding systemic reactions experienced by blood donors. The 6-item scale seems to be useful when the objective is to measure only vasovagal adverse reactions.

© 2014 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular.
All rights reserved.

*Corresponding author at: Universidade de São Paulo, Faculdade de Medicina de Ribeirão Preto, Departamento de Medicina Social, Av. Bandeirantes, 3900, Monte Alegre, 14049-900, Ribeirão Preto, SP, Brazil.

E-mail address: edson@fmrp.usp.br (E.Z. Martinez).

1516-8484/\$ - see front matter © 2014 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. All rights reserved.

DOI: 10.5581/1516-8484.20140031

Introduction

The maintenance of blood component stocks is a constant challenge for blood transfusion services. Recruitment strategies of blood donors must aim to ensure the necessary number of donors considering seasonal variations in the number of donations,¹ and to focus on the quality of the material collected. In this context, the search for repeat blood donors is essential, as those tested and retested donors provide blood bags with a greater safety margin for the recipient, a smaller number of discarded blood components,² and a lower number of positive test results in the screening for infectious diseases.

The loyalty of blood donors can be enhanced using strategies that increase donation accessibility,³ training the professionals at the blood centers,⁴ and fulfilling the expectations and increasing the satisfaction of the donors regarding the service provided.⁴⁻⁷ The literature shows that factors associated with the return of a blood donor for further donations are Rh factor,⁸ age,^{8,9} education,⁸⁻¹⁰ and gender.⁹ In addition, some authors have also highlighted the need to pay attention to the physical effects of blood donation,¹⁰ as uncomfortable reactions are associated with a lower probability to return for further donations.^{11,12} Among these reactions are fainting, vertigo and dizziness.

The Blood Donation Reactions Inventory (BDRI) scale was proposed as part of a study about predictors of psychological reactions in volunteer blood donors.¹³ The scale consists of 11 items, each corresponding to a reaction or feeling regarding the latest blood donation: (1) faintness (such as feeling faint or losing consciousness), (2) dizziness, (3) weakness, (4) facial flush, (5) visual disturbance (such as blurred vision or tunnel vision), (6) difficulty hearing, (7) lightheadedness, (8) rapid or pounding heartbeat, (9) sweating, (10) rapid or difficult breathing and (11) nausea or upset stomach. Donors answer questions on these items after a donation using the 6-point Likert scale ranging from 0 ('not at all') to 5 ('to an extreme degree'). Thus, higher values are associated with greater reaction intensity. The responses are summed to the final score producing values between 0 and 55.

The psychometric properties of BDRI were evaluated by France et al.,¹⁴ who found high internal consistency for the scale, concurrent validity with other measures of reactions to blood donation and construct validity supported by a factor analysis. This study further showed that an abbreviated version of the BDRI containing only items 1, 2, 3 and 7 of the original instrument would have good psychometric properties and could replace the original scale of 11 items. France et al. also concluded that BDRI is brief, easily understood by donors, and quick to administer and score.¹⁴

In another study, France et al.¹⁵ showed that high BDRI scores are associated with a significant reduction in the probability of return of blood donors, suggesting that the instrument is an effective tool for predicting whether a donor will make further donations.

Routinely, the blood donors at the Regional Blood Center (Hemocentro) of Ribeirão Preto, Brazil, and at its satellite units, are observed during donation by a nurse who provides additional care to those who experience an adverse

reaction or accident during blood collection. When a blood donor presents an adverse reaction, the nurse completes a standard form entitled 'Notice of Adverse Reaction during Donation' (NARD). The nurse writes on this form which systemic reactions were experienced, their intensity, vital signs and other signs and symptoms, including possible incidents during blood collection. After completion, the nurse is responsible for adding data to the donor's record so that professionals performing the screening of future donations are informed about the incident. The translated version of BDRI does not intend to replace the NARD because its purpose is different. While the NARD allows routine monitoring of adverse reactions, the BDRI is useful for studies that evaluate reactions aimed to, for example, establish strategies to increase blood donor satisfaction and retention. In this context, France et al.¹⁵ cite some disadvantages of the measures based on the observations of the professionals responsible for blood collection in predicting the return of the donor. These professionals may not be able to detect subjective symptoms such as dizziness, vertigo, stress, nervousness or even excessive distress of the person who has a sharp object in his/her arm, and consequently cannot assess the intensity of the donors' reactions. This means that a 'slight' sensation perceived by the nurse may be so uncomfortable for the donor as to influence his/her decision not to return for further donations. Given that the BDRI assesses the experience from the perspective of the donor, it is more sensitive to the intensity of the reaction.

Thus, the objective of this study was to translate the BDRI into Brazilian Portuguese and study internal consistency, criteria, and construct validation of the translation. In this study, the 11-item version of the BDRI and the abbreviated 6- and 4-item versions proposed by France et al.¹⁴ will be considered. The 6-item scale considers items 1, 2, 3, 5, 7 and 11 of the original instrument with the respective total score ranging from 0 to 30 points. The 4-item scale considers items 1, 2, 3 and 7 with the respective total score ranging from 0 to 20 points.

Methods

Translation

First, three people with knowledge of English and familiarity with medical terms independently worked on the translation of the BDRI scale. It was intended that the meaning of the 11 terms describing the feelings and reactions of each item of the instrument had a literal correspondence between the original English version and the Portuguese translation. This resulted in three different versions, which were later compared in order to produce a final consensual version considering cultural aspects of the target population. Thus, the relevance, appropriateness, and acceptability of the style employed were reevaluated given that the educational and socioeconomic levels are highly variable in the population for which the instrument is intended.

Content validation

Aiming to assess the objectivity and relevance of the items of the translated instrument, its content was independently

evaluated by a panel of eight qualified assessors.¹⁶ All evaluators were nurses with considerable experience in the field of transfusion medicine with at least one year of practice in collecting blood components and complete higher education. Preference was given to professionals with doctorate or master degrees, with published papers related to the theme of blood donation. The evaluators were informed about the purposes of the study and asked to classify as valid or not each one of the items of the instrument. Thus, the content validity index (CVI) was calculated for each item.¹⁷ The items translated with a CVI of 100% were kept for the definitive instrument. The translations of items with a CVI less than 100% and greater than or equal to 80% were discussed, and the translation of items with a CVI of less than 80% were changed.

Data collection

Once the content was validated, the instrument was applied to a sample of 1,001 consecutive blood donors at the Blood Center in Franca (northeastern São Paulo State). The determination of sample size was based on statistical considerations as presented by Schmidt et al.,¹⁸ who reported the minimum number of subjects needed to validate studies.

It was considered that all individuals donating blood at the time of data collection would be able to participate in the study regardless of their literacy, since the data was collected by interviews. Therefore, the inclusion criteria for this study are the same as those in force in Brazil regarding individuals who are able to donate blood: age between 18 and 65 years, weighing more than 50 kilograms and being in good health. The exclusion criteria are naturally the following: being pregnant, donors known to have had Chagas disease, malaria (or have been in an endemic area), hepatitis after 10 years of age, leprosy, AIDS, diabetes, cancer, illicit drug use or sexual risk behavior.

Data collection was performed by four nurses with experience in blood collection for over a year. Blood donors were invited to participate voluntarily in the study after being informed of its purpose. Furthermore it was explained that all the information would be kept anonymous. Consent was established in the blood collection room, immediately after blood donation, with the donors signing an informed consent form. Despite the large sample size, no blood donor refused to participate in the study.

Shortly after the informed consent form was signed, each blood donor was interviewed and asked to complete a structured questionnaire containing questions regarding age, gender, number of previous blood donations and level of education based on the translated version of the BDRI. The interviews were conducted individually in an appropriate place so that the interviewer could read each item to the donor as well as the contents in parentheses (see Appendix A) and, if necessary, discuss the meaning of each item. However, no additional explanation was given about the items. During data collection, modification of the blood donors' routine was prevented in order to avoid affecting the flow of service.

Internal consistency

The internal consistency of the translated instrument was assessed using Cronbach's alpha coefficient.¹⁹ This coefficient is a value ranging between 0 and 1 that indicates whether the instrument items form a consistent set in order to measure the same object. The higher the coefficient, the higher the consistency.²⁰ The item-scale correlation coefficients were also calculated to evaluate the correlation between individual items and the BDRI scale. Cronbach's alpha coefficient was calculated by removing respective items from the BDRI and identifying the likely impact of the removal.²¹ When it is noted that the elimination of any of the items does not imply any significant change in the previously calculated Cronbach's alpha coefficient, it is evidence that the item is not measuring an aspect related to the assessed object. Confidence intervals (CI) for the Cronbach's alpha coefficient were obtained using the approach published by Bonett.²²

Criterion validity

The validity of criteria was achieved by comparing the scores obtained from the BDRI and the number of signs and symptoms observed by the nurse during blood donation reported in the NARD. The agreement between BDRI scores, rated as 0 or 1 point or more, and reports of signs and symptoms were measured using the kappa coefficient²³ with its respective 95% CI.

Construct validity

Construct validity was determined by examining the factor structure of the translated BDRI. Thus, an exploratory factor analysis with varimax rotation was used to value the dimensionality of the translated instrument. The suitability of a one-factor solution was assessed using scree plots (graphs not showed here) and evaluating how many factors had eigenvalues greater than 1. Kaiser's measure of sampling adequacy (MSA)^{24,25} was used to examine data adequacy for factor analysis, where values greater than 0.8 are considered good. Communalities were used to estimate the amount of variance accounted for by the factor solution for each variable.

Ethical considerations

Ethical approval was obtained from the Research Ethics Committee of the Medical School in Ribeirão Preto, Universidade de São Paulo. Informed consent was obtained from the blood donors before their entry in the study.

Statistical analysis

Statistical analyses were performed using the SAS version 9 and the R software.

Results

After the translation process, the Portuguese version of the BDRI was evaluated by the panel of evaluators and the CVI was generated for each item in order to assess the content validity. Eight items were classified as potentially valid by

all eight evaluators (CVI equal to 100%) and were included in the final inventory. Another three items received suggestions for small changes by the assessors (CVI = 87.5%), these suggestions were discussed until a definitive version of the translated BDRI was obtained.

Next, the BDRI was administered to the sample of 1,001 blood donors. Table 1 shows the characteristics of the study sample. Of the donors, 65.8% were male, 30.9% were in the age group of 21-30 years and 27.3% were first-time donors. The

final Portuguese version of the BDRI instrument is shown in Appendix A.

Table 2 shows the analysis of the internal consistency of the Portuguese version regarding the 11-, 6- and 4-item scales. The internal consistency determined by Cronbach's alpha coefficient was satisfactory for all three scales. Cronbach's alpha coefficient was estimated as 0.850 for the 11-item scale, 0.853 for the 6-item scale, and 0.781 for the 4-item scale. The alpha coefficient for deleted items showed that removing

Table 1 - Characteristics of the participants in this study.

	Total		Male		Female	
	n	%	n	%	n	%
<i>Age (completed years)</i>						
18 to 20	183	18.3	144	21.9	39	11.4
21 to 30	310	30.9	186	28.2	124	36.3
31 to 40	236	23.6	149	22.6	87	25.4
41 to 50	169	16.9	112	17.0	57	16.7
More than 50	103	10.3	68	10.3	35	10.2
<i>Education</i>						
No schooling	4	0.4	2	0.3	2	0.6
Incomplete primary school	106	10.6	70	10.6	36	10.5
Complete primary school	107	10.7	77	11.7	30	8.8
Incomplete secondary school	73	7.3	55	8.4	18	5.3
Complete secondary school	477	47.7	340	51.7	137	40.0
University education	233	23.3	114	17.3	119	34.8
<i>Systolic blood pressure (mmHg)</i>						
Less than 100	16	1.6	4	0.6	12	3.5
100 ≤ SBP ≤ 140	949	94.9	624	94.7	325	95.0
140 < SBP ≤ 160	31	3.1	26	3.9	5	1.5
More than 160	5	0.4	5	0.8	-	-
<i>Diastolic blood pressure (mmHg)</i>						
60 ≤ DBP ≤ 90	972	97.2	636	96.5	336	98.2
90 < DBP ≤ 100	29	2.8	23	3.5	6	1.8
<i>Weight (kg)</i>						
Less than 60	141	14.1	41	6.2	100	29.2
60 < weight ≤ 70	262	26.1	143	21.7	119	34.8
70 < weight ≤ 80	270	26.0	196	29.8	74	21.7
80 < weight ≤ 90	171	17.1	143	21.7	28	8.2
More than 90	157	15.7	136	20.6	21	6.1
<i>Number of blood donations^a</i>						
First-time donor	273	27.3	172	26.1	101	29.5
2	116	11.5	66	10.0	50	14.6
3	67	6.7	35	5.3	32	9.4
4	76	7.6	43	6.5	33	9.7
5	49	4.9	29	4.4	20	5.8
More than 6	420	42.0	314	47.7	106	31.0
Total	1001		659		342	

^a Including the current blood donation.

any single item would not yield a higher result. Therefore, considering all the three scales, the items are related enough to be combined into a single construct. It can also be noted in Table 2 that, when considered the 11-item scale, the correlation with total score are equal or less than 0.55 for items 4, 6, 8, 9 and 10. These items are excluded in the 6-item scale.

Table 2 - Internal consistency analysis of the Portuguese version of the Blood Donation Reactions Inventory (BDRI) for the 11, 6 and 4-item scales.

	Correlation with total score	Alpha coefficient if item deleted	Cronbach's alpha (95% CI)
11-item scale			0.850 (0.835-0.863)
1. Faintness	0.6410	0.8316	
2. Dizziness	0.7034	0.8222	
3. Weakness	0.6622	0.8269	
4. Facial flush	0.4854	0.8426	
5. Visual disturbance	0.6805	0.8274	
6. Difficulty hearing	0.5503	0.8389	
7. Lightheadedness	0.5901	0.8355	
8. Rapid or pounding heartbeat	0.3304	0.8566	
9. Sweating	0.4547	0.8496	
10. Rapid or difficult breathing	0.4630	0.8429	
11. Nausea or upset stomach	0.6006	0.8352	
6-item scale			0.853 (0.837-0.866)
1. Faintness	0.6069	0.8359	
2. Dizziness	0.7498	0.8092	
3. Weakness	0.6838	0.8278	
5. Visual disturbance	0.6876	0.8211	
7. Lightheadedness	0.5980	0.8383	
11. Nausea or upset stomach	0.6525	0.8315	
4-item scale			0.781 (0.757-0.802)
1. Faintness	0.5795	0.7452	
2. Dizziness	0.6875	0.6752	
3. Weakness	0.6531	0.7013	
7. Lightheadedness	0.5415	0.7625	
95% CI: 95% confidence interval			

The criterion validity analysis was performed comparing BDRI scores and presence or absence of signs and symptoms observed by the nurse during the donation, as reported in the NARD. Table 3 describes the results of this comparison, considering the 11, 6 and 4-item BDRI scales. All of the Kappa coefficients were relatively low suggesting a poor overall agreement between the two methods. However, only 12 blood donors had reported signs and symptoms from the

NARD, whereas a much higher number reported at least one reaction using the BDRI (243 donors considering the 11-item scale). Of the 12 donors with signs and symptoms according to the NARD, 11 (91.7%) described a reaction when the 11-item BDRI was used and 10 (83.3%) described a reaction when the 6 and 4-item scales were used (Table 3). In addition, of the 989 donors without signs and symptoms as reported by the NARD, expressive proportions did not describe any reaction in the 11, 6 and 4-item BDRI scales (respectively, 76.5%, 86.7% and 87.2%). These results show that the BDRI is more sensitive than the NARD to identify the presence of blood donation reactions. As the translated BDRI was able to identify most of the signs and symptoms reported in the NARD, the results suggest reasonable criteria validity for the Portuguese version of the BDRI despite the low kappa coefficients.

Table 3 - Comparison between reports of signs and symptoms from the Notice of Adverse Reaction during Donation (NARD) and Blood Donation Reactions Inventory (BDRI) scores rated as 0 points or 1 or more points.

BDRI scores	Points	Reports of signs and symptoms from NARD		Total	Kappa coefficient (95% CI)
		Present	Absent		
11-items	0	1 (8.3%)	757 (76.5%)	758	0.065 (0.02-0.10)
	1 or more	11 (91.7%)	232 (23.5%)	243	
6-items	0	2 (16.7%)	857 (86.7%)	859	0.110 (0.04-0.18)
	1 or more	10 (83.3%)	132 (13.3%)	142	
4-items	0	2 (16.7%)	862 (87.2%)	864	0.115 (0.04-0.18)
	1 or more	10 (83.3%)	127 (12.8%)	137	
Total		12 (100.0%)	989 (100.0%)	1001	

Table 4 presents the results of the exploratory factor analysis. Factor loadings with values greater than 0.5 indicating items that were better characterized by the factors. The 11-item scale seems to measure more than one construct as three factors have eigenvalues greater than 1. Factor 1 is better correlated to items 1, 2, 3, 5, 6, 7 and 11 (factor loadings over 0.5). Factor 2 correlates better with items 1, 4, 6 and 9, and factor 3 correlates better with items 8 and 10. Kaiser's MSA was 0.86 for the 11-point scale, which indicates that the data was well suited for factor analysis. The 6- and 4-item scales measure only one construct as only the first eigenvalue is greater than 1. In these cases, all factor loadings are positive and greater than 0.5. In all factor analyses, examination of communality magnitudes showed that the factors accounted for a considerable proportion of the variance in all items.

Discussion

Many studies have shown that the experience of adverse events such as vasovagal reactions can decrease the retention of blood donors.²⁶⁻²⁹ Thus, strategies aimed to know how blood donors react to donation and how the risks of adverse

Table 4 - Exploratory factor analysis of the Portuguese version of the Blood Donation Reactions Inventory (BDRI) for the 11-, 6- and 4-item scales.

BDRI item	Factor pattern (varimax rotation)			Communality
	Factor 1	Factor 2	Factor 3	
11-item scale				
1. Faintness	0.5282	0.5628	0.0643	0.60
2. Dizziness	0.7932	0.1778	0.2331	0.72
3. Weakness	0.7475	0.1408	0.2860	0.66
4. Facial flush	0.1128	0.7994	0.1814	0.68
5. Visual disturbance	0.6969	0.4273	0.0174	0.67
6. Difficulty hearing	0.5332	0.5495	-0.1151	0.60
7. Lightheadedness	0.6901	0.3213	-0.0058	0.58
8. Rapid or pounding heartbeat	0.0852	0.1280	0.8272	0.71
9. Sweating	0.1343	0.7120	0.2062	0.57
10. Rapid or difficult breathing	0.2442	0.1499	0.7662	0.67
11. Nausea or upset stomach	0.7728	-0.0188	0.2679	0.67
Eigenvalue	4.818	1.233	1.072	
Contribution to the total variation	43.8%	11.2%	9.8%	
Variance explained by each factor	3.409	2.142	1.573	
Kaiser's MSA = 0.860				
6-item scale				
1. Faintness	0.7258			0.53
2. Dizziness	0.8447			0.71
3. Weakness	0.7912			0.63
5. Visual disturbance	0.7924			0.63
7. Lightheadedness	0.7294			0.53
11. Nausea or upset stomach	0.7546			0.57
Eigenvalue	3.596			
Contribution to the total variation	59.9%			
Variance explained by each factor	3.596			
Kaiser's MSA = 0.834				
4-item scale				
1. Faintness	0.7640			0.58
2. Dizziness	0.8411			0.71
3. Weakness	0.8147			0.66
7. Lightheadedness	0.7330			0.54
Eigenvalue	2.492			
Contribution to the total variation	62.3%			
Variance explained by each factor	2.492			
Kaiser's MSA = 0.785				
MSA: measure of sampling adequacy.				

reactions can be minimized are of great importance to maintain the blood supply. The BDRI can be a very useful research tool about attitudes towards blood donation and to plan measures to promote satisfaction and retention of blood donors. In the present study, the BDRI was translated into Portuguese; this version is a valid and reliable instrument to collect information regarding systemic reactions experienced by blood donors.

The internal consistency analysis showed similar Cronbach's alpha coefficient estimates for the 11- and 6-item scales (Table 2). The coefficient was lower for the 4-item scale. In addition, the items with lower factor loadings in the 11-item scale are those that do not appear in the 6-item scale. These results suggest that the use of the 4-item scale may cause a loss of some information to measure a given object, whereas the synthetic 6-item scale measures the same object as good as that of 11 items. On the other hand, the decision to use the 11- or 6-item scale demands a greater understanding of the composition of each one, which is assisted by the construct validity.

Considering the 11-item scale, the results of the exploratory factor analysis shown in Table 4 suggest that Factor 1 is better correlated to items that correspond to vasovagal adverse reactions on blood donation in response to bleeding (Items 1, 2, 3, 5, 6, 7 and 11). According to the literature, these reactions occur between 1.4% and 7% of all blood donations.³⁰⁻³⁴ Factor 2 shows high factor loadings for Items 1 (faintness), 4 (facial flush), 6 (difficulty hearing) and 9 (sweating). These symptoms are related to other factors that may affect the process of blood donation such as fear of needles or fear of seeing blood. These are dynamic mental processes that can be transformed into physical symptoms. It is known that strong emotions such as fear can cause neurovegetative symptoms related to alarming situations.^{35,36} The application of psychological strategies to minimize the impact of such fears, and consequently the appearance of unpleasant symptoms in the process of blood donation, are contemporary issues in transfusion medicine studies. Regarding Factor 3, factor loadings were observed with higher values for Items 8 (rapid or pounding heartbeat) and 10 (rapid or difficult breathing). These reactions are common in excited or anxious blood donors.³⁷ Thus, considering the 11-item scale, Factors 1, 2 and 3 correspond to 'vasovagal adverse reactions', 'fear' and 'anxiety/excitation', respectively.

The 6- and 4-item scales have only one factor with an eigenvalue greater than 1. Thus, these scales are useful when the interest is to identify vasovagal symptoms related to blood donation alone, whereas the 11-item scale is useful if the aim is to assess not only vasovagal reactions, but also the donor's psychological state directly related to the occurrence of such reactions.

The results shown in Table 3 suggest that since the 11-item scale addresses signs and symptoms related not only to the physiological response, but also the psychological reactions of the individual in relation to the drawing of blood, the nurse responsible for the donation is unable to have a complete notion about the sensations experienced

by donors. Also, the 11-item scale described reactions in 91.7% of the cases of signs or symptoms recorded in the NARD, and when there was no record of a reaction, the 11-item scale identified that 76.5% of the donors reported no signs or symptoms during donation. This fact suggests that the NARD reported fewer reactions during blood donations than the BDRI, given that it is dependent on the direct observation of the nurse, whereas BDRI includes the subjective reactions of the donor. Therefore, it is noteworthy that one instrument does not replace the other as they have different purposes.

Conclusion

In conclusion, the findings suggest that the Portuguese version of the BDRI is a reliable and valid instrument to measure reactions towards blood donations. This research collaborates by providing a useful tool for research in transfusion medicine in Portuguese speaking countries.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgement

We are grateful to FAEPA (HC-FMRP) and CNPq for financial support.

REFERENCES

- Martinez EZ, Lourençon AF, Achcar JA, Zozolotto HC, Ferreira O. Uso de modelos Bayesianos de volatilidade estocástica no estudo temporal dos registros de doações de sangue. *Cad Saúde Colet.* 2010;18:597-603.
- Ludwig ST, Rodrigues AC. Doação de sangue: uma visão de marketing. *Cad Saúde Pública.* 2005;21:932-9.
- Schlumpf KS, Glynn SA, Schreiber GB, Wright DJ, Randolph Steele W, Tu Y, et al. Factors influencing donor return. *Transfusion.* 2008;48:264-72.
- Ringwald J, Zimmermann R, Eckstein R. Keys to open the door for blood donors to return. *Transfus Med Rev.* 2010;24:295-304.
- Borges VL, Martinez EZ, Bendini MH, Costa MA, Ferreira SC. Avaliação da fidedignidade de um instrumento voltado à satisfação do doador de sangue. *Rev Bras Epidemiol.* 2005;8:177-86.
- Nguyen DD, Devita DA, Hirschler NV, Murphy EL. Blood donor satisfaction and intention of future donation. *Transfusion.* 2008;48:742-8.
- Giacomini L, Lunardi Filho WD. Estratégias para fidelização de doadores de sangue voluntários e habituais. *Acta Paul Enferm.* 2010;23:65-72.
- Ownby HE, Kong F, Watanabe K, Tu Y, Nass CC. Analysis of donor return behavior. *Retrovirus Epidemiology Donor Study. Transfusion.* 1999;39:1128-35.
- Guo N, Wang J, Ness P, Yao F, Dong X, Bi X, et al; NHLBI Retrovirus Epidemiology Donor Study-II, International Component. Analysis of Chinese donors' return behavior. *Transfusion.* 2011;51:523-30.
- Thomson RA, Bethel J, Lo AY, Ownby HE, Nass CC, Williams AE. Retention of 'safe' blood donors. *The Retrovirus Epidemiology Donor Study. Transfusion.* 1998;38:359-67.
- Ogata H, Iinuma N, Nagashima K, Akabane T. Vasovagal reactions in blood donors. *Transfusion.* 1980;20:679-83.
- Trouern-Trend JJ, Cable RG, Badon SJ, Newman BH, Popovsky MA. A case-controlled multicenter study of vasovagal reactions in blood donors: influence of sex, age, donation status, weight, blood pressure, and pulse. *Transfusion.* 1999;39:316-20.
- Meade MA, France CR, Peterson LM. Predicting vasovagal reactions in volunteer blood donors. *J Psychosom Res.* 1996;40:495-501.
- France CR, Ditto B, France JL, Himawan LK. Psychometric properties of the Blood Donation Reactions Inventory: a subjective measure of presyncopal reactions to blood donation. *Transfusion.* 2008;48:1820-6.
- France CR, France JL, Roussos M, Ditto B. Mild reactions to blood donation predict a decreased likelihood of donor return. *Transfus Apher Sci.* 2004;30:17-22.
- Galdeano LE, Rossi LA. Validação de conteúdo diagnóstico: critérios para seleção de expertos. *Ciência, Cuidado e Saúde.* 2006;5:60-6.
- Polit DF, Beck CT. The Content Validity Index: are you sure you know what's being reported? Critique and recommendations. *Res Nurs Health.* 2006;29:489-97.
- Schmidt FL, Hunter JE, Urry VW. Statistical power in criterion-related validation studies. *J Appl Psychol.* 1976;61:473-85.
- Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika.* 1951;16:297-334.
- Bland JM, Altman DG. Cronbach's alpha. *BMJ.* 1997;314:572.
- Pereira JCR. Análise de dados qualitativos: estratégias metodológicas para as Ciências da Saúde, Humanas e Sociais. 3.ed. São Paulo: Editora da Universidade de São Paulo; 2001.
- Bonett DG. Sample size requirements for testing and estimating coefficient alpha. *J Educ Behav Stat.* 2002;27:335-40.
- Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas.* 1960;20:37-46.
- Kaiser HF. A Second Generation Little Jiffy. *Psychometrika.* 1970;35:401-15.
- Kaiser HF, Rice J. Little Jiffy, Mark IV. *Educ Psychol Meas.* 1974;34:111-7.
- Masser BM, White KM, Terry DJ. Beliefs underlying the intention to donate again among first-time blood donors who experience a mild adverse event. *Transfus Apher Sci.* 2013;49:278-84.
- Van Dongen A, Abraham C, Ruiters RA, Veldhuizen IJ. The influence of adverse reactions, subjective distress, and anxiety on retention of first-time blood donors. *Transfusion.* 2013;53:337-43.
- Van Dongen A, Ruiters R, Abraham C, Veldhuizen I. Predicting blood donation maintenance: the importance of planning future donations. *Transfusion.* 2014;54:821-7.

29. Veldhuizen I, Atsma F, Van Dongen A, De Kort W. Adverse reactions, psychological factors, and their effect on donor retention in men and women. *Transfusion*. 2012;52:1871-9.
30. Callahan R, Edelman EB, Smith MS, Smith JJ. Study of the incidence and characteristics of blood donor 'reactors'. *Transfusion*. 1963;3:76-82.
31. Kamel H, Tomasulo P, Bravo M, Wiltbank T, Cusick R, James RC, et al. Delayed adverse reactions to blood donation. *Transfusion*. 2010;50:556-65.
32. Lin JT, Ziegler DK, Lai CW, Bayer W. Convulsive syncope in blood donors. *Ann Neurol*. 1982;11:525-8.
33. Newman BH. Blood donor complications after whole-blood donation. *Curr Opin Hematol*. 2004;11:339-45.
34. Wiltbank TB, Giordano GF, Kamel H, Tomasulo P, Custer B. Faint and prefaint reactions in whole-blood donors: an analysis of predonation measurements and their predictive value. *Transfusion* 2008;48:1799-808.
35. Brignole M, Alboni P, Benditt D, Bergfeldt L, Blanc JJ, Bloch Thomsen PE, et al. For the Task Force on Syncope, European Society of Cardiology. Guidelines on management (diagnosis and treatment) of syncope. *Eur Heart J*. 2001;22:1256-306.
36. Meade MA, France CR, Peterson LM. Predicting vasovagal reactions in volunteer blood donors. *J Psychosom Res*. 1996;40:495-501.
37. Crocco A, D'elia D. Adverse reactions during voluntary donation of blood and/or blood components. *Blood Transfus*. 2007;5:143-52.

Appendix

The Portuguese version of the Blood Donation Reactions Inventory (BDRI).

Indique o grau em que você experimentou as seguintes sensações ao doar sangue hoje circulando um número entre 0 ('nem um pouco') e 5 ('extremadamente').

0 = nem um pouco	3 = forte	
1 = leve	4 = muito forte	
2 = moderado	5 = extremadamente	

1. Desmaio (como se estivesse prestes a desmaiar e ficar inconsciente)	0	1	2	3	4	5
2. Tontura (sensação de desequilíbrio)	0	1	2	3	4	5

3. Fraqueza	0	1	2	3	4	5
4. Rubor na face (como se o rosto estivesse quente)	0	1	2	3	4	5
5. Distúrbios visuais (tais como escurecimento da visão ou visão em túnel)	0	1	2	3	4	5
6. Dificuldade em ouvir	0	1	2	3	4	5
7. Vertigens (sensação que tudo está rodando)	0	1	2	3	4	5
8. Batimento acelerado ou forte do coração	0	1	2	3	4	5
9. Suor, sudorese, transpiração	0	1	2	3	4	5
10. Respiração rápida ou dificuldade para respirar	0	1	2	3	4	5
11. Náuseas (enjoo) ou dor de estômago	0	1	2	3	4	5