


Periprosthetic Knee Infection – Part 1: Risk Factors, Classification and Diagnosis

Infecção periprotética do joelho – Parte 1: Fatores de risco, classificação e diagnóstico

João Maurício Barretto¹ André Luiz Siqueira Campos² Nelson Hiroyuki Miyabe Ooka^{3,4}

¹ Hospital São Vicente da Gávea, Rio de Janeiro, RJ, Brazil

² Department of Orthopedics and Traumatology, Hospital Federal dos Servidores do Estado, Rio de Janeiro, RJ, Brazil

³ Department of Orthopedics and Traumatology, Hospital Universitário Pedro Ernesto, Rio de Janeiro, RJ, Brazil

⁴ Department of Orthopedics and Traumatology, Hospital Universitário Clementino Fraga Filho, Rio de Janeiro, RJ, Brazil

Address for correspondence João Maurício Barretto, PhD, Rua das Laranjeiras, 550/1001, Rio de Janeiro, RJ, 22240-006, Brasil (e-mail: joao.barretto1@gmail.com).

Rev Bras Ortop 2022;57(2):185–192.

Abstract

Keywords

- ▶ arthroplasty, replacement, knee
- ▶ surgical site infection
- ▶ risk factors
- ▶ diagnosis

Infection is one of the most feared complications in the postoperative period of knee arthroplasties. With the progressive aging of the population and the increased incidence of degenerative joint diseases, there is an exponential increase in the number of arthroplasties performed and, consequently, in the number of postoperative infections. The diagnosis of these should follow a hierarchical protocol, with well-defined criteria, which lead to diagnostic conclusion, thus guiding the most appropriate treatment. The aim of the present update article is to present the main risk factors, classifications and, mainly, to guide diagnostic investigation in an organized manner.

Resumo

Palavras-chave

- ▶ artroplastia do joelho
- ▶ infecção de sítio cirúrgico
- ▶ fatores de risco
- ▶ diagnóstico

A infecção é uma das complicações mais temidas no pós-operatório de artroplastias do joelho. Com o envelhecimento populacional progressivo e o aumento da incidência de doenças degenerativas articulares, observa-se um aumento exponencial do número de artroplastias realizadas e, conseqüentemente, do número de infecções pós-operatórias. O diagnóstico destas devem seguir um protocolo hierarquizado, com critérios bem definidos, que conduzam à conclusão diagnóstica, orientando, assim, o tratamento mais adequado. O objetivo do presente artigo de atualização é apresentar os principais fatores de risco, as classificações e, principalmente, guiar de forma organizada a investigação diagnóstica.

received
November 6, 2020
accepted
December 17, 2020

DOI <https://doi.org/10.1055/s-0041-1729935>.
ISSN 0102-3616.

© 2022. Sociedade Brasileira de Ortopedia e Traumatologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Introduction

In the last 20 years, the longevity of the world population has been increasing in developed and developing countries. This fact increases the incidence and prevalence of degenerative diseases in general, including joint diseases.^{1,2} Thus, it is natural and expected an increasing number of primary arthroplasties and revisions³ performed as treatment of these diseases.³⁻⁵

The socioeconomic impact on health systems is significant, particularly in the treatment of possible infections.⁶⁻⁸

The incidence of periprosthetic knee infections in 2001 was of 2.09% and, in 2009, it was of 2.18%, with an increasing trend.⁷ This complication is one of the main causes of rehospitalization,⁹ accounting for between 13 and 25% of the reviews performed.^{3,5,10} The estimated cost of treating

periprosthetic infection is between 3 and 4 times higher than that of primary arthroplasty.^{9,11,12}

The objective of the present work is to review the most current information on the prevention, diagnosis and treatment of periprosthetic knee infection.

Risk Factors

Risk factors for periprosthetic infection may be modifiable or nonmodifiable (► **Figure 1**).¹³

The modifiable factors most constantly found in literature and clinical practice are rheumatoid arthritis, diabetes mellitus, obesity (body mass index [BMI] > 30), corticosteroid therapy, alcoholism, smoking, and malnutrition, with hypoalbuminemia as reference.¹⁴⁻²¹

Modifiable Patient Factors	Factors with Limited Evidence of Association with ISC/IAP
<ul style="list-style-type: none"> * BMI – Strong * Smoking – Strong * Alcohol abuse – Strong * Low income – Strong * Malnutrition (hypoalbuminemia) – Source * History of DM – Strong * History of CVD – Moderate * History of ICC – Strong * History of cardiac arrhythmia – Strong * DVP Story – Strong * Chronic lung disease – Strong * Chronic obstructive pulmonary disease – Strong * History of kidney disease – Strong * History of liver disease/cirrhosis – Strong * AR History – Strong * History of cancer/malignancy – Strong * History of osteonecrosis – Strong * History of depression – Strong * History of psychosis – Strong * History of HIV/AIDS – Strong * Neurological disease (hemiplegia, paraplegia) – Moderate * History of corticosteroid use – Strong * History of intra-articular corticosteroid injection – Moderate * Prior joint surgery – Moderate * Revision arthroplasty – Strong * Previous joint infection – Moderate * Fragility – Moderate * Preoperative anemia – Strong * ASA grade >2 – Strong * Elevated Charlston Comorbidity Index – Strong * Preoperative hyperglycemia and elevated HbA1C – Moderate * Allogeneic blood transfusion – Strong * Warfarin / LMWH prophylaxis – Moderate 	<ul style="list-style-type: none"> * Age (as continuous exposure) – Limited * Hispanic ethnicity – Limited * Native Americans and Eskimo Ethnicity – Limited * Asian breed – Limited * History of drug abuse – Limited * Rural x non-rural domicile - Limited * Under body weight – Limited * History of hypertension – Limited * History of osteoarthritis – Limited * History of post-traumatic arthritis – Limited * Low and high risk dental procedures – Limited * History of UTI – Limited * History of dementia – Limited * Hypercholesterolemia – Limited * Ulcerative peptic disease – Limited * Valvular disease – Limited * Metastatic tumor – Limited * History of coagulopathy – Limited * History of venous thromboembolism – Limited * Pulmonary circulation disorders – Limited * Hypothyroidism – Limited * Hepatitis (B or C) – Limited * Hydroelectrolyte imbalance - Limited * Autologous blood transfusion – Limited
Non-Modifiable Patient Factors	
<ul style="list-style-type: none"> * Age (> 75 years) – Moderate * Male – Strong * Black race – Strong * TKA vs THA – Strong 	

Fig. 1 Risk factors for periprosthetic infection - reproduction of the International Consensus on Musculoskeletal Infections 2018 (ICM-2018).¹³

Table 1 McPherson classification for periprosthetic infection

Factor	Degree	Description
Type	I	Acute infection (< 4 weeks postoperatively)
	II	Acute hematogenous infection (< 4 weeks of symptoms)
	III	Chronic infection (> 4 weeks of symptoms)
Host factors (comorbidities and immunity)	A	Not compromised
	B	Compromised (1–2 comorbidity factors)
	C	Very compromised (> 2 comorbidity factors) or one of the factors below: <ul style="list-style-type: none"> ◦ Neutrophil count < 1,000 ◦ CD4 count < 100 ◦ Intravenous drug user ◦ Active infection elsewhere ◦ Tumor or dysplasia of the immune system
Local Factors	1	Not compromised
	2	Compromised (1–2 comorbidity factors)
	3	Very compromised (> 2 comorbidity factors)

Systemic and local factors are described in ►Table 2.

Some other clinical and social conditions are also described as associated with a higher rate of periprosthetic infection (PPI), such as preoperative American Society of Anesthesiologists (ASA) classification > 2, low income, peripheral vascular disease and others listed in ►Table 1.^{14,21,22}

Active skin lesions, either near the site or at a distance, have a potentially increased risk of periprosthetic articular infection, as well as surgery or previous joint infection.^{16,20,23}

Some studies report an increased rate of infection in total knee arthroplasty (TKA) by post-traumatic arthrosis, mainly with previous surgery and retained implants.^{24,25}

Bergen et al.²⁶ found in a comparative retrospective cohort (109 patients with implants and 109 patients without) an increase in the infection rate in patients undergoing TKA with knee implant (osteosynthesis or osteotomies). However, there was no difference when comparing the groups with previous removal of the implants ($n=43$) with those removed during the TKA procedure ($n=46$).

There is controversy regarding the increase rate of infection after TKA in patients undergoing previous joint infiltration. Some studies have shown an increased risk of infection when TKA is performed up to 3 months after joint infiltration.^{27,28} On the other hand, other studies did not find significant differences, even in short periods after infiltration (10 weeks)²⁹ or in patients submitted to multiple infiltrations.³⁰ The II-ICM-2018 (II International Consensus on Musculoskeletal Infection – 2018) suggests waiting at least 3 months after infiltration to perform arthroplasty.³¹

Regarding the nonmodifiable factors, age alone does not seem to be a predisposing factor for infection.^{15,16} Regarding gender, some studies show a higher infection rate in men than in women^{15,16} Afro descendants also have higher percentage of infection when compared with Caucasians.²¹

Komnos et al.³² retrospectively evaluated patients with arthroplasties in more than one joint. In this study, they

concluded that periprosthetic infection of a joint may predispose to hematogenous infection in another prosthetic site. The risk situations for this complication are: female gender, rheumatoid arthritis, Methicillin-Resistant *Staphylococcus aureus* (MRSA) infection and patients with fever at the time of diagnosis of the first infected joint.

Classification

Classifications are important to stratify and guide management in various clinical conditions, as well as to standardize communication between colleagues.³³

Segawa et al.,³⁴ in 1999, published a retrospective study proposing a classification and its respective treatment based on the chronology of the infection and its etiology, dividing periprosthetic infections into: positive cultures in revision perioperative harvest, acute superficial infection, acute deep infection, chronic infection, and acute hematogenous infection. However, the classification does not consider the conditions of the patient, local and systemic, or the etiological agent.³³


McPherson et al.^{35,36} described a classification system for hip and knee periprosthetic infection based on a retrospective analysis of cases evaluating three factors: type of infection (acute, acute or chronic hematogenous), host factors, and local factors (►Table 2). This classification was validated by the International Consensus on Musculoskeletal Infection with moderate evidence index and 74% of panel agreement.³⁷

Alt et al.³⁸ proposed a new classification based on the TNM classification for tumors, adapting it to periprosthetic infection, which emphasizes the pathogenicity of the etiological agent.

In the proposed classification, "T" would be tissue evaluation, "N," non-human cellular factor (etiological agent), and "M", host morbidity, according to the Charlson comorbidities classification (►Figure 2).^{38–40}

Table 2 Local and systemic factors for the McPherson classification

Factor	Description
Systemic host involvement (comorbidity or immunity)	Age >80 years old
	Alcoholism
	Dermatitis or active chronic cellulitis
	Permanent catheter
	Chronic malnutrition (albumin < 3.0 g/dL)
	Chronic nicotine use (inhaled or oral)
	Diabetes mellitus (requiring drug treatment)
	Liver failure (cirrhosis)
	Use of immunosuppressive drugs (corticosteroids, methotrexate, cyclosporine)
	Malignancy (active or history)
	Pulmonary insufficiency (SaO ₂ < 60% in room air)
	Chronic renal failure on dialysis
	Systemic inflammatory disease (Rheumatoid Arthritis, Systemic Lupus Erythematosus)
	Systemic immune impairment by infection or immunodeficiency (AIDS, acquired immunodeficiencies)
Involvement of the affected limb (wound and limb conditions)	Active infection (> 3–4 months)
	Multiple incisions – skin bridges
	Loss of soft tissue due to previous trauma
	Subcutaneous abscess (extension > 8 cm ²)
	Cutaneous synovial fistula
	Previous periarticular fracture or previous joint trauma (crushing)
	Previous local irradiation
	Peripheral vascular insufficiency – arterial or venous

**Fig. 2** TNM classification for PPI.⁴⁰

The idea of classifying these three factors seems the most appropriate approach; however, we did not find any studies validating this classification.

Diagnosis

The diagnosis of post knee arthroplasty infection has been a challenge. In the first weeks postoperatively, the occurrence of pain, local heat, and functional disability may be normal and not related to any type of bacterial infection.

Thus, it is essential to define reproducible and objective criteria that characterize the presence of infection.

Following the concept that the diagnosis of infection is often a multifactorial analysis - clinical, laboratory, imaging data and synovial fluid analysis - it is very important to hierarchize the actions in order to build this same diagnosis within a logical and progressive clinical reasoning.

We consider the strategy defined by ICM-2018 the most appropriate research option. In addition to guiding the researcher to the next step towards the diagnostic's conclusion, it brings scientific knowledge of better available evidence and the experience of hundreds of orthopedists, infectologists and microbiologists around the world. The proposed algorithm was tested and validated, presenting high sensitivity (96.9%) and specificity (99.5%) rates.^{37,41}

Following this criterion, the diagnosis of infection is defined by the presence of one of the so-called major criteria - fistula with joint communication or two positive cultures for the same microorganism identified using culture media - or by scoring clinical, serum or scoring from analysis of the synovial fluid obtained by joint puncture (► **Figure 3**).³⁷

The occurrence of fistula is found in ~ 13% of cases.³⁸ In its absence, when the patient presents with pain in the operated knee, heat and, often, decreased range of motion, it is imperative to request blood tests for evaluation of the white series, Erythrocyte Sedimentation Rate (ESR), C-reactive protein (PCR), and D-dimer.³⁷

ESR and PCR are inflammatory markers used as the first line in the screening of patients with suspected infection, with a sensitivity of ~75 88% and a specificity of 70 74%, respectively.⁴² The sensitivity of both combined ranges from 84 to 86%, and the specificity from 47 to 72.3%.^{43,44} C-reactive protein reaches its highest value on the 3rd postoperative day and remains above normal for ~ ≥ 3 weeks.⁴⁵ VHS remains elevated for at least 6 weeks.⁴⁶ Noting that, in patients with inflammatory arthritis, the cutoff value of these markers may be higher due to the influence of the underlying disease,⁴⁷ just as the use of antibiotics can generate false-negatives.⁴⁸ A recent study showed greater sensitivity (89%) and specificity (93%) for D-dimer in relation to traditional ESR and PCR.⁴³

Major Criteria (at least one of the following)			Decision
Two positive cultures of the same organism using standard culture methods			Infected
Fistula with evidence of communication with the joint or visualization of the prosthesis			

Minor Criteria	Threshold		Punctuation	Decision
	Acute [€]	Cronic		
Serum CRP (mg/L) Or D-Dimer (ug/L)	100 Unknown	10 860	2	Sum of pre- and post-operative points: ≥ 6 infected 3 to 5 inconclusive* < 3 not infected
ESR elevation (mm/hr)	Not important	30	1	
Elevated synovial leukocytes (cells/μL) Or Leukocyte Esterase Or Alpha-defensin (qualitative / quantitative)	10.000 ++ 1.0	3.000 ++ 1.0	3	
Elevated Synovial PMN (%)	90	70	2	
Only Positive Culture			2	
Positive Histology			3	
Positive Intraoperative PurulenceV			3	

[¶] This criterion has never been validated in acute infections. V No role in suspected local adverse tissue reaction.

* Consider more molecular diagnostics with next-generation sequencing.

Fig. 3 Diagnostic criteria of the Musculoskeletal Infection Society (CIIM reproduction).

Another study observed a decline in basal D-dimer levels on the second postoperative day.⁴⁶ It is important to note that ~ 2.5% of the infections do not present alterations in the aforementioned tests.³⁷

The next step following the investigation is arthrocentesis, sending the synovial fluid for laboratory analysis of cellularity (cytometry) and culture/antibiogram. There is no formal contraindication to joint aspiration.^{37,49} In this procedure, which concludes the diagnosis in 65% of the cases, it is essential that the criteria of maximum barrier to contamination are met, performed by an experienced professional with adequate packaging, and immediate shipment of the material to the laboratory.³⁷

In the acute phase, the presence in the synovial fluid of $\geq 10,000$ leukocytes/ μL , with at least 90% polymorphonuclear (PMN), and $\geq 3,000$ leukocytes/ μL in the chronic phase, with at least 70% PMN, indicate infection.³⁷

As for the culture of the aspirated liquid, some criteria should be followed to minimize the risk of false-negative. It is important to perform a prolonged time of culture, since most of the negative samples are infections by bacteria of time-consuming growth, interrupted before the appropriate time.⁵⁰

The collected joint fluid can also be used for 2 other tests: alpha-defensin and leukocyte esterase.³⁷

Alpha-defensin is an antimicrobial peptide produced by neutrophils in response to pathogens.^{37,51,52} This marker can be researched in the synovial fluid by laboratory immunoassay or by the lateral flow test, which is a rapid test with a specific kit that can be performed in the operating room with results in a few minutes. The lateral flow test presents a sensitivity rate of 78.5% and a specificity rate of 93.3%, according to a systematic review conducted by the CIIM-2018 with grouped data from 486 patients.⁴¹ Immunoassay has a sensitivity rate of 98.1% and a specificity rate of 96.4%.⁵⁴ Alpha-defensin is not influenced by recent antibiotic use, traces of blood in the sample, or comorbidities such as inflammatory diseases. The rapid test requires a small volume of synovial fluid (15 μL), which can be a great advantage in cases of absence of joint effusion.^{53,54} On the other hand, in the presence of metallomy, it may present false-negative in up to 30% of cases; it can also be influenced by crystal arthropathy (gout) and should not be done in hematoma aspirate.⁵³

Leukocyte esterase is a test with a sensitivity of 85.7% and a specificity of 94.4% according to the systematic review conducted by the CIIM-2018 with grouped data from 2,061 patients.⁴¹ This test is also not influenced by recent antibiotic use, but the presence of blood in the sample alters the readability of the test, and centrifugation may be necessary to neutralize erythrocyte interference.^{37,56}

In cases in which it is not possible to aspirate enough content for analysis (dry puncture) or whose cultures are negative, which correspond to ~ 17% of cases, intraoperative findings of pus, histological analysis, tissue culture, and new generation sequencing may help in the diagnosis of infection.⁴¹ It is not appropriate to perform joint washing in cases of dry puncture.⁴¹

Even with the entire arsenal of tests and the algorithm structured and validated, in ~ 5% of cases the diagnosis of infection cannot be confirmed.⁴¹

Some imaging tests may help in the planning of treatment but have low specificity regarding diagnosis.⁵⁷ Signs of early release on conventional radiography lead to suspected infection.⁵⁸ Computed tomography (CT) (especially arthroctomography) and magnetic resonance imaging (MRI) with metal suppression may also show signs of loosening, bone defects and, occasionally, osteomyelitis;⁵⁷ however, due to their high cost and low specificity, these tests are not recommended as diagnostic measures.^{58,59}

On the other hand, other tests have been used to differentiate aseptic loosening from infection, especially in cases of dry puncture, such as the combination of scintigraphy with marked leukocytes and single photon emission computed tomography (SPECT-CT), which also has the advantage of showing the extent of infection impairment, both in bone and soft tissue, and may be of great value in the planning of surgery.⁵⁷

Financial Support

These authors declare they have received no financial support from public, commercial, or non-profit sources.

Conflict of Interests

They have no conflict of interests to declare. Dr. Barreto reports personal fees from Stryker Latin America, outside of the submitted work.

References

- 1 Beard JR, Officer A, de Carvalho IA, et al. The World report on ageing and health: a policy framework for healthy ageing. *Lancet* 2016;387(10033):2145–2154
- 2 Cross M, Smith E, Hoy D, et al. The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014;73(07):1323–1330
- 3 Bozic KJ, Kurtz SM, Lau E, et al. The epidemiology of revision total knee arthroplasty in the United States. *Clin Orthop Relat Res* 2010;468(01):45–51
- 4 Patel A, Pavlou G, Mújica-Mota RE, Toms AD. The epidemiology of revision total knee and hip arthroplasty in England and Wales: a comparative analysis with projections for the United States. A study using the National Joint Registry dataset. *Bone Joint J* 2015; 97-B(08):1076–1081
- 5 Delanois RE, Mistry JB, Gwam CU, Mohamed NS, Choksi US, Mont MA. Current Epidemiology of Revision Total Knee Arthroplasty in the United States. *J Arthroplasty* 2017;32(09):2663–2668
- 6 Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am* 2007;89(04): 780–785
- 7 Kurtz SM, Lau E, Watson H, Schmier JK, Parvizi J. Economic burden of periprosthetic joint infection in the United States. *J Arthroplasty* 2012;27(8, Suppl):61–5.e1
- 8 Hoy DG, Smith E, Cross M, et al. The global burden of musculoskeletal conditions for 2010: an overview of methods. *Ann Rheum Dis* 2014;73(06):982–989
- 9 Kurtz SM, Lau EC, Ong KL, Adler EM, Kolisek FR, Manley MT. Which Clinical and Patient Factors Influence the National Economic Burden of Hospital Readmissions After Total Joint Arthroplasty? *Clin Orthop Relat Res* 2017;475(12):2926–2937

- 10 Evangelopoulos DS, Ahmad SS, Krismer AM, et al. Periprosthetic Infection: Major Cause of Early Failure of Primary and Revision Total Knee Arthroplasty. *J Knee Surg* 2019;32(10):941–946
- 11 Kapadia BH, Banerjee S, Cherian JJ, Bozic KJ, Mont MA. The Economic Impact of Periprosthetic Infections After Total Hip Arthroplasty at a Specialized Tertiary-Care Center. *J Arthroplasty* 2016;31(07):1422–1426
- 12 Akindolire J, Morcos MW, Marsh JD, Howard JL, Lanting BA, Vasarhelyi EM. The economic impact of periprosthetic infection in total hip arthroplasty. *Can J Surg* 2020;63(01):E52–E56
- 13 Parvizi J, Gehrke T. Anais do 2o Encontro do Consenso Internacional (ICM) em Infecção Musculoesquelética [Internet]. ICM Philly 2019 [citado 22 de março de 2020]. Disponível em: <https://icmphilly.com/document/icm-document-translation-s/icm-document-portugues-brasil-translation/>
- 14 Blanco JF, Díaz A, Melchor FR, da Casa C, Pescador D. Risk factors for periprosthetic joint infection after total knee arthroplasty. *Arch Orthop Trauma Surg* 2020;140(02):239–245
- 15 Chen J, Cui Y, Li X, et al. Risk factors for deep infection after total knee arthroplasty: a meta-analysis. *Arch Orthop Trauma Surg* 2013;133(05):675–687
- 16 Kunutsor SK, Whitehouse MR, Blom AW, Beswick ADINFORM Team. Patient-Related Risk Factors for Periprosthetic Joint Infection after Total Joint Arthroplasty: A Systematic Review and Meta-Analysis. *PLoS One* 2016;11(03):e0150866
- 17 Zhu Y, Zhang F, Chen W, Liu S, Zhang Q, Zhang Y. Risk factors for periprosthetic joint infection after total joint arthroplasty: a systematic review and meta-analysis. *J Hosp Infect* 2015;89(02):82–89
- 18 Rotevatn TA, Bøggild H, Olesen CR, et al. Alcohol consumption and the risk of postoperative mortality and morbidity after primary hip or knee arthroplasty - A register-based cohort study. *PLoS One* 2017;12(03):e0173083
- 19 Nelson CL, Elkassabany NM, Kamath AF, Liu J. Low Albumin Levels, More Than Morbid Obesity, Are Associated With Complications After TKA. *Clin Orthop Relat Res* 2015;473(10):3163–3172
- 20 Cordtz RL, Zobbe K, Højgaard P, et al. Predictors of revision, prosthetic joint infection and mortality following total hip or total knee arthroplasty in patients with rheumatoid arthritis: a nationwide cohort study using Danish healthcare registers. *Ann Rheum Dis* 2018;77(02):281–288
- 21 Zainul-Abidin S, Amanatullah DF, Anderson MB, et al. General Assembly, Prevention, Host Related General: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S13–S35
- 22 Namba RS, Inacio MC, Paxton EW. Risk factors associated with deep surgical site infections after primary total knee arthroplasty: an analysis of 56,216 knees. *J Bone Joint Surg Am* 2013;95(09):775–782
- 23 Ares O, Arnold WV, Atilla B, et al. General Assembly, Prevention, Host Related Local: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S3–S12
- 24 Bala A, Penrose CT, Seyler TM, Mather RC 3rd, Wellman SS, Bolognesi MP. Outcomes after Total Knee Arthroplasty for post-traumatic arthritis. *Knee* 2015;22(06):630–639
- 25 Aalirezaie A, Anoushiravani A, Cashman J, et al. General Assembly, Prevention, Host Risk Mitigation - Local Factors: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S37–S41
- 26 Bergen MA, Ryan SP, Hong CS, Bolognesi MP, Seyler TM. Conversion Total Knee Arthroplasty: A Distinct Surgical Procedure With Increased Resource Utilization. *J Arthroplasty* 2019;34(7S):S114–S120
- 27 Cancienne JM, Werner BC, Luetkemeyer LM, Browne JA. Does Timing of Previous Intra-Articular Steroid Injection Affect the Post-Operative Rate of Infection in Total Knee Arthroplasty? *J Arthroplasty* 2015;30(11):1879–1882
- 28 Richardson SS, Schairer WW, Sculco TP, Sculco PK. Comparison of Infection Risk with Corticosteroid or Hyaluronic Acid Injection Prior to Total Knee Arthroplasty. *J Bone Joint Surg Am* 2019;101(02):112–118
- 29 Khanuja HS, Banerjee S, Sodhi GS, Mont MA. Do Prior Intra-Articular Corticosteroid Injections or Time of Administration Increase the Risks of Subsequent Periprosthetic Joint Infections after Total Knee Arthroplasty? *J Long Term Eff Med Implants* 2016;26(03):191–197
- 30 Kokubun BA, Manista GC, Courtney PM, Kearns SM, Levine BR. Intra-Articular Knee Injections Before Total Knee Arthroplasty: Outcomes and Complication Rates. *J Arthroplasty* 2017;32(06):1798–1802
- 31 Cizmic Z, Feng JE, Huang R, et al. Hip and Knee Section, Prevention, Host Related: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S255–S270
- 32 Komnos GA, Manrique J, Goswami K, et al. Periprosthetic Joint Infection in Patients Who Have Multiple Prostheses in Place: What Should Be Done with the Silent Prosthetic Joints. *J Bone Joint Surg Am* 2020;102(13):1160–1168
- 33 Meinberg EG, Agel J, Roberts CS, Karam MD, Kellam JF. Fracture and Dislocation Classification Compendium-2018. *J Orthop Trauma* 2018;32(Suppl 1):S1–S170
- 34 Segawa H, Tsukayama DT, Kyle RF, Becker DA, Gustilo RB. Infection after total knee arthroplasty. A retrospective study of the treatment of eighty-one infections. *J Bone Joint Surg Am* 1999;81(10):1434–1445
- 35 McPherson EJ, Tontz W Jr, Patzakis M, et al. Outcome of infected total knee utilizing a staging system for prosthetic joint infection. *Am J Orthop* 1999;28(03):161–165
- 36 McPherson EJ, Woodson C, Holtom P, Roidis N, Shufelt C, Patzakis M. Periprosthetic total hip infection: outcomes using a staging system. *Clin Orthop Relat Res* 2002;403:8–15
- 37 Amanatullah D, Dennis D, Oltra EG, et al. Hip and Knee Section, Diagnosis, Definitions: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S329–S337
- 38 Alt V, Rupp M, Langer M, Baumann F, Trampuz A. Can the oncology classification system be used for prosthetic joint infection?: The PJI-TNM system *Bone Joint Res* 2020;9(02):79–81
- 39 Alt V, Rupp M, Langer M, Baumann F, Trampuz A. Infographic: Can the oncology classification system be used for prosthetic joint infection?: The PJI-TNM system *Bone Joint Res* 2020;9(02):77–78
- 40 Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40(05):373–383
- 41 Abdel Karim M, Andrawis J, Bengoa F, et al. Hip and Knee Section, Diagnosis, Algorithm: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S329–S350
- 42 Berbari E, Mabry T, Tsaras G, et al. Inflammatory blood laboratory levels as markers of prosthetic joint infection: a systematic review and meta-analysis. *J Bone Joint Surg Am* 2010;92(11):2102–2109
- 43 Shahi A, Kheir MM, Tarabichi M, Hosseinzadeh HRS, Tan TL, Parvizi J. Serum D-Dimer Test Is Promising for the Diagnosis of Periprosthetic Joint Infection and Timing of Reimplantation. *J Bone Joint Surg Am* 2017;99(17):1419–1427
- 44 Huerfano E, Bautista M, Huerfano M, Bonilla G, Llinas A. Screening for Infection Before Revision Hip Arthroplasty: A Meta-analysis of Likelihood Ratios of Erythrocyte Sedimentation Rate and Serum C-reactive Protein Levels. *J Am Acad Orthop Surg* 2017;25(12):809–817
- 45 Barretto JM, Loures FB, Albuquerque RS, Bezerra FD, Faro RV, Cavanellas NT. Evaluation of serum levels of C-reactive protein after total knee arthroplasty. *Rev Bras Ortop* 2017;52(02):176–181
- 46 Lee YS, Lee YK, Han SB, Nam CH, Parvizi J, Koo KH. Natural progress of D-dimer following total joint arthroplasty: a baseline for the diagnosis of the early postoperative infection. *J Orthop Surg Res* 2018;13(01):36

- 47 Cipriano CA, Brown NM, Michael AM, Moric M, Sporer SM, Della Valle CJ. Serum and synovial fluid analysis for diagnosing chronic periprosthetic infection in patients with inflammatory arthritis. *J Bone Joint Surg Am* 2012;94(07):594–600
- 48 Shahi A, Deirmengian C, Higuera C, et al. Premature Therapeutic Antimicrobial Treatments Can Compromise the Diagnosis of Late Periprosthetic Joint Infection. *Clin Orthop Relat Res* 2015;473(07):2244–2249
- 49 Yui JC, Preskill C, Greenlund LS. Arthrocentesis and Joint Injection in Patients Receiving Direct Oral Anticoagulants. *Mayo Clin Proc* 2017;92(08):1223–1226
- 50 Yoon HK, Cho SH, Lee DY, et al. A Review of the Literature on Culture-Negative Periprosthetic Joint Infection: Epidemiology, Diagnosis and Treatment. *Knee Surg Relat Res* 2017;29(03):155–164
- 51 Gehrke T, Lausmann C, Citak M, Bonanzinga T, Frommelt L, Zahar A. The Accuracy of the Alpha Defensin Lateral Flow Device for Diagnosis of Periprosthetic Joint Infection: Comparison with a Gold Standard. *J Bone Joint Surg Am* 2018;100(01):42–48
- 52 Frangiamore SJ, Gajewski ND, Saleh A, Farias-Kovac M, Barsoum WK, Higuera CA. α -Defensin Accuracy to Diagnose Periprosthetic Joint Infection-Best Available Test? *J Arthroplasty* 2016;31(02):456–460
- 53 Bauer TW, Bedair H, Creech JD, et al. Hip and Knee Section, Diagnosis, Laboratory Tests: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S351–S359
- 54 Shahi A, Parvizi J, Kazarian GS, et al. The Alpha-defensin Test for Periprosthetic Joint Infections Is Not Affected by Prior Antibiotic Administration. *Clin Orthop Relat Res* 2016;474(07):1610–1615
- 55 Shahi A, Alvand A, Ghanem E, Restrepo C, Parvizi J. The Leukocyte Esterase Test for Periprosthetic Joint Infection Is Not Affected by Prior Antibiotic Administration. *J Bone Joint Surg Am* 2019;101(08):739–744
- 56 Li X, Li R, Ni M, et al. Leukocyte Esterase Strip Test: A Rapid and Reliable Method for the Diagnosis of Infections in Arthroplasty. *Orthopedics* 2018;41(02):e189–e193
- 57 Diaz-Ledezma C, Espinosa-Mendoza R, Gallo J, et al. General Assembly, Diagnosis, Imaging: Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34(2S):S215–S223
- 58 Springer BD. The Diagnosis of Periprosthetic Joint Infection. *J Arthroplasty* 2015;30(06):908–911
- 59 Della Valle C, Parvizi J, Bauer TW, et al; American Academy of Orthopaedic Surgeons. American Academy of Orthopaedic Surgeons clinical practice guideline on: the diagnosis of periprosthetic joint infections of the hip and knee. *J Bone Joint Surg Am* 2011;93(14):1355–1357