



Reply to Letter to the Editor Regarding the Article: “Preoperative Prediction of Gartland IV Supracondylar Fractures of the Humerus: Is it Possible?”

Resposta à carta ao editor referente ao artigo: “Previsão Pré-Operatória de Fraturas Supracondilares de Úmero Gartland IV: É Possível?”

Sitanshu Barik¹ Gobinder Singh² Vivek Singh³

¹Department of Orthopedics, All India Institute of Medical Sciences, Deoghar, Jharkhand, India

²Orthopedics, Dayanand Medical College and Hospital, Ludhiana, Punjab, India

³Orthopedics, All India Institute of Medical Sciences, Rishikesh, Uttarakhand, India

Address for correspondence Gobinder Singh, MBBS, MS (Orthopedics), Department of Orthopedics, Dayanand Medical College and Hospital, Ludhiana, Punjab, India, 141001 (e-mail: gobindergs@gmail.com).

Rev Bras Ortop 2022;57(5):898.

To the Editor,

It was a delight to go through a critical analysis of the article by Barik et al.¹ The findings in the study are not in tune with the previously published studies on multidirectional unstable supracondylar humerus fracture.^{2,3}

The fractures are known to be created iatrogenically during intraoperative reduction of type III Gartland supracondylar humerus fractures.² It has been highlighted by Barik et al.¹ that the competency of the operating surgeon can play an important role in the creation of these type IV fractures. The discrepancy noted in the rate of neurovascular status of type IV fractures may be attributed to the sample size, compared with that of other studies. We agree that the management protocol as envisaged by Flynn et al. should be a guiding light for less-experienced surgeons, which would prevent iatrogenic conversion of type III to type IV fractures.⁴

To conclude, as advised, appropriate referral systems for these fractures should be instituted so that they are managed at specialized centers by well-trained surgeons who manage these fractures frequently.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Barik S, Singh G, Maji S, Azam MQ, Singh V. Preoperative Prediction of Gartland IV Supracondylar Fractures of Humerus: Is it Possible? Rev Bras Ortop (Sao Paulo) 2021;56(02): 230–234
- 2 Leitch KK, Kay RM, Femino JD, Tolo VT, Storer SK, Skaggs DL. Treatment of multidirectionally unstable supracondylar humeral fractures in children. A modified Gartland type-IV fracture. J Bone Joint Surg Am 2006;88(05):980–985
- 3 Mitchell SL, Sullivan BT, Ho CA, Abzug JM, Raad M, Spon-seller PD. Pediatric Gartland Type-IV Supracondylar Humeral Fractures Have Substantial Overlap with Flexion-Type Fractures. J Bone Joint Surg Am 2019;101(15): 1351–1356
- 4 Flynn JM, Skaggs DL, Waters PM. Rockwood and Wilkins': Fractures in Children. 8th. Philadelphia, PA:Wolters Kluwer;2015. [cited 2021 Oct 5]; Available from: <https://orthopaedics.lwwhealthlibrary.com/book.aspx?bookid=918>

received

October 12, 2021

accepted

November 22, 2021

published online

February 15, 2022

DOI <https://doi.org/>

10.1055/s-0042-1742601.

ISSN 0102-3616.

© 2022. Sociedade Brasileira de Ortopedia e Traumatologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil