

Table 1 Crude and adjusted factors associated with alcohol misuse among indigenous people with tuberculosis, Mato Grosso do Sul state, Brazil, 2011-2016

Variable	Crude PR (80%CI)	p-value	Adjusted PR (95%CI)	p-value
Gender				
Male	2.0 (1.5-2.7)	< 0.001	1.4 (1.1-2.0)	0.075
Female	1			
Age group				
10-24 years	1			
25-49 years	1.5 (1.1-2.1)	0.099	--	--
50 or older	0.9 (0.6-1.5)	0.842	--	--
Cash transfer programs*				
Yes	1			
No	1.7 (1.4-2.1)	< 0.001	1.5 (1.1-2.0)	0.005
Smoking (≥ 10 cigarettes/day)				
Yes	5.1 (3.8-6.7)	< 0.001	4.6 (3.0-7.1)	< 0.001
No	1		1	
Severity of tuberculosis				
Severe	1.6 (1.2-2.1)	0.046	--	--
Other	1		1	

CI = confidence interval; PR = prevalence ratio.

*Bolsa Família program or rural retirement benefits.

Maximiliano L. de Souza,¹  Jesem D. Orellana,²
Paulo C. Basta³

¹Fundação Oswaldo Cruz (FIOCRUZ) Ceará, Eusébio, CE, Brazil.

²Instituto Leônidas e Maria Deane, FIOCRUZ, Manaus, AM, Brazil.

³Escola Nacional de Saúde Pública Sergio Arouca (ENSP),
FIOCRUZ, Rio de Janeiro, RJ, Brazil.

Submitted Feb 04 2020, accepted Mar 09 2020, Epub May 11 2020.

Acknowledgements

This study received financial support from the Programa de Desenvolvimento e Inovação em Saúde, Escola Nacional de Saúde Pública Sergio Arouca (ENSP), Fundação Oswaldo Cruz (FIOCRUZ; process ENSP 013-LIV-10).

Disclosure

The authors report no conflicts of interest.

How to cite this article: de Souza ML, Orellana JD, Basta PC. Alcohol misuse by Amerindians with tuberculosis: relations to cash transfer programs in Brazil. *Braz J Psychiatry*. 2020;42:569-570. <http://dx.doi.org/10.1590/1516-4446-2020-0928>

References

- Cormier M, Schwartzman K, N'Diaye DS, Boone CE, Dos Santos AM, Gaspar J, et al. Proximate determinants of tuberculosis in Indigenous peoples worldwide: a systematic review. *Lancet Glob Health*. 2019;7:e68-e80.
- Lagarde M, Haines A, Palmer N. The impact of conditional cash transfers on health outcomes and use of health services in low and middle income countries. *Cochrane Database Syst Rev*. 2009;(4):CD008137.
- Hochberg NS, Sarkar S, Horsburgh CR Jr, Knudsen S, Pleskunas J, Sahu S, et al. Comorbidities in pulmonary tuberculosis cases in

Puducherry and Tamil Nadu, India: opportunities for intervention. *PLoS One*. 2017;12:e0183195.

4 Verdum R. Estudos etnográficos sobre o programa bolsa família entre povos indígenas.2016 [cited 2019 Aug 10]. www.academia.edu/29808772/Relat%C3%B3rio_Final_-_Estudos_Etnogr%C3%A1ficos_sobre_o_Programa_Bolsa_Fam%C3%ADlia_entre_Povos_Ind%C3%ADgenas

5 Kreter AC, Bacha CJC. Avaliação da equidade da previdência no meio rural do Brasil. *Rev Econ Sociol Rural*. 2006;44:467-502.

Brazilian research on child and adolescent suicide: looking at the past to plan the future

Braz J Psychiatry. 2020 Sep-Oct;42(5):570-572
doi:10.1590/1516-4446-2020-1024



Suicide in children and adolescents is a major public health concern across low- and middle-income and high-income countries alike, with approximately 140,000 youth aged 10-24 years dying by suicide every year worldwide.^{1,2} Although rare in absolute numbers, suicide in children and adolescents is the second or third leading cause of death among youth aged 14-24 years in many countries, including Brazil. Furthermore, the prevalence of suicide attempt and suicidal ideation in this age range is among the highest across the lifespan. Preventing suicidal attempt and ideation is a key public health issue, as they are the most important and consistently identified risk factors for suicide. Additionally, apart from the increased likelihood of suicide, suicide attempt and ideation in youth have a profound impact on young people's

lives during a sensitive period for the establishment of adult health. For example, studies have shown that adolescents who attempt suicide are more likely to report long-term socioeconomic problems, including a higher likelihood of needing welfare assistance, dropping out from high school, and being unemployed.³

In a recent issue of the *Brazilian Journal of Psychiatry*, Piccin et al.⁴ examined the research output on suicide in children and adolescents in Brazil. The authors performed a systematic search of the literature indexed in PubMed/MEDLINE from the date of database inception to December 31, 2017 and identified 146 studies meeting the inclusion criteria (i.e., studies investigating self-harm or suicide-related outcomes in individuals ≤ 19 years old and including at least 5 Brazilian participants). The description of these studies revealed several important findings on the nature of suicide research in Brazil. First, all of the identified studies can be classified as epidemiological and observational. No other methodologies, such as intervention or neurobiological studies, have been applied to date in child and adolescent populations in Brazil. Second, only a minority of the identified studies specifically investigated suicide-related outcomes in children and adolescents. Third, despite the relevance of suicide attempt and suicidal ideation in child and adolescent-focused research, more than half of the identified studies focused on suicide mortality.

Research on child and adolescent suicide in Brazil has the potential to be of great interest for an international audience and to significantly advance our understanding of such a complex phenomenon. Indeed, despite cultural variability in suicide manifestations, most of what we know about suicide in children and adolescents comes from research conducted in high-income countries, particularly in North America and Northern Europe. A systematic summary of the current state of suicide research in children and adolescents in Brazil, a middle-income country, would therefore be useful to further advance the research effort towards understanding youth suicide. Most importantly, looking at the past allows one to better plan for the future. After systematically reviewing previous work, what direction should research on suicide in Brazil follow? We believe Brazil has much potential to contribute to the field. First, observational epidemiological research is likely to continue providing important information on the epidemiology of suicide and can lead to further advancements, e.g., by exploiting high-quality longitudinal datasets such as those in the Pelotas cohorts.⁵ Refining official statistics to offer accurate representation of suicide rates and trends over time would be a key added value to epidemiological research on pediatric suicide. Second, it is increasingly clear that neurobiological factors are strongly implicated in the etiology of suicide.² Developing this field of research, perhaps relying on existing facilities and resources available in major universities and research centers, would be an asset to advance research on the biological basis of youth suicide. Third, translating current research findings into concrete actions requires a rigorous investigation of the efficacy of population-based and clinical interventions. Most evidence for the efficacy of the available preventive interventions for suicide comes from high-income

countries. Therefore, conducting trials of their efficacy in the Brazilian context is a necessary step to promote suicide prevention and to adapt interventions to the specific reality of Brazil. Finally, given the significant cultural and socioeconomic diversity of Brazilian society, a fine-grained understanding of cultural variations of suicide phenomena – including stigma, risk and protective factors, and barriers/facilitators to the implementation of preventive programs – would be highly informative for youth suicide prevention.

Just as every study, the work of Piccin et al. has some limitations. The most important one, in our view, is the absence of a narrative description of the results of included studies. Indeed, despite their systematic and rigorous literature search, the authors did not present a narrative review of the main findings and research themes. This is a missed opportunity because such a description would have been highly informative about future directions for Brazilian research in pediatric suicide. A second shortcoming is the limitation of the search to studies published in journals indexed in the MEDLINE platform. Although the authors did not restrict their search to articles published in English, widening their search to include gray literature and country-specific databases and repositories would have given a much more complete view of the research on child and adolescent suicide in Brazil. For example, works such as master's and doctoral theses often contain rigorous investigations or pilot research in understudied communities, which are highly informative at the local level even if they are not always published in indexed journals.

In spite of these shortcomings, the work by Piccin et al. is important to remind us that, while valuable information has already been produced on youth suicide in Brazil, we still have much to learn in order to address a problem that will only become more serious as Brazil develops.

Massimiliano Orri,^{1,2}  Gustavo Turecki¹

¹McGill Group for Suicide Studies, Douglas Mental Health University Institute, Department of Psychiatry, McGill University, Montreal, Canada. ²Bordeaux Population Health Research Center, Inserm U1219, University of Bordeaux, France.

Submitted Apr 10 2020, accepted Apr 14 2020, Epub Jun 01 2020.

Disclosure

The authors report no conflicts of interest.

How to cite this article: Orri M, Turecki G. Brazilian research on child and adolescent suicide: looking at the past to plan the future. *Braz J Psychiatry*. 2020;42:570-572. <http://dx.doi.org/10.1590/1516-4446-2020-1024>

References

- Hawton K, O'Connor RC. Self-harm in adolescence and future mental health. *Lancet*. 2012;379:198-9.
- Turecki G, Brent DA. Suicide and suicidal behaviour. *Lancet*. 2016;387:1227-39.
- Mars B, Heron J, Crane C, Hawton K, Lewis G, Macleod J, et al. Clinical and social outcomes of adolescent self harm: population based birth cohort study. *BMJ*. 2014;349:g5954.

- 4 Piccin J, Manfro PH, Caldieraro MA, Kieling C. The research output on child and adolescent suicide in Brazil: a systematic review of the literature. *Braz J Psychiatry*. 2020;42:209-13.
- 5 Gonçalves H, Wehrmeister FC, Assunção MC, Tovo-Rodrigues L, Oliveira IO, Murray J, et al. Cohort profile update: the 1993 Pelotas (Brazil) birth cohort follow-up at 22 years. *Int J Epidemiol*. 2018; 47:1389-90e.

Addressing interpersonal conflict among healthcare workers during the coronavirus pandemic

Braz J Psychiatry. 2020 Sep-Oct;42(5):572-573
doi:10.1590/1516-4446-2020-1109



The coronavirus outbreak is having a severe and profound socioeconomic impact worldwide. There are daily reports in the news media of healthcare services that are overwhelmed due to a lack of protective equipment and an increasing caseload of patients in need of hospitalization or intensive care. Healthcare personnel are on the front lines of the pandemic, and abundant evidence shows they are at increased risk of both infection and acute/chronic psychological distress. Greenberg et al. recently discussed moral issues that healthcare workers may perceive during the pandemic.¹ Herein, we would like to shed light to another relevant issue that healthcare teams must deal with during these challenging times.

Interpersonal conflict, a widely recognized stress factor in human relationships, can be defined as a dynamic process that arises among individuals who experience negative emotional reactions to perceived disagreement.² Although interpersonal conflict is unavoidable, it strongly impacts the quality of and perceived satisfaction with human relationships. Thus, it is unsurprising that efficient management of interpersonal conflicts in institutions and organizations has been associated with a better organizational environment and better results.³ Of course, interpersonal conflict already existed in healthcare teams to a greater or lesser degree before the pandemic emerged, and previous research has revealed its negative impact on absenteeism rates and self-esteem in nurses.⁴ Unfortunately, both new and pre-established stress factors could exacerbate interpersonal conflict between healthcare workers during the pandemic. Thus, in the current scenario, which involves increased loss of frontline healthcare workers due to contamination or contact with suspected cases, it is imperative that health organizations improve programs to positively impact their staff's mental health and mitigate potential causes of absenteeism.

Psychological First Aid and similar tools are well-established interventions to reduce or prevent psychological distress during human disasters and crisis situations. Although these methods focus on enhancing social support, they do not adequately address the complex

dimension of interpersonal conflict and associated psychological distress. Therefore, to better address this issue, we propose an ultra-brief clinical intervention model based on Interpersonal Psychotherapy (IPT). IPT is a brief structured psychotherapy intervention focused on resolving interpersonal problems.⁵ The method has been used clinically for decades and has proven efficacy in major depressive disorder.⁵ Although it has not been tested for interpersonal conflict in organizational scenarios such as healthcare teams, some specific aspects of IPT might adequately lend themselves to this goal. First, IPT uses simple language and centers primarily on emotions and feelings triggered by relationships. Second, one of the four clinical foci of IPT is very similar to interpersonal conflict, namely "role dispute." Third and most importantly, effective and learnable techniques, such as identification/expression of emotion and role playing, are used in IPT but are not specific to IPT. In fact, preliminary evidence suggests that role playing may be effective in healthcare team conflict management.⁶

This is not to say that IPT can be performed without adequate certification and training. However, in these challenging times it is essential to develop alternative means of dealing with the problems encountered in real practice, such as interpersonal conflict between health professionals, while taking good care to implement only evidence-based actions that will either help or cause no harm. Furthermore, liaison psychiatrists and hospital mental health teams could perform these interventions for their healthcare colleagues in need during the outbreak. Thus, when the usual coping techniques are insufficient to resolve interpersonal conflicts between members of healthcare teams, a clinical and minimally adapted IPT intervention could be tried as an alternative, since it is based on a psychotherapy model focused on interpersonal conflict.

Paulo Marcos Brasil Rocha,¹  Humberto Correa^{1,2}

¹Departamento de Saúde Mental, Faculdade de Medicina, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil. ²Programa de Pós-Graduação em Medicina Molecular, Faculdade de Medicina, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil.

Submitted May 05 2020, accepted May 23 2020, Epub Jun 03 2020.

Disclosure

The authors report no conflicts of interest.

How to cite this article: Rocha PMB, Correa H. Addressing interpersonal conflict among healthcare workers during the coronavirus pandemic. *Braz J Psychiatry*. 2020; 42:572-573. <http://dx.doi.org/10.1590/1516-4446-2020-1109>

References

- Greenberg N, Docherty M, Gnanapragasam S, Wessely S. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ*. 2020;368:m1211.
- Barki H, Hartwick J. Conceptualizing the construct of interpersonal conflict. *Int J Confl Manag*. 2004;15:216-44.