

Helminth fauna parasitizing *Pimelodus pohli* (Actinopterygii: Pimelodidae) from the upper São Francisco River, Brazil

Helmintofauna de *Pimelodus pohli* (Actinopterygii: Pimelodidae) do alto rio São Francisco, Brasil

Claudia Silveira São Sabas¹; Marília Carvalho Brasil-Sato^{2*}

¹Curso de Pós-graduação em Ciências Veterinárias, Departamento de Parasitologia Animal, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brasil

²Departamento de Biologia Animal, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brasil

Received July 7, 2014

Accepted August 5, 2014

Abstract

The parasite fauna of catfish, *Pimelodus pohli*, from the São Francisco River Basin is presented. A total of 45 catfish from the upper São Francisco River (45°15'44"W 18°13'25"S), were examined from July 2009 to September 2011. Forty-three catfish (95.5%) were infected by at least one parasite species, with 885 parasite specimens being found, distributed across 17 species: Monogenea (*Demidosperrmus uncusvalidus*, *Pavanelliella pavanellii*, and *Scleroductus* sp.); Eucestoda (plerocercoids of Proteocephalidea); Digenea (metacercariae of *Austrodiplostomum compactum*, adults of *Auriculostoma platense* and *Kalipharynx* sp., and juvenile of *Prosthenhystera obesa*); Nematoda (larvae of *Contraecum* sp., *Hysterothylacium* sp., *Procamallanus pimelodus*, *Procamallanus* sp., and unidentified of Cucullanidae, and adults of *Cucullanus caballeri*, *Philometra* sp., and *Procamallanus freitasi*); and Acanthocephala (adults of *Neoechinorhynchus pimelodi*). *Procamallanus freitasi* and *Scleroductus* sp. were the taxa with the highest prevalence. *Demidosperrmus uncusvalidus*, *P. freitasi*, and *Scleroductus* sp. were the dominant species. The host's sex did not influence parasitic indexes; however, the total length of the catfish did appear to have some influence. The parasites, with except for *P. obesa*, were registered for the first time in *P. pohli*, as well as the occurrence of *Kalipharynx* sp. and *C. caballeri* among pimelodid hosts from São Francisco River and South America.

Keywords: Parasites of endemic fish, helminth parasites of freshwater fish, parasite community.

Resumo

A fauna parasitária de mandis, *Pimelodus pohli* da bacia do rio São Francisco é apresentada. Um total de 45 mandis do alto rio São Francisco (45°15'44"W 18°13'25"S), foram examinados entre julho de 2009 e setembro de 2011. Quarenta e três mandis (95,5%) estavam infectados por pelo menos uma espécie de parasito tendo sido encontrados 885 parasitos de 17 espécies: Monogenea (*Demidosperrmus uncusvalidus*, *Pavanelliella pavanellii*, and *Scleroductus* sp.); Eucestoda (plerocercóides de Proteocephalidea); Digenea (metacercárias de *Austrodiplostomum compactum*, adultos de *Auriculostoma platense* e *Kalipharynx* sp. e juvenil de *Prosthenhystera obesa*); Nematoda (larvas de *Contraecum* sp., *Hysterothylacium* sp., *Procamallanus pimelodus*, *Procamallanus* sp., larvas não identificadas de Cucullanidae e adultos de *Cucullanus caballeri*, *Philometra* sp. e *Procamallanus freitasi*); e Acanthocephala (adultos de *Neoechinorhynchus pimelodi*). *Procamallanus freitasi* e *Scleroductus* sp. foram os táxons com prevalência mais elevada. *Demidosperrmus uncusvalidus*, *P. freitasi* e *Scleroductus* sp. foram as espécies dominantes. O sexo dos hospedeiros não influenciou os índices parasitários, mas houve alguma influência do comprimento total sobre eles. Os parasitos, com exceção de *P. obesa*, são registrados pela primeira vez em *P. pohli*, bem como a ocorrência de *Kalipharynx* sp. e *C. caballeri* em hospedeiros pimelodídeos do rio São Francisco e da América do Sul.

Palavras-chave: Parasitos de peixe endêmico, helmintos de peixes de água doce, comunidade parasitária.

*Corresponding author: Marília Carvalho Brasil-Sato, Departamento de Biologia Animal, Universidade Federal Rural do Rio de Janeiro – UFRRJ, BR 465, Km 7, CP 74539, CEP 23851-970, Seropédica, RJ, Brasil, e-mail: mcbsato@gmail.com

Introduction

Pimelodus pobli Ribeiro & Lucena, 2006, a fish species popularly known as white catfish, or “mandi-branco”, is found at several localities along the São Francisco River Basin (RIBEIRO; LUCENA, 2006). According to these authors, *P. pobli* is a species identified by Britski et al. (1988) as *Pimelodus* sp.. As was recently described, studies referring to its biological, behavioral, and parasitological aspects are unknown, and a comparison of its fauna with that of *Pimelodus maculatus* Lacepède, 1803 (congeneric of the same hydric system) and other pimelodids, for which its parasite faunas are already known, is urgent. Observations of the gastric content of *P. pobli* in the present study have revealed the use of aquatic invertebrates (insects, small arachnids, and crustaceans) and sediments as elements of its diet. Taking into account the study on the feeding habits of fish in the upper São Francisco River, as conducted by Alvim (1999), *P. pobli* may be considered a generalistic invertivorous and iliophagous. Thus, the endohelminthic fauna recorded in that study must reflect these feeding habits.

In the São Francisco River Basin, the records on the parasite fauna of pimelodids were established for *Conorhynchus conirostris* (Valenciennes, 1840) by Pinto et al. (1974), Pinto and Noronha (1976), Vicente et al. (1985), and Brasil-Sato and Santos (2005); for *Pseudoplatystoma corruscans* (Spix & Agassiz, 1829) by Rego and Gibson (1989), Kohn et al. (1997), and Corrêa and Brasil-Sato (2008); for *Bergiaria westermanni* (Lütken, 1874) by Moreira et al. (1991) and Vicente and Pinto (1999); for *P. maculatus* by Brasil-Sato and Pavanelli (1998, 1999, 2000, 2004), Brasil-Sato (2003), and Monteiro et al. (2010); and for *P. pobli* by Martins et al. (2012). In this last work, the authors have established the first record of a parasite species, *Prosthenhystera obesa* (Diesing, 1850), in the gall bladder of *P. pobli*.

Studies on parasitism in pimelodids have been also conducted in other localities of Brazil, such as some helminth parasites of species of *Pimelodus* Lacepède, 1803: *P. blochii* Valenciennes, 1840 in Paraná River (KOHN et al., 1997); *P. fur* (Lütken, 1874) in Paraná River (KOHN et al., 1997); *P. maculatus* in Amazon Basin (KOHN et al., 2011); in Guaíba River (KOHN; FRÓES, 1986; FORTES et al., 1993; FORTES; HOFFMANN, 1995; SARMENTO et al., 1995); in Guandu River (SANTOS et al., 2007; ALBUQUERQUE et al., 2008; AZEVEDO et al., 2010); in Itajaí-Açu River (BACHMANN et al., 2007); in Mogi Guaçu River (TRAVASSOS; KOHN, 1965; KOHN et al., 1985; KOHN; FERNANDES, 1987; AGUIAR et al., 2011); in Paraíba do Sul River (VENANCIO et al., 2010); in Paraná River (TRAVASSOS; KOHN, 1965; PAVANELLI; MACHADO DOS SANTOS, 1992; MORAVEC et al., 1993, 1997; KOHN et al., 1997; BRASIL-SATO; PAVANELLI, 2004; COHEN; KOHN, 2008; KOHN et al., 2011; LOPES et al., 2011); in Paranapanema River (RAMOS et al., 2013); *Pimelodus microstoma* Steindachner, 1877 in Mogi Guaçu River (AGUIAR et al., 2011) [note of the authors: this fish species was cited by the authors as *P. heraldoi* Azpelicueta, 2001, that is a junior synonym of *P. microstoma* Stein., *sensu* Ribeiro and Lucena (2010)]; *Pimelodus ornatus* Kner, 1858 in Paraná River (PAVANELLI; TAKEMOTO, 1996; PAVANELLI et al., 1997;

FERNANDES; KOHN, 2001; KOHN et al., 2011); *Pimelodus* sp. in Amazon Basin (VICENTE et al., 1978). Through a literature review, it was possible to verify at least 131 species of helminths recorded among the pimelodids distributed in the Brazilian basins.

In this study, the helminth fauna of *P. pobli* is presented for the first time in the São Francisco River Basin, contributing to the knowledge on the parasite species of fish in the São Francisco River, and on the parasite species of pimelodids in the hydrographic basins in South America.

Materials and Methods

A total of 45 specimens of *P. pobli* were collected from July 2009 to September 2011 in the upper São Francisco River, downstream from the Três Marias Reservoir (45°15'44"W, 18°13'25"S) in the municipality of Três Marias, State of Minas Gerais, Brazil. The fish were captured by local fishers and were sent to the Centro Integrado de Recursos Pesqueiros e Aquicultura (CIRPA) of the Companhia de Desenvolvimento dos Vales do São Francisco e do Parnaíba (CODEVASF), where they were weighted (average body weight: 26.6 ± 11 g), measured (average total length: 14.5 ± 1.6 cm), and necropsied to investigate their parasite fauna. Voucher specimens of *P. pobli* were deposited in the Coleção de Peixes of the Museu de Zoologia of the Universidade de São Paulo (MZUSP: 105895), State of São Paulo, Brazil. Parasite voucher specimens were deposited in the Coleção Helminológica of the Instituto Oswaldo Cruz (CHIOC), Rio de Janeiro, Brazil, under the numbers indicated in Table 1 of the Results section.

Ecological descriptors (prevalence, mean abundance, and mean intensity) were calculated according to Bush et al. (1997). The dispersion index was calculated for each species in order to determine the distribution standard of the species; it was tested by calculating the *d* statistic (LUDWIG; REYNOLDS, 1988). The interaction between couples of species co-occurring within the hosts was evaluated via the chi-square test (χ^2) with Yates's correction and a 2 × 2 contingency table, and by Spearman's rank correlation coefficient (*r_s*). Student's *t*-test was used to evaluate the total length among the male and female hosts. Spearman's rank correlation coefficient (*r_s*) was used to verify possible correlations between the total length of hosts and parasite abundance and intensity. Pearson's correlation coefficient (*r*) was used to determine possible correlations between the total length of the hosts and parasite prevalence. The influence of the host's sex on parasite abundance and prevalence was tested using the normal approximation of *Z* via the Mann–Whitney *U* test and the chi-square test (χ^2) with Yates's correction and a 2 × 2 contingency table, respectively. The statistical analysis was performed according to Zar (1996), and was applied to the study of parasite infracommunities among *P. pobli*, but only for species with a prevalence rate higher than 10% (BUSH et al., 1990). The adopted significance level was *p* < 0.05.

Results

Among the 45 examined specimens of *P. pobli*, 18 (40%) were male and 27 (60%) were female (representing a ratio of 1:1.5). Male and female catfish presented with a mean total length of

Table 1. Ecological descriptors (prevalence, mean intensity, intensity amplitude, mean abundance), and site of infestation/infection of helminth parasites (the developmental stages of the parasite specimens, except for adults, were indicated in parentheses following the nominal taxa) of *Pimelodus pohli* Ribeiro & Lucena, 2006 from the upper São Francisco River, State of Minas Gerais, Brazil.

Parasite	Prevalence (%)	Mean intensity	Intensity amplitude	Mean abundance	Site
Monogenea					
<i>Demidospermus uncusvalidus</i> Gutierrez & Suriano, 1992 (CHIOC n.37922, 37923)	51.1	17.22±19.73	1-78	8.80±16.44	Gill filaments
<i>Pavanelliella pavanellii</i> Kritsky & Boeger, 1998 (CHIOC n.37921)	4.4	1	1	0.04±0.21	Nasal cavity
<i>Scleroductus</i> sp.(CHIOC n.37924)	62.2	4.86±5.76	1-23	3.02±5.10	Body surface
Digenea					
<i>Austrodiplostomum compactum</i> (Lutz, 1928) (metacercariae) (CHIOC n.37927)	35.5	2.25±2.57	1-10	0.80±1.85	Eyes
<i>Auriculostoma platense</i> (Szidat,1954) (CHIOC n.37926)	6.6	19.67±31.47	1-56	1.31±8.34	Anterior intestine
<i>Prosthenhystera obesa</i> (Diesing, 1850) (juvenile) (CHIOC n.37732)	2.2	1	1	0.02±0.15	Gall bladder
<i>Kalipharynx</i> sp.	2.2	3	1-3	0.07±0.45	Coelom
Cestoda					
Proteocephalidea (plerocercoids) (CHIOC n.37928)	6.7	2.70±2.10	1-5	0.18±0.80	Intestine
Nematoda					
<i>Contraecaecum</i> sp. (larvae) (CHIOC n.35922)	17.8	3.00±2.14	1-6	0.53±1.44	Coelom
<i>Cucullanus caballeri</i> Petter, 1977 (CHIOC n.35915, 35916)	6.7	1	1	0.02±0.15	Anterior and posterior intestine
Cucullanidae (larvae unidentified) (CHIOC n.35914)	4.4	1	1	0.04±0.21	Coelom
<i>Hysterothylacium</i> sp. (larvae) (CHIOC n.35917)	2.2	1	1	0.02±0.15	Coelom
<i>Philometra</i> sp. (CHIOC n.35923)	2.2	1	1	0.02±0.15	Coelom
<i>Procamallanus freitasi</i> (Moreira, Oliveira & Costa, 1991) (CHIOC n.35920, 35921)	64.4	5.55±5.78	1-24	3.58±5.34	Anterior, middle and posterior intestine
<i>Procamallanus pimelodus</i> (Pinto, Fábio, Noronha & Rolas, 1974) (larvae) (CHIOC n.35918)	6.7	1.33±0.58	1-2	0.09±0.36	Middle intestine
<i>Procamallanus (Spirocamallanus)</i> sp. (larvae) (CHIOC n.35919)	24.4	3.82 ± 3.65	1-11	0.93± 2.41	Anterior and middle intestine
Acanthocephala					
<i>Neoechinorhynchus pimelodi</i> Brasil-Sato & Pavanelli, 1998 (CHIOC n. 37925)	11.6	1.20±0.44	1-2	0.13±0.40	Anterior intestine

14.3 ± 1.4 cm and 14.7 ± 1.7 cm, respectively, and a mean body weight of 25.6 ± 9.8 g and 28.2 ± 13.2 g, respectively. There was no significant difference in the total length ($t = 0.84$; $p = 0.20$) and weight ($t = 0.75$; $p = 0.45$) between male and female hosts.

Among the examined hosts, 43 (95.5%) were infected by at least one parasite species. A total of 885 parasites were found, and those represented the following taxa: Monogenea (three species); Eucestoda (one species); Digenea (four species); Nematoda (eight species); and Acanthocephala (one species) (Table 1).

Nematoda and Monogenea were the taxa whose prevalence values were higher, representing 82% and 75%, respectively, of the total fauna.

In Nematoda, the adult specimens of *Procamallanus freitasi* (Moreira, Oliveira & Costa, 1991) presented with a higher prevalence rate, mean abundance, and mean intensity, followed by *Procamallanus (Spirocamallanus)* sp. and *Contraecaecum* sp., respectively, which consisted of specimens at the larval stage. In Monogenea, *Scleroductus* sp. presented with the highest prevalence rate, and *Demidospermus uncusvalidus* Gutiérrez & Suriano, 1992,

was the most abundant species with the highest mean intensity (Table 1).

The parasite community of *P. pohli* demonstrated the standard aggregated distribution pattern, except for the uniform distribution of *Neoechinorhynchus pimelodi* Brasil-Sato & Pavanelli, 1998. The dominance frequency and mean relative dominance of the parasite community were most marked in *D. uncusvalidus*, followed by *P. freitasi* and *Scleroductus* sp.. The frequency of shared dominance was highest in *P. freitasi* and *Scleroductus* sp., respectively (Table 2).

There was no correlation between the host's sex and the parasitic indexes prevalence and abundance (*D. uncusvalidus*: $\chi^2 = 0.18$, $Z = -0.81$; *Scleroductus* sp.: $\chi^2 = 1.14$, $Z = -0.75$; *A. compactum* $\chi^2 = 0.33$, $Z = -0.65$; *Contraecaecum* sp.: $\chi^2 = 0.06$, $Z = -0.08$; *P. freitasi*: $\chi^2 = 0.01$, $Z = -0.20$; *Procamallanus* (S.) sp.: $\chi^2 = 0.49$, $Z = -0.35$; *N. pimelodi*: $\chi^2 = 0.64$, $Z = 0.99$, respectively, $p > 0.05$ for each result).

The total length of the hosts negatively influenced the parasitic indexes of the parasite fauna of *P. pohli* – that is, the intensity and abundance of *D. uncusvalidus*, the abundance of *Contraecaecum*

sp., and the prevalence of *P. freitasi*, were highest in fish with a smaller total length (Table 3).

Among the co-occurring species, the couples *D. uncusvalidus* - *Scleroductus* sp. and *P. freitasi* - *N. pimelodi* presented their correlated abundances (Table 4).

Discussion

Nematodes, both at their larval stage and as adults, are common parasites of freshwater fish, and they can attack all organs in the

body; many of them are known to be disease agents in fish and humans (MORAVEC, 1998). Nematoda was the prevailing taxon in several studies on the helminth fauna of *P. maculatus*. In Brazil, Kohn and Fernandes (1987) recorded the occurrence of *Cucullanus pinnai* Travassos, Artigas & Pereira, 1928, as the most prevailing parasite in the Mogi Guaçu River, State of São Paulo, followed by *Cucullanus* sp. and *Procamallanus pimelodus* (Pinto, Fabio, Noronha & Rolas, 1974); Bachmann et al. (2007) recorded the highest prevalence of camallanids (*P. pimelodus*) in the Itajaí-Açu River, State of Santa Catarina. The highest prevalence was noted among cucullanids (*C. pinnai*) by Santos et al. (2007) and

Table 2. Values of dominance frequency, mean relative dominance, shared dominance frequency, dispersion index, and *d* statistic of helminth parasites of *Pimelodus pohli* Ribeiro & Lucena, 2006 from the upper São Francisco River, State of Minas Gerais, Brazil.

Parasite	Dominance Frequency	Mean relative-dominance	Shared dominance frequency	Dispersion Index	<i>d</i> Statistic
Monogenea					
<i>Demidospermus uncusvalidus</i>	17	0.26±0.29	12	30.72	41.61*
<i>Scleroductus</i> sp.	6	0.20±0.24	22	8.62	17.15*
Digenea					
<i>Austrodiplostomum compactum</i>	3	0.07±0.14	13	4.29	9.06*
Nematoda					
<i>Contracaecum</i> sp.	2	0.05±0.17	5	3.88	8.10*
<i>Procamallanus freitasi</i>	7	0.21±0.26	22	7.97	16.10*
<i>Procamallanus (Spirocamallanus)</i> sp.	1	0.04±0.11	7	6.2	12.98*
Acanthocephala					
<i>Neoechinorhynchus pimelodi</i>	0	0.01±0.03	4	1.22	0.01

*Significant values: $d > 1.96$

Table 3. Influence of host length in prevalence (Pearson's (*r*) correlation coefficient), intensity, and abundance (Spearman's rank (*rs*) correlation coefficient) of helminth parasites of *Pimelodus pohli* Ribeiro & Lucena, 2006 from the upper São Francisco River, State of Minas Gerais, Brazil.

Parasite	Prevalence		Intensity		Abundance	
	<i>r</i>	<i>p</i>	<i>rs</i>	<i>p</i>	<i>rs</i>	<i>p</i>
Monogenea						
<i>Demidospermus uncusvalidus</i>	-0.96	0.18	-0.42	0.04*	-0.38	0.009*
<i>Scleroductus</i> sp.	-0.88	0.25	-0.29	0.13	0.08	0.6
Digenea						
<i>Austrodiplostomum compactum</i>	0.97	0.12	0.24	0.36	0.21	0.15
Nematoda						
<i>Contracaecum</i> sp.	-0.30	0.8	-0.24	0.56	-0.33	0.02*
<i>Procamallanus freitasi</i>	-0.99	0.02*	0.04	0.83	-0.14	0.34
<i>Procamallanus (Spirocamallanus)</i> sp.	-0.55	0.62	-0.05	0.87	-0.16	0.29
Acanthocephala						
<i>Neoechinorhynchus pimelodi</i>	0.97	0.12	0.35	0.51	-0.44	0.77

*significant values: $p < 0.05$.

Table 4. Couples of co-occurring species of helminth parasites of *Pimelodus pohli* Ribeiro & Lucena, 2006 from the upper São Francisco River, State of Minas Gerais, Brazil

Co-occurring Species	Prevalence		Abundance	
	χ^2	<i>p</i>	<i>rs</i>	<i>p</i>
Ectoparasites (adults)				
<i>Demidospermus uncusvalidus</i> - <i>Scleroductus</i> sp.	2.74	0.09	0.34	0.02*
Endoparasites (adults)				
<i>Procamallanus freitasi</i> - <i>Neoechinorhynchus pimelodi</i>	0.72	0.39	0.35	0.02*
Endoparasites (larvae)				
<i>Contracaecum</i> sp. - <i>Procamallanus (Spirocamallanus)</i> sp.	0.19	0.66	0.27	0.07

(Chi-square test (χ^2) with Yates's correction; Spearman's rank (*rs*) correlation coefficient, *significant values: $p < 0.05$).

Albuquerque et al. (2008) in the Guandu River, State of Rio de Janeiro; Venancio et al. (2010) found that *C. pinnai* was the sole helminth in the fauna, presenting with a high prevalence rate in this host in the Paraíba do Sul River, State of Rio de Janeiro. In addition to the occurrence of these nematodes in high indexes in several hydrographic basins, *Cucullanus* spp. [especially *C. pinnai* (cucullanids) and *Procamallanus* spp. (camallanids)] have been constantly presented in the parasite fauna of *P. maculatus*. In this study, *P. freitasi* was the species that presented the highest prevalence among the endoparasites of *P. pohli*, with remarkable record of larvae of *Procamallanus* sp. and *P. pimelodus*, as well as adults of *Cucullanus caballeroi* Petter, 1977; this pattern was relevant, as it showed the same structural composition of the representative cucullanid and camallanid species in the helminth fauna of congeneric pimelodid hosts. This parasite–host relationship had been cited for pimelodid hosts in the Mogi Guaçu (KOHN; FERNANDES, 1987), Paraná (PAVANELLI et al., 1997), and São Francisco Rivers (BRASIL-SATO, 2003) when, in addition to the Eucestoda larvae, Nematoda was highlighted as a common parasite that was represented by the larvae of the species allotted to Anisakidae, the species of *Procamallanus* Baylis, 1923, *Cucullanus* Müller, 1777 and, among the latter, *C. pinnai* (BRASIL-SATO, 2003).

In addition to the ecological aspects that involve the use of alimentary items that predispose congeneric hosts to the same types of parasites, and which lead to the parasite–host specificity in several hydrographic basins listed in this study, it is equally significant to add that *P. freitasi* indexes were significantly higher in smaller specimens of *P. pohli* – an endemic species to the São Francisco River.

Procamallanus spp. perforate the intestinal wall of hosts with their buccal capsule and feed on blood; these nematodes are commonly found upon necropsy, and are red or dark brown in color due to such hematophagous activity (BRASIL-SATO, 2003). They can cause inflammatory reactions at the place where they adhere to, and they may cause anemia in the host due to blood loss. They can occupy the intestinal lumen in small or juvenile fish and, if they do not cause death under such conditions, they can affect growth (THATCHER, 2006). Besides the availability of the intermediate host (crustaceans and insects), which justifies the constant record of *C. pinnai* in *Pimelodus* spp., Albuquerque et al. (2008) have also suggested the possibility of a histotrophic phase in *C. pinnai* cycle. As most Brazilian rivers are related to reservoirs with hydroelectric potential, the abundance of nematodes may be enhanced, i.e. the parasitic transmission can be faster and/or more efficient among the hosts kept in smaller areas (e.g., due to fragmentation of the rivers and also due changing the flow of the rivers). Host abundance is favored in lentic environments, such as in lakes and dams (MORAVEC, 1998), and thus high indexes of infection by nematodes are found in the fish species that dwell in areas close to the reservoirs of hydroelectric power stations (FELTRAN et al., 2004). Therefore, all of these aspects must be considered in proposals that suggest raising *Pimelodus* spp. under confined conditions, and a lot of attention must be paid to parasitism in endemic species, such as *P. pohli* (endemic to the São Francisco River), when dealing with biodiversity preservation in aquatic systems.

The development stage and the parasitic indexes of *P. freitasi* and *C. caballeroi* have suggested that *P. pohli* serves as the definitive host for those nematodes, and as an intermediate host for the larvae of *Contracaecum* sp. and *Hysterothylacium* sp., which will become adult specimens in birds, according to Moravec (1998).

Santos et al. (2007) and Gutiérrez and Martorelli (1999) recorded high prevalence values for *D. uncusvalidus* and low prevalence rates for *Scleroductus* sp. in *P. maculatus* in the Guandu River, Brazil and in the la Plata River, Argentina, respectively. In the present study, the prevalence of *Scleroductus* sp. and the prevalence and mean intensity of *D. uncusvalidus* were high. In the upper São Francisco River, these two direct-transmission ectoparasite species (THATCHER, 2006) are the ones that characterized the ectoparasite community of *P. pohli* (mainly in fish with smaller body sizes), and the high values of their parasitic indexes optimized the observed co-occurrence.

The biometric data of siluriform hosts have been analyzed and were related to the parasitic indexes in several works (MACHADO et al., 1994; BRASIL-SATO; PAVANELLI, 2004; BRASIL-SATO; SANTOS, 2005; MOREIRA et al., 2005; SANTOS; BRASIL-SATO, 2006; BACHMANN et al., 2007). In the São Francisco River, parasitic indexes have been minimally influenced by fish size, and the possible cumulative effect of parasites, as proposed by Rohde (1993), has not been observed in parasite communities of fish in that hydric system. In this study, smaller fish presented with a higher abundance of ectoparasites (*D. uncusvalidus*) and, except for *A. compactum*, endohelminths also had their abundance favored, with the number of larval specimens of *Contracaecum* sp. and adults of *P. freitasi* being significantly higher in smaller *P. pohli* (Table 4). For camallanid nematodes, it is possible that the smaller fish presented with predilection for some alimentary items such as copepods, which were indicated as intermediate hosts for these parasites by Moravec (1998).

The sex of siluriform hosts has not been a determining factor in the structure of fish in the São Francisco River. In the present study, and in the studies conducted by Brasil-Sato and Santos (2005) and Santos and Brasil-Sato (2004, 2006), in the São Francisco River, there was no influence of host sex on the parasitic indexes, and this fact was supported by the similarities in feeding strategies and habits observed between male and female hosts.

Sympatric or related hosts must be similar in composition to their parasite faunas due to the level of similarity in their biological and behavioral features. In the São Francisco River, Pinto et al. (1974), Pinto and Noronha (1976), and Vicente et al. (1985) reported on the larvae of *Procamallanus* (*Spirocamallanus*) sp. in *Conorhynchos conirostris*, and Moreira et al. (1991) and Vicente and Pinto (1999) recorded *P. freitasi* in *B. westermanni*. Brasil-Sato (2003) recorded the following helminths in *P. maculatus*: *Demidospermus* sp., *Vancleaveus* sp., *Pavanelliella pavanellii* Kritsky & Boeger, 1998; *Monticellia loyolai* Pavanelli & Machado dos Santos, 1992 and *Nomimoscolex* sp. – two proteocephalid cestode species, one them presenting with larval specimens; *Auriculostoma platense* (Szidat, 1954) Scholz, Aguirre-Macedo and Choudhury, 2004 (cited as the original combination: *Crepidostomum platense* Szidat, 1954), *Creptotrema creptotrema* Travassos, Artigas & Pereira, 1928, *Sanguinicola coelomica* (Szidat, 1951) Gibson and Bray, 2002 (cited as the original combination: *Plehnella coelomica* Szidat,

1951), *Prosthenhystera obesa* (Diesing, 1850), *Thometrema overstreeeti* Brooks, Mayers & Thorson, 1979, a digenean species identified in this study (*Kalipharynx* sp.), metacercariae of *Clinostomum* sp. and *Diplostomum* sp. (= *Austrodiplostomum* sp.); *C. pinnai*, *Dichelyne* sp., *Philometra* sp., *P. freitasi*, larval specimens of unidentified nematodes; and *N. pimelodi*. Larvae of *Procamallanus* (*Spirocamallanus*) sp. were recorded by Brasil-Sato and Santos (2005) in *C. conirostris*. Thus, among the species already recorded in pimelodids in the São Francisco Basin, the following parasites were found in *P. pobli* in this study: *P. pavanellii*, plerocercoids of Proteocephalidea, *A. platense*, *P. obesa*, *Kalipharynx* sp., *Philometra* sp., *P. freitasi*, larvae of *Contraecaecum* sp., and *N. pimelodi*.

In other localities, some these common helminths species were also found in pimelodids. Regarding the Monogenea, in the Baía (type-locality) and Paraná Rivers, *Pavanelliella pavanellii* was described of *P. corruscans* by Kritsky and Boeger (1998) and they also reported this monogenean species of *Callophysus macropterus* (Lichtenstein, 1819) from Solimões River; *P. pavanellii* was recorded by Lopes et al. (2009) from Negro and Solimões Rivers in *Pseudoplatystoma tigrinum* (Spix & Agassiz, 1829) (*sic*) and *Pseudoplatystoma punctifer* (Linnaeus, 1766) (*sic*) [note of the authors: *P. tigrinum* as cited by the authors is *P. tigrinum* (Valenciennes, 1840), and *P. punctifer* refers to *P. punctifer* (Castelnau, 1855) *sensu* Buitrago-Suárez and Burr (2007)]. In the Guandu River, Santos et al. (2007) found *Demidospermus uncusvalidus* Gutiérrez & Suriano, 1992 and *Scleroductus* sp.; and in the Mogi Guaçu River, Aguiar et al. (2011) recorded *P. pavanellii* of *P. maculatus*. With respect to the Digenea, in the Guaíba River, *A. platense* (as *C. platense*) was found parasitizing *P. maculatus* by Kohn and Fróes (1986) and Fortes et al. (1993), respectively; in the Paraná Basin, the six Digenea species cited in *P. maculatus* from São Francisco River were common in the fauna of *P. maculatus* and their indexes were recorded and compared by Brasil-Sato and Pavanelli (2004); and in the Guandu River, Santos et al. (2007) and Azevedo et al. (2010), recorded *A. compactum* parasitizing *P. maculatus*. With relation to the Nematodes, in the Paraná Basin, Machado et al., (1994, 1995, 1996) recorded *Contraecaecum* sp. and *Procamallanus* (S.) sp. of *P. corruscans*; and Guidelli et al. (2003) recorded larvae of *Contraecaecum* sp. in *Hemisorubim platyrhynchos* (Valenciennes, 1840); and In the Aquidauana River, Campos et al. (2008) recorded larvae of *Contraecaecum* sp. in *Pseudoplatystoma fasciatum* (L., 1766). Among the helminths recorded in *P. pobli*, *N. pimelodi* – the sole acanthocephalan species – was described from the parasite specimens of the congeneric *P. maculatus* and was recorded by Santos and Brasil-Sato (2006) in *Franciscodoras marmoratus* Reinhardt, 1874 (Siluriformes, Doradidae) in the São Francisco River (type-locality). Recently, Lopes et al. (2011) recorded *N. pimelodi* in *P. maculatus* and *B. westermanni* in the Paraná Basin. In the parasite community of *P. pobli*, *N. pimelodi* was the only one with a uniform distribution, but in the congeneric host *P. maculatus*, its distribution was widely dispersed, both during periods of flooding and drought in the upper São Francisco River region (BRASIL-SATO; PAVANELLI, 1999).

In this study, it is suggested that the co-occurrence of endoparasites *P. freitasi* and *N. pimelodi* was due to the use of arthropods, intermediate hosts, in the diet of *P. pobli*.

The significance of the analysis of parasitic indexes among host species with a wide distribution in the South American hydrographic basins, and among congeneric and sympatric species in this region, is highlighted to detect the main groups of parasite species that are able to affect fish. This type of predictive analysis indicates the relative specificity of parasite–host relationships in the hydrographic basins. The parasitism analysis of *P. pobli* highlighted that the smaller specimens were significantly more infested by Monogenea and infected by Nematoda; *P. pobli* is an endemic pimelodid host, whose niche and immunological plasticity is related to the available habitats that is restricted to the São Francisco River in relation to congeneric species or other pimelodids that present with a wide geographic distribution in the South and Central Americas. It was reinforced that, among the endoparasites, cucullianid and camallanid nematodes are the parasite representatives that prevail in the parasite fauna of the *Pimelodus* species. For *P. pobli*, the presence and parasitic indexes of those nematodes (as well as the endoparasite community) prove its invertivorous alimentary habits, which were not previously known.

In this study, the parasite species of *P. pobli* were presented, as well as those that are common in the fauna of *P. maculatus* in the São Francisco River and other basins. With the exception of *P. obesa*, which was recorded in *P. pobli* by Martins et al. (2012), this study revealed the parasite community of *P. pobli* and highlighted the first record of *Kalipharynx* sp. and *C. caballeroi* in the parasite community of fish in the upper São Francisco River, and in the pimelodids from South America.

Acknowledgments

The authors are grateful to Dr Yoshimi Sato (CODEVASF, Três Marias, MG) for the resources and scientific information available about the host and to the working arrangement CEMIG-GT/CODEVASF for providing logistical and material support; to Dr Osvaldo Takeshi Oyakawa and to Dr Marcelo Knoff for his kindness in receiving the voucher specimens sent for deposit at the MZUSP and CHIOC, respectively. To CAPES for scholarship to first author and for funded this research (Edital 032/2010 – Parasitologia Básica).

References

- Aguiar JC, Ceccarelli PS, Luque JL. Two new species of *Pavanelliella* (Monogenea, Dactylogyridae) parasitic on pimelodid fishes from Mogi Guaçu river, southeastern Brazil, and notes on the morphology of *P. pavanellii*. *Neotrop Helminthol* 2011; 5(2): 213-224.
- Albuquerque MC, Santos MD, Monteiro CM, Martins AN, Ederli NB, Brasil-Sato MC. Helminths endoparasitos de *Pimelodus maculatus* Lacépède, 1803 (Siluriformes, Pimelodidae) de duas localidades (lagoa e calha) do Rio Guandu, Estado do Rio de Janeiro, Brasil. *Rev Bras Parasitol Vet* 2008; 17(S1): 113-119. PMID:20059829.
- Alvim MCC. *Composição e alimentação da ictiofauna em um trecho ao alto rio São Francisco, município de Três Marias - MG* [Dissertação]. São Carlos: Universidade Federal de São Carlos; 1999.
- Azevedo RK, Abdallah VD, Luque JL. Acanthocephala, Annelida, Arthropoda, Myxozoa, Nematoda and Platyhelminthes parasites of

- fishes from the Guandu river, Rio de Janeiro, Brazil. *Check List* 2010; 6(4): 659-667.
- Bachmann F, Greinert JA, Bertelli PW, Filho HHS, Lara NOT, Ghiraldelli L, et al. Parasitofauna de *Pimelodus maculatus* (Osteichthyes: Pimelodidae) do rio Itajaí-Açu em Blumenau, Estado de Santa Catarina, Brasil. *Acta Sci Biol Sci* 2007; 29(1): 109-114.
- Brasil-Sato MC. Parasitos de peixes da bacia do São Francisco. In: Godinho HP, Godinho AL. *Águas, peixes e pescadores do São Francisco das Minas Gerais*. Belo Horizonte: PucMinas; 2003. p. 149-166.
- Brasil-Sato MC, Pavanelli GC. *Neoechinorhynchus pimelodi* sp. n. (Eoacanthocephala, Neoechinorhynchidae) parasitizing *Pimelodus maculatus* Lacépède, “mandi-amarelo” (Siluroidei, Pimelodidae) from the basin of the São Francisco river, Três Marias, Minas Gerais, Brazil. *Rev Bras Zool* 1998; 15(4): 1003-1011. <http://dx.doi.org/10.1590/S0101-81751998000400019>
- Brasil-Sato MC, Pavanelli GC. Ecological and reproductive aspects of *Neoechinorhynchus pimelodi* Brasil-Sato & Pavanelli (Eoacanthocephala, Neoechinorhynchidae) of *Pimelodus maculatus* Lacépède (Siluroidei, Pimelodidae) of the São Francisco river, Brasil. *Rev Bras Zool* 1999; 16(1): 73-82. <http://dx.doi.org/10.1590/S0101-81751999000100002>
- Brasil-Sato MC, Pavanelli GC. *Pavanelliella pavanellii* Kritsky e Boeger, 1998 (Monogenea: Dactylogyridae) parasito das cavidades nasais de *Pimelodus maculatus* Lac., 1803, “mandi”, das bacias do rio São Francisco e do rio Paraná, Brasil. *Parasitol. dia* 2000; 24(3-4): 123-126.
- Brasil-Sato MC, Pavanelli GC. Digenea de *Pimelodus maculatus* (Osteichthyes, Pimelodidae) das bacias dos rios São Francisco e Paraná, Brasil. *Parasitol Latinoam* 2004; 59(3-4): 123-131.
- Brasil-Sato MC, Santos MD. Metazoan parasites of *Conorhynchus conirostris* (Valenciennes, 1840), an endemic siluriform fish of the São Francisco Basin, Brazil. *Rev Bras Parasitol Vet* 2005; 14(4): 160-166.
- Britski HA, Sato Y, Rosa ABS. *Manual de identificação de peixes da região de Três Marias*: com chaves de identificação para os peixes da bacia do São Francisco. 3. ed. Brasília: Camara dos Deputados, Codevasf; 1988.
- Buitrago-Suárez UA, Burr BM. Taxonomy of the catfish genus *Pseudoplatystoma* Bleeker (Siluriformes: Pimelodidae) with recognition of eight species. *Zootaxa* 2007; 1512: 1-38.
- Bush AO, Aho JM, Kennedy CR. Ecological versus phylogenetic determinants of helminth parasite community richness. *Evolut Ecol* 1990; 4(1): 1-20. <http://dx.doi.org/10.1007/BF02270711>
- Bush AO, Lafferty KD, Lotz JM, Shostak AW. Parasitology meets ecology on its own terms: Margolis et al. revisited. *J Parasitol* 1997; 83(4): 575-583. PMID:9267395. <http://dx.doi.org/10.2307/3284227>
- Campos CM, Fonseca VE, Takemoto RM, Moraes FR. Fauna parasitária de cachara *Pseudoplatystoma fasciatum* (Siluriforme: Pimelodidae) do rio Aquidauana, Pantanal Sul Mato-Grossense, Brasil. *Acta Sci Biol Sci* 2008; 30(1): 91-96.
- Cohen SC, Kohn A. New data on species of *Demidospermus* (Dactylogyridae: Monogenea) parasitizing fishes from the reservoir of the Itaipu hydroelectric power station, Paraná State, Brazil, with new synonymies. *Rev Bras Parasitol Vet* 2008; 17(3): 167-170. <http://dx.doi.org/10.1590/S1984-29612008000300011>
- Corrêa RFSS, Brasil-Sato MC. Digenea in the surubim *Pseudoplatystoma corruscans* (Spix & Agassiz, 1829) (Siluriformes: Pimelodidae) of the upper São Francisco River, State of Minas Gerais, Brazil. *Braz Arch Biol Technol* 2008; 51(6): 1181-1185. <http://dx.doi.org/10.1590/S1516-89132008000600013>
- Feltran RB, Marçal O Jr, Pinese JF, Takemoto RM. Prevalência, abundância, intensidade e amplitude de infecção de nematóides intestinais em *Leporinus frederici* (Bloch, 1794) e *L. obtusidens* (Valenciennes, 1836) (Pisces, Anostomidae), na represa de Nova Ponte (Perdizes, MG). *Rev Bras Zool* 2004; 6(2): 169-179.
- Fernandes BMM, Kohn A. On some trematodes parasites of fishes from Paraná river. *Braz J Biol* 2001; 61(3): 461-466. <http://dx.doi.org/10.1590/S1519-69842001000300016>
- Fortes E, Hoffmann RP. 7 Levantamento da fauna parasitária de peixes do Lago Guaíba, Porto Alegre, Rio Grande do Sul, Brasil. *Rev Bras Med Vet* 1995; 17(3): 107-111.
- Fortes E, Hoffmann RP, Sarmiento, TM. *Cucullanus fabregasi* sp. n. Nematoda parasita do intestino do peixe *Pimelodus maculatus* Lacépède, 1803 do Rio Guaíba, Porto Alegre, Rio Grande do Sul, Brasil. *Arq Fac Vet UFRGS* 1993; 21: 71-76.
- Guidelli GM, Isaac A, Takemoto RM, Pavanelli GC. Endoparasite infracommunities of *Hemisorubim platyrhynchos* (Valenciennes, 1840) (Pisces: Pimelodidae) of the baía river, upper Paraná River floodplain, Brazil: specific composition and ecological aspects. *Braz J Biol* 2003; 63(2): 261-268. PMID:14509848. <http://dx.doi.org/10.1590/S1519-69842003000200011>
- Gutiérrez PA, Martorelli SR. The structure of the monogenean community on the gills of *Pimelodus maculatus* in Río de la Plata (Argentina). *Parasitology* 1999; 119(2): 177-182. <http://dx.doi.org/10.1017/S0031182099004606>
- Kohn A, Fernandes BMM. Estudo comparativo dos helmintos parasitos de peixes do Rio Mogi Guassu, coletados nas excursões realizadas entre 1927 e 1985. *Mem Inst Oswaldo Cruz* 1987; 82(4): 483-500. PMID:3507917. <http://dx.doi.org/10.1590/S0074-02761987000400006>
- Kohn A, Frões OM. *Saccocoeloides godoyi* n. sp. (Haploporidae) and other trematodes parasites of fishes from the Guaíba estuary, RS, Brazil. *Mem Inst Oswaldo Cruz* 1986; 81(1): 67-72. <http://dx.doi.org/10.1590/S0074-02761986000100009>
- Kohn A, Fernandes BMM, Macedo B, Abramson B. Helminths parasites of freshwater fishes from Pirassununga, SP, Brazil. *Mem Inst Oswaldo Cruz* 1985; 80(3): 327-336. <http://dx.doi.org/10.1590/S0074-02761985000300009>
- Kohn A, Fernandes BMM, Baptista-Farias MFD. Redescription of *Prosthenbystera obesa* (Diesing, 1850) (Callodistomidae, Digenea) with new host records and data on morphological variability. *Mem Inst Oswaldo Cruz* 1997; 92(2): 171-179. <http://dx.doi.org/10.1590/S0074-02761997000200008>
- Kohn A, Moravec F, Cohen SC, Canzi C, Takemoto RM, Fernandes BMM. Helminths of freshwater fishes in the reservoir of the hydroelectric power station of Itaipu, Paraná, Brazil. *Check List* 2011; 7(5): 681-690.
- Kritsky DC, Boeger WA. Neotropical Monogenoidea. 35. *Pavanelliella pavanellii*, a new genus and species (Dactylogyridae, Ancyrocephalinae) from the nasal cavities of Siluriform fishes in Brazil. *J Helminthol Soc Wash* 1998; 65(2): 160-163.
- Lopes LPC, Varella AMB, Malta JCO. Metazoan parasites of *Pseudoplatystoma punctifer* (Linnaeus, 1766) and *Pseudoplatystoma tigrinum* (Spix & Agassiz, 1829) Siluriformes: Pimelodidae) of the Central Amazon Basin, Brazil. *Biol Geral Exper* 2009; 9(2): 3-15.
- Lopes MS, Fernandes BMM, Bastos OM, Cohen SC, Kohn A. New hosts for two species of Acanthocephala of fishes from Paraná River, State of Paraná, Brazil. *Rev Bras Zool* 2011; 13(1-2-3): 29-32.
- Ludwig JA, Reynolds JF. *Statistical ecology: a primer in methods and computing*. New York: Wiley-Interscience; 1988.
- Machado MH, Pavanelli GC, Takemoto RM. Influence of host's sex and size on endoparasitic infrapopulations of *Pseudoplatystoma corruscans* and

- Schizodon borelli* (Osteichthyes) of the high Paraná river, Brazil. *Rev Bras Parasitol Vet* 1994; 3(2): 143-148.
- Machado MH, Pavanelli GC, Takemoto RM. Influence of the type of environment and the hydrological level variation in endoparasitic infrapopulations of *Pseudoplatystoma corruscans* (Agassiz) and *Schizodon borelli* (Boulenger) (Osteichthyes) of the high Paraná River, Brazil. *Rev Bras Zool* 1995; 12(4): 961-976. <http://dx.doi.org/10.1590/S0101-81751995000400023>
- Machado MH, Pavanelli GC, Takemoto RM. Structure and diversity of endoparasitic infracommunities and the trophic level of *Pseudoplatystoma corruscans* and *Schizodon borelli* (Osteichthyes) of the High Paraná River. *Mem Inst Oswaldo Cruz* 1996; 91(4): 441-448. PMID:9070406. <http://dx.doi.org/10.1590/S0074-02761996000400010>
- Martins AN, Sabas CSS, Brasil-Sato MC. *Prosthenhystera obesa* (Diesing, 1850) (Digenea, Callodistomidae) in the São Francisco River Basin, Brazil: New host records and their ecological parameters. *Neotrop Helminthol* 2012; 6(1): 31-41.
- Monteiro CM, Kritsky DC, Brasil-Sato MC. Neotropical Monogenoidea. 55. Dactylogyrids parasitising the pintado-amarelo *Pimelodus maculatus* Lacépède (Actinopterygii: Pimelodidae) from the Rio São Francisco, Brazil. *Syst Parasitol* 2010; 76(3): 179-190. PMID:20532849. <http://dx.doi.org/10.1007/s11230-010-9250-2>
- Moravec F. *Nematodes of freshwater fishes of the Neotropical Region*. Praha: Academia; 1998.
- Moravec F, Kohn A, Fernandes BMM. Nematode parasites of fishes of the Paraná river, Brazil. Part 3. Camallanoidea and Dracunculoidea. *Folia Parasitol* 1993; 40(3): 211-229.
- Moravec F, Kohn A, Fernandes BMM. New observations on seuratoid nematodes parasitic in fishes of the Paraná River, Brasil. *Folia Parasitol* 1997; 44(3): 209-223.
- Moreira NIB, Oliveira CL, Costa HMA. A new helminth parasite of fish: *Spirocammallanus freitasi* sp. n. (Nematoda-Camallanidae). *Mem Inst Oswaldo Cruz* 1991; 86(4): 429-431. <http://dx.doi.org/10.1590/S0074-02761991000400009>
- Moreira ST, Ito KF, Takemoto RM, Pavanelli GC. Ecological aspects of the parasites of *Iberingichthys labrosus* (Lütken, 1874) (Siluriformes: Pimelodidae) in reservoirs of Paraná basin and upper Paraná floodplain, Brazil. *Acta Sci Biol Sci* 2005; 27(4): 317-322.
- Pavanelli GC, Machado dos Santos MH. *Goezeella agostinhoi* sp. n. e *Monticellia loyolai* sp. n., cestóides proteocefalídeos parasitas de peixes pimelodídeos do rio Paraná, Paraná, Brasil. *Rev Bras Parasitol Vet* 1992; 1(1): 45-50.
- Pavanelli GC, Takemoto RM. *Spasskyellina mandi* n. sp. (Proteocephalidea: Monticelliidae), parasite of *Pimelodus ornatus* Kner, 1857 (Pisces: Pimelodidae) of the Paraná River, Paraná, Brazil. *Mem Inst Oswaldo Cruz* 1996; 91(6): 723-726. PMID:9283653. <http://dx.doi.org/10.1590/S0074-02761996000600013>
- Pavanelli GC, Machado MH, Takemoto RM. Fauna helmíntica de peixes do rio Paraná, região de Porto Rico, Paraná. In: Vazzoler AEAM, Agostinho AA, Hahn NS. *A planície de inundação do alto rio Paraná: aspectos físicos, biológicos e socioeconômicos*. Maringá: EDUEM; 1997. p. 307-329.
- Pinto RM, Noronha D. *Procamallanus* brasileiros (Nematoda, Camallanoidea): considerações finais, com chave para determinação das espécies. *Mem Inst Oswaldo Cruz* 1976; 74(3-4): 323-339. <http://dx.doi.org/10.1590/S0074-02761976000300011>
- Pinto RM, Fábio SP, Noronha D, Rolas FJT. *Procamallanus* brasileiros. Parte I (Nematoda, Camallanoidea). *Mem Inst Oswaldo Cruz* 1974; 72(3-4): 205-211. <http://dx.doi.org/10.1590/S0074-02761974000200005>
- Ramos IP, Franceschini L, Zago AC, Zica EOP, Wunderlich AC, Carvalho ED, et al. New host records and a checklist of fishes infected with *Austrodroplostomum compactum* (Digenea: Diplostomidae) in Brazil. *Rev Bras Parasitol Vet* 2013; 22(4): 511-518. PMID:24473875. <http://dx.doi.org/10.1590/S1984-29612013000400010>
- Rego AA, Gibson DI. Hyperparasitism by helminths: new records of cestodes and nematodes in proteocephalid cestodes from South American siluriform fishes. *Mem Inst Oswaldo Cruz* 1989; 84(3): 371-376. <http://dx.doi.org/10.1590/S0074-02761989000300012>
- Ribeiro FRV, Lucena CAS. A new species of *Pimelodus* LaCépède, 1803 (Siluriformes: Pimelodidae) from the rio São Francisco drainage, Brazil. *Neotrop Ichthyol* 2006; 4(4): 411-418. <http://dx.doi.org/10.1590/S1679-62252006000400004>
- Ribeiro FRV, Lucena CAS. *Pimelodus heraldoi* Azpelicueta, 2001, a junior synonym of *Pimelodus microstoma* Steindachner, 1877 (Siluriformes: Pimelodidae). *Neotrop Ichthyol* 2010; 8(2): 277-281. <http://dx.doi.org/10.1590/S1679-62252010000200005>
- Rohde K. *Ecology of marine parasites: an introduction to marine parasitology*. 2nd ed. Wallingford: CAB International; 1993.
- Santos MD, Brasil-Sato MC. Parasitos metazoários de *Franciscodoras marmoratus* (Reinhardt, 1874), "serrudo" (Siluriformes: Doradidae) do Rio São Francisco, Brasil. *Rev Bras Parasitol Vet* 2004; 13(1): 18-22.
- Santos MD, Brasil-Sato MC. Parasitic community of *Franciscodoras marmoratus* (Reinhardt, 1874) (Pisces: Siluriformes, Doradidae) from the upper São Francisco river, Brazil. *Braz J Biol* 2006; 66(3): 931-938. PMID:17119841. <http://dx.doi.org/10.1590/S1519-69842006000500019>
- Santos MD, Lemos-Pita SRLC, Brasil-Sato MC. Metazoan parasite fauna of *Pimelodus maculatus* La Cépède, 1803 (Siluriformes, Pimelodidae) from the Guandu River, Rio de Janeiro State, Brazil. *Acta Sci Biol Sci* 2007; 29(1): 101-107.
- Sarmento TM, Fortes E, Hoffmann RP. *Cucullanus debacoi* sp. n. Nematoda, parasito do intestino do *Pimelodus maculatus* Lacépède, 1803 (Pisces, Pimelodidae) do rio Guaíba, Porto Alegre, Rio Grande do Sul. *A Hora Vet* 1995; 15(85): 38-40.
- Thatcher VE. *Amazon fish parasites*. 2nd. ed. Moscow: Pensoft; 2006.
- Travassos LP, Kohn A. Lista dos helmintos parasitos de peixes encontrados na Estação Experimental de Biologia e Piscicultura de Emas, Pirassununga, Estado de São Paulo. *Papéis Avulsos Zool* 1965; 17: 35-52.
- Venancio ACP, Aguiar GR, Lopes OS, Alves DR. Metazoan parasites of mandi-amarelo *Pimelodus maculatus* and of jundiá *Rhamdia quelen* (Osteichthyes: Siluriformes) of Paraíba do Sul river, Volta Redonda, Rio de Janeiro. *Rev Bras Parasitol Vet* 2010; 19(3): 157-163. PMID:20943019. <http://dx.doi.org/10.1590/S1984-29612010000300006>
- Vicente JJ, Pinto RM. Nematóides do Brasil. Nematóides de peixes: atualização 1985-1998. *Rev Bras Zool* 1999; 16(3): 561-610. <http://dx.doi.org/10.1590/S0101-81751999000300001>
- Vicente JJ, Rodrigues HO, Gomes DC. Nematóides do Brasil. 1ª parte: nematóides de peixes. *Atas Soc Biol Rio de Janeiro* 1985; 25: 1-79.
- Vicente JJ, Santos E, Souza SV. Helmintos de peixes de rios amazônicos da coleção helmintológica do Instituto Oswaldo Cruz. I. Trematoda. *Atas Soc Biol Rio de Janeiro* 1978; 19: 9-15.
- Zar JH. *Biostatistical analysis*. 3rd ed. New Jersey: Prentice-Hall; 1996.