

Soil genesis and iron nodules in a karst environment of the Apodi Plateau¹

Gênese de solos e nódulos de ferro em ambiente cárstico na Chapada do Apodi

Rodrigo de Oliveira Girão^{2*}, Leo Jakson da Silva Moreira³, Ana Leônia de Araújo Girão², Ricardo Espíndola Romero² e Tiago Osório Ferreira⁴

ABSTRACT - The Apodi Plateau, located in the northeastern part of the state of Ceará in Brazil, has limestone as its parent material, with significant variation in the soils formed in this region, as well as the presence of iron nodules, having been identified. The iron oxides have been used as indicators for evaluating the weathering processes of some soils, with the nodules being sources of information about pedogenic processes. Seeking to evaluate the influence of the karstic parent material and of the iron nodules on soil genesis in the Apodi Plateau, the morphological, physical and chemical properties of five profiles were evaluated, as well as of the iron nodules present in some horizons. The profiles were classified as RED-YELLOW ARGISOL Abruptic eutrophic plinthosol (P1), Tb HAPLIC CAMBISOL Eutrophic latosol (P2), RED-YELLOW ARGISOL Abruptic eutrophic plinthosol (P3), Tb HAPLIC CAMBISOL Eutrophic latosol (P4) and RED ARGISOL Eutrophic nitosol (P5). Changes in chemical, physical and morphological characteristics were observed at depth in some profiles, suggesting variation in the characteristics of the parent material. Nodules of class C1 (0.053 to 0.25 mm) and C2 (0.25 to 2.00 mm) were concentrated on the surface, alternating with a sub-surface concentration of class C3 (2.00 to 4.76 mm), C4 (4.76 to 7.90 mm) and C5 (7.90 to 19.1 mm). Profile P5, considered as more evolved, showed a predominance of classes C1, C2 and C3 in the surface horizons, indicating degradation of the larger-diameter nodules with the advancing weathering process.

Key words: Pedogenic processes. Limestone. Soil classification.

RESUMO - A Chapada do Apodi, localizada na porção nordeste do Ceará, possui o calcário como material de origem e tem-se identificado relevante variação nos solos formados nessa região, bem como a presença de nódulos de ferro. Os óxidos de ferro têm sido usados como indicadores para avaliação dos processos de intemperismo de alguns solos, e esses nódulos são fontes de informações sobre processos pedogenéticos. Buscando avaliar a influência do material de origem cárstico e dos nódulos de ferro na gênese dos solos da Chapada do Apodi, foram avaliadas as propriedades morfológicas, físicas, e químicas de cinco perfis, bem como de nódulos de ferro presentes em alguns horizontes. Os perfis foram classificados como ARGISSOLO VERMELHO-AMARELO Eutrófico abruptico plintossólicos (P1), CAMBISSOLO HÁPLICO Tb Eutrófico latossólico (P2), ARGISSOLO VERMELHO AMARELO Eutrófico abruptico plintossólicos (P3), CAMBISSOLO HÁPLICO Tb Eutrófico latossólico (P4) e ARGISSOLO VERMELHO Eutrófico nitossólico (P5). Foram observadas alterações de características químicas, físicas e morfológicas em profundidade de alguns perfis, sugerindo uma variação nas características do material de origem. Os nódulos das classes C1 (0,053 - 0,25 mm) e C2 (0,25 - 2 mm) concentraram-se superficialmente, alternando para uma concentração sub-superficial das classes C3 (2 - 4,76 mm), C4 (4,76 - 7,90 mm) e C5 (7,90 - 19,1 mm). O perfil considerado mais evoluído (P5) apresentou predomínio das classes C1, C2 e C3 nos horizontes mais superficiais, indicando degradação dos nódulos de maior diâmetro com o avançar do processo de intemperismo.

Palavras-chave: Processos pedogenéticos. Calcário. Classificação de solos.

*Autor para correspondência

¹Recebido para publicação em 28/08/2013; aprovado em 26/06/2014

Parte da Dissertação de Mestrado em Solos e Nutrição de Plantas apresentada à Universidade Federal do Ceará desenvolvida com recursos do projeto financiado pelo Banco do Nordeste, intitulado "Atributos dos Solos Relacionados às Condições Paleo-Climáticas da Chapada do Apodi-CE"

²Departamento de Ciências do Solo, Universidade Federal do Ceará/UFC, Campus do Pici, Av. Mister Hull, 2977, Bloco 807, Campus do Pici, Fortaleza-CE, Brasil, 60.021-970, rogiarao@gmail.com, analeonia@yahoo.com.br, rromero@ufc.br

³Departamento de Solos, Universidade Federal de Viçosa/UFV, Campus UFV, Viçosa-MG, Brasil, 36.570-000, leopontocom2004@hotmail.com

⁴Departamento de Ciência do Solo, Escola Superior de Agricultura Luiz de Queiroz/ESALQ, Universidade de São Paulo, Avenida Pádua Dias, 11, Vila Independência, Piracicaba-SP, Brasil, toferreira@usp.br

INTRODUCTION

Iron oxides can be used as pedogenic indicators as they make possible a better understanding of the evolution of the processes of weathering, being sensitive to variations in the environment, mainly to changes in the redox conditions of the soil, and so allow us to infer changes in the past forms of landscapes (COELHO; VIDAL-TORRADO, 2003; INDA-JUNIOR; KÄMPF, 2003).

The segregation and accumulation of iron can take place in many ways, being generally related to mottling, nodules, concretions and petroplinthite, among other classifications (CONSTANTINI; PRIORI, 2007). The formation of iron accumulated through nodules is influenced by seasonal variations in the water table (TAN *et al.*, 2006), where materials having diffuse internal structures are formed by the continuous supply of iron. These materials can provide information about very common pedogenic processes in the soils of tropical and subtropical regions (CONSTANTINI *et al.*, 2006; HUANG *et al.*, 2008).

Colour is an important aspect of soils derived from limestone, and is used as one of the defining elements of the classification 'Terra rossa', employed for soils which have their origins in calcareous materials, with a clayey texture and reddish colours of varying hues ranging from 5YR to 10R, and which are found in the Mediterranean region (AYDINALP; FITZPATRICK, 2009), being related to soils influenced by such iron oxides as hematite or goethite. Soils with similar characteristics, developed in other areas and also originating from calcareous materials have been documented, however under semi-arid climates (KHORMALI *et al.*, 2003).

In Brazil, studies can be found related to soils which developed on calcareous materials with an occurrence of iron oxides; these generally being clayey soils of a reddish colour (5YR) and of oxide mineralogy (hematite and goethite), kaolinitic and micaceous. Lynch (2009) found Chernosols, Nitosols and Latosols in the Planaltina region of the state of Goiás. Oliveira *et al.* (2000) analysed Cambisols, Red Argisols and Red Latosols developed on limestone in the north of the state of Minas Gerais, and highlighted the presence of nodules and concretions of iron-manganese in the Cambisol profile. Pereira *et al.* (2013) carried out evaluations on the Serra da Bodoquena, in Mato Grosso do Sul, finding Organosols, Chernosols and Gleysols. Running further evaluations on the same region of Mato Grosso do Sul, Silva *et al.* (2013) found Red Argisols and Haplic Cambisols and stressed polygenesis as having strongly influencing this variation.

The Apodi plateau, located in the northeast of the state of Ceará in Brazil, is a fruit-producing region of importance in the state, and in soil surveys carried out in the 70s, large areas of Cambisols (BRASIL, 1973) were identified. Some more recent studies carried out in the

region however, have shown variations in soil class and the presence of iron nodules (ALENCAR, 2002; MOTA *et al.*, 2007), albeit with few detailed evaluations as to what factors may be contributing to these changes. Thus, with the aim of assessing the influence of the source material and of the iron nodules on the genesis of soils of the Apodi Plateau, morphological, chemical and physical analyses were carried out in five profiles formed from limestone.

MATERIAL AND METHODS

Area under study


The Apodi Plateau is inserted in the Potiguar Basin and is part of the Rift System of the Brazilian northeast. The emerged part of the Potiguar Basin displays the Acu and Jandaíra rock formations, the first consisting of red and green, clayey, micaceous and kaolinitic fine-grained sandstones and of siltstones of mostly fluvial origin, while the Jandaíra formation is characterised by the presence of calcite and dolomite, fine to medium grain limestones, which may be inter-bedded with cross-stratified sandy horizons (BRASIL, 1981; MOTA *et al.*, 2008).

The area under study is of approximately 10 ha, planted with banana, and located in the town of Limoeiro do Norte, Ceará. The climate in the region according to the Köppen classification is BSw'h' (hot and semiarid), with an average annual temperature and precipitation of 28.5 °C and 772 mm respectively. The dominant topography of the region is flat, with a declivity ranging from 0.005 to 0.015 m⁻¹ and an approximate elevation of 150 m.

Sampling and Analysis

A transect of approximately 300 metres was established, where five trenches were opened (Figure 1), described and collected as per Santos *et al.* (2005). The area is of flat terrain with a declivity of around 0.01 m⁻¹, the typical vegetation is hyper-xerophilic caatinga, and the small distances between profiles do not suggest variations in the climate. Samples from each horizon were broken up and passed through a 2 mm sieve in order to obtain fine, air-dried soil (FADS), in accordance with Empresa Brasileira de Pesquisa Agropecuária (1997). The soil profiles were classified according to Empresa Brasileira de Pesquisa Agropecuária (2013).

Granulometric composition was determined by the pipette method. The pH was determined in water and in a 1 mol L⁻¹ KCl solution in the ratio of 1:2.5. The organic carbon was obtained using the method proposed by Yeomans and Bremner (1988). The Ca²⁺ and Mg²⁺ cations and exchangeable acidity (Al³⁺) were

Figure 1 - Representation of profiles P1, P2, P3, P4 and P5, located in the area under study, the Apodi Plateau, Ceara, Brazil

extracted with a 1 mol L⁻¹ KCl solution and determined by titration. The exchangeable Na⁺ and K⁺ were analysed by photometric method in a soil extract obtained with a Mehlich-1 solution. The potential acidity was determined by the volumetric method, with extraction by 0.5 mol L⁻¹ calcium acetate at a pH of 7.0.

The levels of Fe in the fine-soil fraction were determined by atomic-absorption spectrophotometry after extraction with a solution of sodium dithionite-citrate-bicarbonate (DCB), in accordance with Mehra and Jackson (1960), and with Tamm's solution in the absence of light, according to Urley and Drees (2008). From the Fe content, extracted by ammonium oxalate (Fe_o) and dithionite-citrate-bicarbonate (Fe_d), the Fe_o/Fe_d and Fe_d/clay ratios were calculated, and used as a qualitative index of the degree of crystallinity of the oxides (INDA-JUNIOR, KÄMPF, 2005).

Using methodology proposed by Empresa Brasileira de Pesquisa Agropecuária (1997), the values for SiO₂, Fe₂O₃, Al₂O₃, MnO₂ and TiO₂ were quantified by inductively coupled plasma optical emission spectroscopy (ICP OES).

Iron Nodules

The nodules of iron present in some horizons were separated from the soil matrix with the use of water-jet sieves and were then grouped into five diameter classes (C1 - 0.053 to 0.25, C2 - 0.25 to 2.00, C3 - 2.00 to 4.76, C4 - 4.76 to 7.90 and C5 - 7.9 to 19.10 mm).

After separation of the nodules the relative mass, density of the nodules was determined by the volumetric flask method (EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA, 1997) together with the levels of iron extracted by the ammonium oxalate (Fe_o) and dithionite-citrate-bicarbonate (Fe_d) (MEHRA; JACKSON, 1960; URLEY; DREES, 2008).

RESULTS AND DISCUSSION

Genesis and Morphology

The soils have depths exceeding 150 cm and display well-developed A and B horizons, thus indicating an evolving pedogenesis; with the exception of profile P3 there was no contact with the parent material of these soils. In general, the soils are of red and reddish colourations, indicating the presence of iron oxides, mainly hematite.

The soils of the Apodi Plateau were generally classified as Cambisols (BRASIL, 1973). However, on the transect defined in the study area, two profiles were classified as Cambisols, two soils identified as red-yellow Argisols and one as a red Argisol. Given the negligible changes in the other factors of formation, this would suggest that changes in source material are influencing this variability, also been reported by other authors (ACHURRA *et al.*, 2009; ALENCAR, 2002; MOTA *et al.*, 2007).

The occurrence of variations in limestone rocks reported by Bautista *et al.* (2011), where different layers may vary from centimetres to a few metres, can be considered as a cause of the variation seen in profile depth, such as in profile P2 (4.3+ m) and profile P3 (1.9 m) (Table 1), separated by a distance of 70 m.

The prevailing hue of the profiles was 2.5 YR (red) (Table 1). These reddish colours in soils developed on a karst environment are strongly related to a rapid precipitation of iron oxides due to the high pH, which among other factors, favours the precipitation of ferrihydrate, a precursor of hematite, instead of goethite (SILVA NETO *et al.*, 2008; TARDY; NAHON, 1985).

In profiles P2 and P3, vertic horizons were described just below the horizons that showed a greater accumulation of concretions (Table 1). Yaalon (2009)

characterises vertic features as common in the lower parts of the landscape, due to the lateral and surface flow of solutions rich in Ca^{2+} and Mg^{2+} , which favours the formation of 2:1 clays. The position of vertic horizons may be related to obstruction of the vertical

flow of the soil solution resulting in the formation of such types of clay, further complemented by the alternating conditions of intense rainfall and prolonged dry periods which is characteristic of this environment, as reported by Anjos *et al.* (2007).

Table 1 - Morphological and physical characteristics of profiles P1, P2, P3, P4 and P5

Hor	Depth	Colour		Structure	Sand	Silte	Clay
	cm	Moist	Dry				
P1 - RED-YELLOW ARGISOL Abruptic eutrophic plintisol (PVAe)							
AP	0-13	5YR 3/4	5YR 4/4	mod sm bl suba mod vry sm gra	466	125	409
AB	13-23	2,5YR 3/4		mod lge vry lge bl ang	454	116	430
Btc1	23-50	2,5YR 4/6		mod sm md bl suba	220	142	638
Btcf2	50-70	5YR 5/6		mod sm md bl suba	258	107	635
B/Ccf3	70-103	5YR 4/6		mod vry sm bl suba	299	168	533
B/Cf	103-153+	5YR 4/6		mod sm bl ang suba	532	122	346
P2 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)							
Apc1	0-18	5YR 3/4	5YR 4/6	mod md bl suba	540	177	283
Ac2	18-37	5YR 4/4	5YR 5/6	mod md lge bl suba	561	115	324
Bic1	37-60	5YR 4/6	5YR 5/8	mod md bl suba	269	181	450
Bic2	60-95	5YR 4/6	5YR 6/8	mod md bl suba	356	206	438
Bi3	95-166	5YR 4,5/8	5YR 6/8	mod md bl suba	337	220	443
Bicf4	166-180	10YR 7/8	7,5YR 6/8	mod md lge bl ang suba	309	268	423
Bicf5	180-253	10YR 6/8	10YR 8/8	mod md lge bl ang suba	352	217	431
Cv1	253-285	10YR 5/6	10YR 6/6	str lge pris	190	232	578
Cr2	285-335	10YR 8/1	10YR 8/1		158	716	126
Cr3	335-430+	10YR 7/5	10YR 7/1		342	471	187
P3 - RED-YELLOW ARGISOL Abrupt eutrophic plintisol (PVAe)							
AP	0-29	7,5YR 4/3	7,5YR 4/5	str lge bl suba	460	190	350
Bt1	29-51	2,5YR 4/7	7,5YR 5/8	str md lge bl suba	310	120	570
Bt2	51-86	5YR 4/6	5YR 5/8	mod md lge bl suba	300	170	530
Bt3	86-110	5YR 5/8	5YR 5/8	mod md lge bl suba	300	140	560
Btc4	110-140	10YR 6/8	10YR 7/8		260	210	530
Cv1	140-160	10YR 5/8	10YR 6/8	str md lge pris bl ang	220	160	620
Cr2	160-190				306	457	237
P4 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)							
AP	0-31	5YR 3/4	5YR 4/6	mod md lge bl suba	450	170	380
BA	31-54	2,5YR 4/6	5YR 5/8	mod md lge bl suba	330	150	520
Bi1	54-93	2,5YR 4/6	5YR 5/7	mod md bl suba	240	160	600
Bi2	93-135	2,5YR 4,5/8	5YR 6/8	mod md bl suba	290	130	580
Bi3	135-162+	5YR 4,5/6	7,5YR 5/8	mod md bl suba	280	190	530

Table 1 continued

P5 - RED ARGISOL Eutrophic nitosol (PVe)							
A	0-12	10R 3/4	5YR 4/6	mod sm md lge bl suba	428	140	432
BA	12-44	2,5YR 4/6	2,5YR 4/8	mod md lge bl suba	228	96	676
B1	44-75	2,5YR 4/7	2,5YR 5/8	mod md bl suba	212	89	699
B2	75-124	2,5YR 4/8	2,5YR 5/7	mod md lge bl ang suba	269	130	601
B3	124-170+	2,5YR 4/8	2,5YR 5/8	mod md lge bl ang suba	363	105	532

Hor-horizon; bl-blocks; ang-angular; suba-subangular; pris-prismatic; gra-granular; vry-very; sm-small; md-medium; gde-large; mod-moderate; str-strong

Chemistry and physics

All of the profiles displayed high values for V% (Table 2), a fact also noted by Foster, Chittleborough and Barovich (2004), who found high CEC in soils identified as ‘Terra rossa’, especially for Ca^{2+} , to which is attributed the ability for particle aggregation.

The texture of the profiles is predominantly clayey and very clayey, reaching values of the order of 700 g kg^{-1} in profile P5 (Table 1). Moreover, the texture gradient present in profiles P1, P3 and P5, and the significant amount of waxiness in the last profile are evidence of clay illuviation, a feature also identified by Delgado *et al.* (2003) as one of the determining factors in the genesis of soils on karst environments in Spain.

The values for the CaCO_3 found were low, with the hypo-carbonate feature (EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA, 2013) only being identified in some of the deeper horizons of profiles P2 and P3 (Table 2). The reason for the low values may be due to better drainage in the profiles favouring solubilisation and leaching.

The Cr layers of profiles P2 and P3, materials which are still little weathered, look to be different material to that which gave rise to the overlying horizons, since they appeared strongly stratified, which may be related to wind deposition and a large selection of grains, as evidenced by the silt content (Table 1). Feng (2011), when evaluating insoluble residues of dolomitic limestones in China, found a predominance of fine materials, mainly sand and silt.

The Cr2 and Cr3 horizons of profile P2, and the Cv1 horizon of profile P3 showed an abrupt change in the values of the Fe_o/Fe_d ratio (Table 3). As they represent the basal horizons of the profiles, this increase may be related to greater humidity or suggest strata variation in the source material.

In the Cv1 horizon of profile P2, and the Cv1 horizon of profile P3, there is a strong reduction in the

amounts of iron extracted through sulphuric acid attack, and in the underlying horizons of the same profiles the reduction is even greater, together with a decrease in the values of silica, reinforcing the hypothesis of variation in the properties of the source material (Table 4).

The values for TiO_2 in horizons Cr2 and Cr3 of profile P2, and Cr2 of profile P3 show a clear contrast (Table 4), as their values are of the order of 2 g kg^{-1} in these horizons, while in the above horizons they are of the order of 6 g kg^{-1} , confirming the observations of Jiménez-Millán and Nieto (2008). Further, variation was observed in the values of TiO_2 in the above profiles, from the surface to where horizons showing a presence of concretions were identified, suggesting that the source material from the surface to the concretions differs from the source material of the last layers of these profiles.

Among the studied profiles, the Ki index (Table 4), used to relate the weathering intensity, shows that P5 is the profile displaying the highest degree of pedogenetic evolution, this can be evidenced by the lower CEC of the profile. The Kr index greater than 0.75 indicates the presence of kaolinitic soils (MOTA *et al.*, 2007).

Iron nodules

The occurrence and size of iron nodules differ for depth in the profiles under study (Figure 2). Nodules having smaller diameters, included in the sand fraction and belonging to classes C1 and C2 (Figure 2), tend to concentrate at the top of the profiles. The larger nodules, with a size equivalent to the gravel fraction (classes C3, C4 and C5), showed the highest values in horizons B and C. A similar condition was noted by Arocena, Pawluk and Dudas (1994) when evaluating nodules rich in iron in soils in Canada. Variations in such chemical conditions as oxidation / reduction, which are also influenced by such physical conditions as the wet and dry cycles, may have favoured this separation, as perhaps did the fragmentation caused by agricultural activities.

Table 2 - Chemical characteristics of profiles P1, P2, P3, P4 and P5

Hor	Depth	pH		OC	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	Al ³⁺ +H ⁺	S	T	V	CaCO ₃
	cm	H ₂ O	KCl	g kg ⁻¹	-----cmol _c kg ⁻¹ -----					%	g kg ⁻¹		
P1 - RED-YELLOW ARGISOL Abrupt eutrophic plintisol (PVAe)													
AP	0-13	8.0	7.1	10.00	6.8	3.3	1.1	0.4	0.0	11.7	11.7	100	38
AB	13-23	7.6	6.7	7.10	6.2	2.7	0.9	0.8	0.0	10.7	10.7	100	33
Btc1	23-50	7.2	6.1	2.50	6.5	3.1	0.3	0.8	0.0	10.8	10.8	100	36
Btcf2	50-70	5.7	5.0	2.40	5.2	2.9	0.1	0.7	1.8	9.0	10.8	83	38
B/Ccf3	70-103	5.4	5.0	1.60	4.3	3.3	0.1	0.6	1.5	8.4	9.9	85	37
B/Cf	103-153+	5.7	5.1	1.00	3.0	3.4	0.1	0.8	1.0	7.3	8.3	88	38
P2 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)													
Ap1	0-18	7.6	6.6	13.68	3.1	2.1	0.7	0.1	0.0	6.0	6.1	98	30
A2	18-37	7.6	6.4	7.69	2.4	2.6	0.3	0.1	0.0	5.5	5.6	98	33
Bi1	37-60	7.6	6.3	6.34	2.5	1.3	0.2	0.1	0.0	4.2	4.4	96	31
Bi2	60-95	7.0	5.9	5.65	2.6	0.9	0.1	0.2	0.0	3.8	3.9	98	36
Bi3	95-166	5.6	4.5	2.81	1.5	2.0	0.1	0.3	1.2	3.9	5.1	77	43
Bicf4	166-180	5.8	4.7	5.50	2.2	1.0	0.1	0.4	1.3	3.7	5.0	75	36
Bicf5	180-253	5.9	4.9	4.06	1.6	2.8	0.1	0.2	1.1	4.8	6.0	81	35
Cv1	253-285	7.3	5.9	3.56	7.5	10.2	0.2	0.5	0.1	18.5	18.6	99	52
Cr2	285-335	8.8	7.9	1.84	3.0	2.7	0.1	0.1	0.0	6.1	6.1	100	149
Cr3	335-430+	8.7	7.6	2.69	3.8	4.4	0.2	0.3	0.0	8.7	8.7	100	149
P3 - RED-YELLOW ARGISOL Abrupt eutrophic plintisol (PVAe)													
AP	0-29	8.0	7.2	18.78	3.1	1.6	0.3	0.1	1.0	5.2	6.2	87	29
Bt1	29-51	7.9	6.7	7.44	4.7	1.6	0.2	0.3	1.0	6.9	7.9	87	31
Bt2	51-86	7.6	6.3	4.28	5.0	2.3	0.1	0.4	2.0	8.0	10.0	77	29
Bt3	86-110	6.4	5.4	4.31	4.0	2.4	0.1	0.4	2.0	6.9	8.9	74	29
Btc4	110-140	6.4	5.3	2.75	3.2	3.2	0.1	0.3	2.0	6.8	8.8	78	26
Cv1	140-160	6.1	4.9	2.91	3.7	5.6	0.1	0.6	2.0	11	13.1	86	38
Cr2	160-190	8.6	7.6	2.22	6.6	3.8	0.1	0.3	0.0	11	11	100	149
P4 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)													
AP	0-31	7.7	6.8	14.31	5.4	2.4	0.3	0.1	0.0	8.2	8.2	100	10
BA	31-54	7.7	6.5	6.65	4.4	0.9	0.2	0.1	0.0	5.7	5.7	98	32
Bi1	54-93	6.6	5.6	3.94	3.4	1.4	0.2	0.1	0.0	5.1	5.1	98	30
Bi2	93-135	5.9	4.8	4.31	2.9	1.5	0.2	0.1	2.1	4.8	6.8	69	29
Bi3	135-162+	6.2	5.1	4.37	3.1	1.8	0.1	0.1	2.1	5.3	7.4	71	32
P5 - RED ARGISOL Eutrophic nitosol (PVe)													
A	0-12	7.0	5.8	18.81	3.2	2.5	0.1	0.1	0.0	5.8	5.8	99	33
BA	12-44	7.7	6.1	8.25	3.1	1.7	0.1	0.1	0.0	5.0	5.1	98	33
B1	44-75	6.1	4.9	5.56	2.1	1.8	0.1	0.1	1.0	4.0	5.1	79	30
B2	75-124	5.2	4.6	4.19	1.8	2.2	0.1	0.1	2.1	4.1	6.2	66	28
B3	124-170+	5.2	4.7	4.12	1.7	2.0	0.1	0.1	2.2	3.9	6.1	64	29

Hor-Horizon; OC-organic carbon; S-sum of bases; T-cation exchange capacity; V-Base saturation


Table 3 - Values for iron in sodium dithionite-citrate-bicarbonate (Fe_d), iron in ammonium oxalate (Fe_o), ratio of iron in the oxalate to iron in the dithionite (Fe_o/Fe_d) and ratio of iron in the dithionite to clay ($Fe_d/Clay$) in profiles P1, P2, P3, P4 and P5

Horizon	Depth	Fed	Feo	Feo/Fed	Fed/Clay
	cm	-----g kg ⁻¹ -----			
P1 - RED-YELLOW ARGISOL Abrupt eutrophic plintisol (PVAe)					
Ap	0-13	20.41	2.58	0.13	0.05
AB	13-23	43.91	2.91	0.07	0.10
Btc1	23-50	27.84	2.54	0.09	0.04
Btcf2	50-70	26.34	1.92	0.07	0.04
B/Ccf3	70-103	22.79	1.59	0.07	0.04
B/Cf	103-153+	22.34	1.63	0.07	0.06
P2 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)					
Ap1	0-18	28.24	2.23	0.08	0.10
A2	18-37	29.57	2.28	0.08	0.09
Bi1	37-60	29.28	2.72	0.09	0.07
Bi2	60-95	23.48	1.71	0.07	0.05
Bi3	95-166	26.62	1.08	0.04	0.06
Bicf4	166-180	30.80	1.89	0.06	0.07
Bicf5	180-253	31.95	1.80	0.06	0.07
Cv1	253-285	17.16	1.73	0.10	0.03
Cr2	285-335	2.59	0.39	0.15	0.02
Cr3	335-430+	3.34	3.79	1.14	0.02
P3 - RED-YELLOW ARGISOL Abrupt eutrophic plintisol (PVAe)					
Ap	0-29	20.27	2.39	0.12	0.06
Bt1	29-51	18.87	2.72	0.14	0.03
Bt2	51-86	18.00	2.25	0.13	0.03
Bt3	86-110	27.40	1.82	0.07	0.05
Btc4	110-140	35.29	2.88	0.08	0.07
Cv1	140-160	20.66	4.35	0.21	0.03
Cr2	160-190	10.36	0.79	0.08	0.04
P4 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)					
Ap	0-31	38.88	2.67	0.07	0.10
BA	31-54	23.97	2.85	0.12	0.05
Bi1	54-93	24.82	2.63	0.11	0.04
Bi2	93-135	31.26	2.44	0.08	0.05
Bi3	135-162+	20.54	2.72	0.13	0.04
P5 - RED ARGISOL Eutrophic nitosol (PVe)					
A	0-12	22.24	3.31	0.15	0.05
BA	12-44	17.84	2.80	0.16	0.03
B1	44-75	26.12	1.85	0.07	0.04
B2	75-124	20.84	2.32	0.11	0.03
B3	124-170+	24.82	1.51	0.06	0.05

Table 4 - Percentage of SiO₂, Fe₂O₃, Al₂O₃, MnO₂ and TiO₂ and the molecular indices Kr and Ki determined in the extract from the sulphuric acid attack for profiles P1, P2, P3, P4 and P5

Horizon	Depth	SiO ₂	Fe ₂ O ₃	Al ₂ O ₃	MnO ₂	TiO ₂	Ki	Kr
	cm	-----g kg ⁻¹ -----						
P1 - RED-YELLOW ARGISOL Abrupt eutrophic plintisol (PVAe)								
Ap	0-13	164.0	73.8	151.3	0.57	5.20	1.84	1.41
AB	13-23	267.0	79.8	179.8	0.52	5.43	2.52	1.97
Btc1	23-50	275.8	81.6	217.5	0.45	6.13	2.16	1.74
Btcf2	50-70	310.7	83.5	224.4	0.71	6.37	2.35	1.90
B/Ccf3	70-103	242.0	80.5	202.1	0.35	6.45	2.04	1.62
B/Cf	103-153+	268.5	75.6	191.4	0.41	5.65	2.38	1.90
P2 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)								
Ap1	0-18	155.4	79.7	146.7	0.94	5.18	1.80	1.34
A2	18-37	161.3	86.6	153.1	0.93	5.28	1.79	1.32
Bi1	37-60	223.5	79.4	186.3	0.64	6.54	2.04	1.60
Bi2	60-95	207.2	88.2	196.4	0.64	7.05	1.79	1.39
Bi3	95-166	260.5	68.9	171.7	0.39	6.03	2.58	2.05
Bicf4	166-180	257.1	61.9	188.9	0.29	5.02	2.31	1.91
Bicf5	180-253	271.6	99.6	220.4	0.44	6.42	2.10	1.63
Cv1	253-285	323.3	49.7	122.7	1.95	4.48	4.48	3.56
Cr2	285-335	-	10.1	28.2	0.27	1.33	0.00	-
Cr3	335-430+	10.8	17.9	50.2	0.31	2.18	0.37	0.30
P3 - RED-YELLOW ARGISOL Abrupt eutrophic plintisol (PVAe)								
Ap	0-29	184.5	67.9	137.6	0.81	5.28	2.28	1.73
Bt1	29-51	214.1	80.9	173.5	0.69	6.17	2.10	1.62
Bt2	51-86	261.4	86.7	187.8	0.46	6.22	2.37	1.83
Bt3	86-110	278.1	86.0	195.9	0.37	6.01	2.41	1.89
Btc4	110-140	268.5	97.8	207.5	0.91	6.25	2.20	1.69
Cv1	140-160	290.4	88.0	203.9	0.89	7.26	2.42	1.90
Cr2	160-190	7.0	15.3	43.0	0.15	1.81	0.28	0.22
P4 - HAPLIC CAMBISOL Tb Eutrophic latosol (CXbe)								
Ap	0-31	182.5	72.1	145.4	0.92	5.02	2.13	1.62
BA	31-54	271.0	77.5	210.3	0.52	6.13	2.19	1.77
Bi1	54-93	285.6	79.4	219.8	0.49	6.76	2.21	1.80
Bi2	93-135	249.7	64.1	175.9	0.50	5.07	2.41	1.96
Bi3	135-162+	266.4	71.8	195.2	0.44	5.39	2.32	1.88
P5 - RED ARGISOL Eutrophic nitosol (PVe)								
A	0-12	187.4	87.8	179.9	1.04	6.37	1.77	1.35
BA	12-44	295.0	87.1	245.0	0.67	6.54	2.05	1.67
B1	44-75	276.7	95.2	259.5	0.53	7.02	1.81	1.47
B2	75-124	301.4	95.1	260.7	0.47	6.52	1.97	1.59
B3	124-170+	305.4	94.9	254.6	0.37	5.89	2.04	1.65


Figure 2 - Representation of the distribution of nodules divided by size class (C1, C2, C3, C4, and C5) in profiles P1, P2, P3, P4 and P5



Yaalon (2009) reports that nodules of different sizes, quantities, hardness and distribution are common in soils developed on carbonate rocks and in an arid climate. Jiménez-Millán and Nieto (2008), studying concretions of iron-manganese in carbonate formations in southeastern Spain, report that these concretions are not only common, but strongly related to differences in facies in the source material, which even in the same rock presents small chemical and structural differences.

Average values for density were around 2.6 Mg m^{-3} and similar to those found by Singh and Gilkes (1996). It can also be seen that nodule density is greater on the surface, becoming smaller with depth (Figure 3). The increase in density takes place with a decrease in diameter. This inverse relationship between the size and density of the concretions is related to incorporation of the soil matrix, mainly quartz and kaolinite, during the formation and development

Figure 3 - Representation of the average density of nodules divided by class (C1, C2, C3, C4, and C5) in profiles P1, P2, P3, P4 and P5



of the nodules increasing the pore space, thereby resulting in a lower density.

In profiles P1, P2 and P3, high proportions of rounded-habit nodules were seen in the gravel fraction (classes C3, C4 and C5), which displayed a low mechanical resistance. However, nodules with smaller diameters, belonging to the sand fraction, were mechanically resistant, resembling lead shot.

Given the conditions for development of the limestone that makes up the Jandaíra Formation, such as shallow-water marine environments, the formation of rounded limestone structures is common and often associated with ferrous compounds (CASSAB, 2003). Also characteristic of this environment is a rise in water temperature, favouring the precipitation of carbonates, mainly Ca and Fe, while the movement of the waves may have rounded the bodies that were formed.


Analysing the levels of Fe_d in the nodules of profiles P2 and P4, classified as Cambisols (Figure 4), values increase with diameter for classes C1, C2 and C3, suggesting greater variation in the degree of crystallinity. Whereas for profiles P3 and P5, classified as Argisols, there is greater homogeneity, especially in the last profile which has characteristics related to more evolved soil.

In profile P5 the horizons displayed a homogeneous distribution for nodule mass and density, a fact which demonstrates progress of the weathering processes, a slight increase can also be seen in the quantity of nodules of classes C1, C2 and C3 on the surface, which may be related to an environmental condition which favours their preservation, since in this horizon conditions of hydromorphism are virtually absent when compared to the subsurface layers.

Also in profile P5 no nodules of classes C4 and C5 were recorded (Figure 2), which confirms that, both in the horizons and in the more advanced profiles, there is a reduction in larger-diameter iron nodules, suggesting their degradation with the advancement of the weathering process.

From the observations made here, a significant relationship can be seen between the processes of oxidation/reduction in the dissolution of limestone material, and consequently a change in the dynamics of the iron accumulating in the form of nodules in horizons of currently oxidic conditions. In this study, the number of nodules identified and their characteristics are not seen as limiting the development of agricultural activities; a fact which is relevant faced with the economic importance of this region, however genetic peculiarities in the nodules should be pointed out which are giving rise to more extensive approaches.

Figure 4 - Average values for iron in the sodium dithionite-citrate-bicarbonate (Fe_d) and iron in the ammonium oxalate (Fe_o) for classes C1, C2, C3, C4 and C5 of profiles P1, P2, P3, P4 and P5



CONCLUSIONS

1. The influence of the factors of formation on the process of genesis has the source material as a factor of determining importance; with the soils presenting variations in morphological, chemical and physical data, indicating a change in the characteristics of the source material, with emphasis on variations in the amounts of iron and titanium;
2. The reduction in the amount and diameter of the iron nodules in the more evolved horizons and profiles is evidence that such nodules are in the process of degradation, influenced by the advanced weathering process.

REFERENCES

- ACHURRA, L. E. *et al.* Manganese nodules in the Miocene Bahía Inglesa Formation, north-central Chile: Petrography, geochemistry, genesis and palaeoceanographic significance. **Sedimentary geology**, v. 217, n. 1, p. 128-139, 2009.
- ALENCAR, E. L. L. **Química e mineralogia de três pedons originários de calcário da Chapada do Apodi-CE**. 2002. 83 f. Dissertação (Mestrado em Solos e Nutrição de Plantas) - Universidade Federal do Ceará, Fortaleza, 2002.
- ANJOS, L. H. C. *et al.* Caracterização e classificação de plintossolos no município de Pinheiro-MA. **Revista Brasileira de Ciências do Solo**, v. 31, n. 5, p. 1035-1044, 2007.
- AROCENA, J. M.; PAWLUK, S.; DUDAS, M. J. Iron oxides in iron rich nodules of sandy soils from Alberta (Canada). *In*: RINGROSE-VOASE, A. J.; HUMPHREYS, G. S. (Ed.), **Soil Micromorphology: Studies in Management and Genesis**. Elsevier, 1994. p. 83-97.
- AYDINALP, C.; FITZPATRICK, E. A. Pedogenesis and characteristics of the Terra Rossas developed on different physiographic position and their classification. **Agrociencia**, v. 43, n. 2, p. 97 -105, 2009.
- BAUTISTA, F. *et al.* Spatial distribution and development of soils in tropical karst areas from the Peninsula of Yucatan, Mexico. **Geomorphology**, v. 135, n. 3/4, p. 308-321, 2011.
- BRASIL. MINISTÉRIO DA AGRICULTURA. **Levantamento Exploratório-Reconhecimento de solos do estado do Ceará**. Recife, 1973. 297 p. v. 1.
- BRASIL. MINISTÉRIO DAS MINAS E ENERGIA. SECRETARIA GERAL. **Projeto RADAMBRASIL**. Folhas SB. 24/25 Jaguaribe/Natal; geologia, geomorfologia, pedologia, vegetação e uso potencial da terra. Rio de Janeiro, 1981. 774 p. v. 23.
- CASSAB, R. C. T. **Paleontologia da Formação Jandaíra, cretáceo superior da bacia potiguar, com ênfase na paleobiologia dos gastrópodos**. 2003. 184 f. Tese (Doutorado em Geologia) Universidade Federal do Rio de Janeiro, Rio de Janeiro.
- COELHO, M. R.; VIDAL-TORRADO, P. Caracterização e gênese de perfis plínticos desenvolvidos de arenito do grupo Bauru. II - Mineralogia. **Revista Brasileira de Ciência do Solo**, v. 27, n. 3, p. 495-507, 2003.
- CONSTANTINI, E. A. C. *et al.* Using the analysis of iron and iron oxides in paleosols (TEM, geochemistry and iron forms) for the assessment of present and past pedogenesis. **Quaternary International**, v. 156/157, p. 200-211, 2006.
- CONSTANTINI, E. A. C.; PRIORI, S. Pedogenesis of plinthite during early Pliocene in the Mediterranean environment: Case study of a buried paleosol at Podere Reniere, central Italy. **Catena**, v. 71, n. 3, p. 425-443, 2007.
- DELGADO, R. *et al.* Genesis of the terra rossae of the Sierra Gádor (Andalusia, Spain). **European Journal of Soil Science**, v. 54, n. 1, p. 1-16, 2003.
- EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA. Centro Nacional de Pesquisa de solos **Manual de métodos de análise de solo**. 2. Ed. Rio de Janeiro: Embrapa Solos, 1997. 212 p.
- EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA. Centro Nacional de Pesquisa de solos 2013. **Sistema Brasileiro de Classificação de Solos**. 3. ed. Brasília, 2013. 353 p.
- FENG, J. L. Trace elements in ferromanganese concretions, gibbsite spots, and the surrounding terra rossa overlying dolomite: Their mobilization, redistribution and fractionation. **Journal of Geochemical Exploration**, v. 108, n. 1, p. 99-111, 2011.
- FOSTER, J.; CHITTLEBOROUGH, D. J.; BAROVICH, K. **Genesis of a Terra Rossa soil over marble and the influence of a neighbouring texture contrast soil at Delamere, South Australia**. 2004. Australia. *In*: AUSTRALIAN NEW ZEALAND SOILS CONFERENCE, Australia: University of Sydney, Australia, 2004. p. 5-9. 1 CD-ROM.
- HUANG, L. *et al.* Characteristics of micromorphology and element distribution of iron-manganese cutans in typical soils of subtropical China. **Geoderma**, v. 146, n. 1/2, p. 40-47, 2008.
- INDA JUNIOR, A. V.; KÄMPF, N. Avaliação de procedimentos de extração dos óxidos de ferro pedogênicos com ditionito-citrato-bicarbonato de sódio. **Revista Brasileira de Ciências do Solo**, v. 27, n. 6, p. 1139-1147, 2003.
- INDA JUNIOR, A. V.; KÄMPF, N. Variabilidade de goethita e hematita via dissolução redutiva em solos de região tropical e subtropical. **Revista Brasileira de Ciências do Solo**, v. 29, n. 6, p. 851-866, 2005.
- JIMÉNEZ-MILLÁN, J.; NIETO, L. M. Geochemical and mineralogical evidence of tectonic and sedimentary factors controlling the origin of ferromanganese crusts associated to stratigraphic discontinuities (Betic Cordilleras, SE of Spain). **Chemie der Erde - Geochemical**, v. 68, n. 3, p. 323-336, 2008.

- KHORMALI, F. *et al.* Argillic horizon development in calcareous soils of arid and semiarid regions of southern Iran. **Catena**, v. 53, n. 3, p. 273-301, 2003.
- LYNCH, L. S. **Gênese e geoquímica de solos em ambiente cárstico no cerrado na região de Planaltina de Goiás**. 2009. 155 f. Tese (Doutorado em Solos e Nutrição de Plantas) - Universidade Federal de Viçosa, Viçosa, 2009.
- MEHRA, O. P.; JACKSON, M. L. Iron oxide removal from soils and clays by a dithionite-citrate system buffered with sodium bicarbonate. **Clay Clay Mineralogy**, v. 72, p. 317-327, 1960.
- MOTA, J. C. A. *et al.* Atributos mineralógicos de três solos explorados com a cultura do melão na chapada do Apodi-RN. **Revista Brasileira de Ciências do Solo**, v. 31, n. 3, p. 445-454, 2007.
- MOTA, J. C. A. *et al.* Algumas propriedades físicas e hídricas de três solos na chapada do Apodi, RN, cultivados com melão. **Revista Brasileira de Ciências do Solo**, v. 32, n. 1, p. 49-58, 2008.
- OLIVEIRA, C. V. *et al.* Atributos micromorfológicos de solos do projeto Jaíba, norte de Minas gerais. **Revista Brasileira de Ciências do Solo**, v. 24, n. 1, p. 117-128, 2000.
- PEREIRA, M. G. *et al.* Caracterização e classificação de solos em uma topossequência sobre calcário na serra da bodoquena, MS. **Revista Brasileira de Ciências do Solo**, v. 37, n. 1, p. 25-36, 2013.
- SANTOS, R. D. *et al.* **Manual de descrição e coleta de solo no campo**. 5. ed. Viçosa: Sociedade Brasileira de Ciência do Solo, 2005. 100 p.
- SILVA, M. B. *et al.* Gênese e classificação dos solos de uma topossequência em área de carste na Serra da Bodoquena, MS. **Revista Brasileira de Ciências do Solo**, v. 37, n. 1, p. 1464-1480, 2013.
- SILVA NETO, L. F. Óxidos de ferro em latossolos tropicais e subtropicais brasileiros em plantio direto. **Revista Brasileira de Ciências do Solo**, v. 32, p. n. 5, 1873-1881, 2008.
- SINGH, B.; GILKES, R. J. Nature and properties of iron rich glaebules and mottles from some south-west Australian soils. **Geoderma**, v. 71, n. 1/2, p. 95-120, 1996.
- TAN, W. *et al.* Elemental composition and geochemical characteristics of Iron-Manganese nodules in main soils of China. **Pedosphere**, v. 16, n. 1, p. 72-81, 2006.
- TARDY, Y.; NAHON, D. B. Geochemistry of laterites. Stability of Al-goethite, Al-hemathite and Fe³⁺-kaolinite in bauxites and ferricretes. An approach to the mechanism of concretion formation. **American Journal of Science**, v. 285, p. 865-903, 1985.
- URLEY, A. L.; DREES, L. R. **Methods of Soil Analysis**. Soil Science Society of America. Madison, 2008. 521 p.
- YAALON, D. H. Brief comments on red Mediterranean soils. **Catena**, v. 76, n. 3, p. 224, 2009.
- YEOMANS, J. C.; BREMNER, J. M. A rapid and precise method for routine determination of organic carbon in soil. **Communication in Soil Science Plant Analysis**, v. 19, n. 13, p.1467-1476, 1988.