

## ORIGINAL

## Clinical Nutrition

## Editor

Alex Harley Crisp

## Conflict of interest

The authors declare that there is no conflict of interests.

## Received

February 9, 2024

## Final version






June 17, 2024

## Approved

July 16, 2024

# Relationship between ultrasound measurement of quadriceps muscle and nutritional status in ICU patients in a high-complexity trauma care hospital

*Relação entre medida do músculo quadríceps pelo ultrassom e estado nutricional em pacientes internados em UTI de um hospital de alta complexidade de atendimento ao trauma*

Milton Alves Danziato-Neto<sup>1</sup> , Priscilla Sousa Santos Caldas<sup>1,2</sup> , Juliana Magalhães da Cunha Rêgo<sup>1</sup> , Antonio Augusto Ferreira Carioca<sup>3</sup> , Cristiane Rodrigues Silva Câmara<sup>1,3</sup> 

<sup>1</sup> Instituto Doutor José Frota. Fortaleza, CE, Brasil. Correspondence to: MA DANZIATO-NETO. E-mail: <danziatomilton@gmail.com>.

<sup>2</sup> Escola de Saúde Pública do Ceará. Fortaleza, CE, Brasil.

<sup>3</sup> Universidade de Fortaleza, Centro de Ciências da Saúde. Fortaleza, CE, Brasil.

**How to cite this article:** Danziato-Neto MA, Caldas PSS, Rêgo JMC, Carioca AAF, Câmara CRS. Relationship between ultrasound measurement of quadriceps muscle and nutritional status in ICU patients in a high-complexity trauma care hospital. Rev Nutr. 2024;37:e240005. <https://doi.org/10.1590/1678-9865202437e240005>

## ABSTRACT

### Objective

The present study aimed to investigate the association of quadriceps muscle thickness, measured by ultrasound, with the nutritional status of critically ill patients in a referral high-complexity trauma care hospital.

### Methods

A cross-sectional observational study was conducted in the intensive care units in a tertiary hospital in Brazil. The sample comprised 30 critically ill trauma patients admitted between February and March 2022. The methodology involved evaluating muscle mass and comparing nutritional status through mid-upper arm circumference measurements and ultrasound assessments. Specifically, the quadriceps muscle thickness was quantified using ultrasound at a predefined site between the iliac crest and the proximal border of the patella.

### Results

The Kruskal-Wallis test indicated variability in quadriceps muscle thickness between the nutritional status groups, with statistical significance reached after excluding the overweight group ( $H(2) = 7.532, p=0.023$ ). The moderate malnutrition group exhibited notably lower quadriceps muscle thickness. A positive correlation was found between quadriceps muscle

thickness and mid-upper arm circumference adequacy ( $p < 0.05$ ), demonstrating fair to moderately strong correlation ( $r_s = 0.531$ ).

### Conclusion

Significant changes in quadriceps muscle thickness were detected by ultrasound assessment in moderate malnutrition patients compared to patients of other nutritional statuses. Ultrasound may be a valuable technique for monitoring muscle integrity in critically ill patients.

**Keywords:** Imaging. Intensive Care Units. Malnutrition. Muscle, skeletal. Nutritional status. Quadriceps muscle. Ultrasonics.

## RESUMO

### Objetivo

O presente estudo teve como objetivo investigar a associação da espessura do músculo quadríceps, medida por ultrassom, com o estado nutricional de pacientes críticos em um hospital de referência de alta complexidade para o cuidado de traumas.

### Métodos

Um estudo observacional transversal foi conduzido nas unidades de terapia intensiva de um hospital terciário no Brasil. A amostra compreendeu 30 pacientes críticos de trauma admitidos entre fevereiro e março de 2022. A metodologia envolveu a avaliação da massa muscular e a comparação do estado nutricional por meio de medições da circunferência do braço e avaliações por ultrassom. Especificamente, a espessura do músculo quadríceps foi quantificada usando ultrassom em um local pré-definido entre a crista ilíaca e a borda proximal da patela.

### Resultados

O teste de Kruskal-Wallis indicou variabilidade na espessura do músculo quadríceps entre os grupos de estado nutricional, com significância estatística alcançada após a exclusão do grupo com sobrepeso ( $H(2) = 7.532$ ,  $p = 0,023$ ). O grupo com desnutrição moderada exibiu um resultado menor em espessura do músculo quadríceps. Foi encontrada uma correlação positiva entre a espessura do músculo quadríceps e a adequação da circunferência do braço médio-superior ( $p < 0,05$ ), demonstrando uma correlação de razoável a moderadamente forte ( $r_s = 0,531$ ).

### Conclusão

Mudanças significativas na espessura do músculo quadríceps foram detectadas pela avaliação de ultrassom em pacientes com desnutrição moderada em comparação com pacientes de outros estados nutricionais. O ultrassom pode ser uma técnica valiosa para monitorar a integridade muscular em pacientes críticos.

**Palavras-chave:** Imagem. Unidades de Terapia Intensiva. Desnutrição. Músculo esquelético. Estado nutricional. Músculo quadríceps. Ultrassom.

## INTRODUCTION

Critically ill patients admitted to an Intensive Care Unit (ICU) present both physiological instability and a high risk of mortality [1]. These patients typically experience changes in body composition, mainly caused by hypermetabolism resulting from disease-associated inflammation, reduced food intake and mobility, and drug effects. These changes can lead to an intense loss of muscle mass and function [2,3].

A broad scientific literature suggests that malnutrition and reduced muscle mass in hospitalized patients serve as predictors of worsening clinical outcomes, particularly among those receiving intensive care, leading to an increased risk of morbidity, mortality, and impaired long-term functionality [4]. Therefore, it is essential that muscle mass evaluation be part of the nutritional diagnosis [5] and the monitoring muscle losses be used to guide interventions for muscle recovery [6].

Nutritional assessment comprises evaluation of body composition, which can be challenging in critically ill patients since anthropometric measurements can be altered by hemodynamic changes

commonly observed during their ICU stay [7]. In addition, trauma injuries can limit or make it impossible to use many instruments and techniques involved in nutritional assessment [8].

Mid-Upper Arm Circumference (MUAC) is a standard protocol in nutritional assessment as it can be used to determine the patient's nutritional status and provides a necessary variable for estimating weight using predictive equations [9,10]. However, MUAC measurements can be affected by edema, a clinical condition commonly present in ICU patients [8], and it does not provide specific information on muscle composition.

Ultrasound (US) is progressively becoming a promising tool for assessing muscle mass in various clinical populations, including critically ill patients. This device confers good reproducibility, reliability, and diagnostic accuracy for both intra and inter-examiners [11,12]. US is a modality that allows measurements in various sections of muscles, organs, and adipose tissue. The resulting images offer a direct visualization of the region under study, enabling the observation of compositional differences in muscular mass or adiposity, among other parameters [13].


The evaluation of Quadriceps Muscle Thickness (QMT) has been reported to have good reliability and validity for the diagnosis and monitoring of acute muscle wasting in critically ill patients, particularly when assessed at the bedside using US [14]. Several authors have highlighted it as an excellent region that is both practical and accessible for the application of US in studying patients admitted to the ICU [3,15,16].

Vallet et al. [17] emphasize the possibility of using US to assess muscle mass in patients with sarcopenia, a progressive condition, often age-induced, that compromises both muscle mass and function and may have a significant adverse effect on outcomes in ICU, especially in older patients. In this context, evaluating QMT may be a key parameter of nutritional assessment in ICU in many cases since other common anthropometric measurements could not be useful considering the high incidence of subcutaneous edema.

In the care of critically ill patients who are at high nutritional risk, the need for procedures that specifically monitors muscle mass integrity is paramount to establish nutritional diagnosis and to guide dietary treatment. Thus, the purpose of this study was to investigate the association of QMT, measured by US, with nutritional status of critically ill patients in a referral high-complexity trauma care hospital. This study method can allow us to evaluate the sensitivity of this muscle region to changes in nutritional status, which can improve the nutritional diagnosis method. Besides, the follow-up of changes in the US can help assess the effect of nutritional therapy on the muscles.

## METHODS

This observational quantitative cross-sectional study was conducted on patients admitted to the ICUs of a highly complex public hospital, which serves as a reference in trauma for the North and Northeastern regions of Brazil, between February and March 2022. The study was carried out with the approval of the hospital's ethics committee (CAAE 41909321.8.0000.5047) and informed consent forms were signed by the legal guardians of the participants. A non-probabilistic convenience sampling method was applied, with inclusion criteria consisting of adult and elderly patients ( $\geq 18$  years old) of both genders. Exclusion criteria comprised individuals with amputated upper or lower limbs, edema in both arms, burned patients, pregnant women, and those whose condition did not permit access to the quadriceps for US evaluation (Figure 1).

**Figure 1** – Flowchart of sample inclusion and exclusion criteria.

Note: ICU: Intensive Care Unit; QMT: Quadriceps Muscle Thickness; US: Ultrasound.

### *Data collection*

The collection of clinical data and identification from the medical records was performed through the utilization of a data collection form. The data collected were gender, age, hospitalization date, medical diagnosis, comorbidities, and physical exam (e.g., presence of edema, muscle mass atrophy, and fat tissue excess).

### *Mid-upper arm circumference measurement*


Mid-upper arm circumference was measured to classify patient's nutritional status based on its adequacy, according to Blackburn and Thornton [18]. The measurement of MUAC was conducted following the method described by Lohman [9]. For accurate measurements, an inelastic measuring tape marked in centimeters (cm) and yielding a precision of 0.1cm was utilized.

### *Ultrasound measurements*

The quadriceps muscle thickness (QMT) quantification was obtained by using a portable Mindray® ultrasound device, Model M6, equipped with a 19" screen, which allowed for obtaining an image in mode B, with a (7L4s) transducer and a sampling frequency of 5-10 MHz. Precise measurements, in centimeters, were taken perpendicularly and transversely on the evaluated

muscles. The assessment was performed with the patient in dorsal decubitus, with the elbows and knees in passive extension, to allow examination of the muscle when it is extended and relaxed. Sterile gel was applied to the US transducer to facilitate contact with the skin surface. The transducer was placed on the anterior surface of the thigh, transversely in relation to the muscle length and perpendicularly to the longitudinal axis, at the point corresponding to two-thirds of the distance between the iliac crest and the upper border of the patella [19-21] (Figure 2A).

**Figure 2** – Illustration of the anatomic site used for QMT quantification and a QMT image obtained in the present study.



Note: Figure A: Point corresponding to two-thirds of the distance between the iliac crest and the patella. Figure B: Quadriceps muscle thickness as detected by a participant's ultrasound image. The dotted line represents the full thickness of the quadriceps (rectus femoris and vastus intermedius complex). RF: rectus femoris; VI: vastus intermedius; VL: Vastus Lateralis; VM: Vastus Medialis. In the image, only small portions of the VM and VL are visible. These images are licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

With the aim of standardizing measurements for all participants, the position of the US transducer was determined after demarcating the midpoint with the aid of an inelastic measuring tape graduated in centimeters (cm) and with a precision of 0.1cm. To calculate the median, US measurements were performed in duplicate. US images of the region where the measurements were

performed were also collected. Figure 2B illustrates one of the ultrasound images obtained in the present study, demarcating the QMT. The use of only vastus intermedius and rectus femoris among all quadriceps muscles in QMT assessment is justified for its clinical practicality and relevance, as supported by studies such as Toledo et al. [3] and Paris et al. [22].

### *Statistical analysis*

Data were analyzed using IBM SPSS Statistics 26.0 and sensitivity analyses by R software version 4.3.1. Categorical variables were presented as relative frequencies. The Shapiro-Wilk test assessed the normality of the distribution of continuous variables. Given the non-normal distribution of some variables, measures of central tendency and dispersion were presented as mean  $\pm$  standard deviation (SD) for normal data and median with interquartile range (IQR) for skewed data.

The Kruskal-Wallis test was utilized to identify differences between more than two independent groups, accounting for the non-normal distribution. Additionally, Spearman's rank correlation coefficient explored associations between quantitative muscle thickness (QMT) with both mid-upper arm circumference (MUAC) and its adequacy (MUACA). The interpretation of the Spearman's correlation coefficients was based on the guidelines provided by Chan [23]. The significance level was set at  $p < 0.05$ .

The interpretation of Spearman's correlation coefficients according to Chan [23], is as follows. A Spearman's correlation coefficient ( $\rho$ ) of  $\pm 1$  indicates a perfect correlation, meaning there is a perfect monotonic relationship between two variables. Coefficients between  $\pm 0.8$  and  $\pm 0.9$  signify a very strong correlation. Coefficients between  $\pm 0.6$  and  $\pm 0.7$  denote a moderately strong correlation. Coefficients between  $\pm 0.3$  and  $\pm 0.5$  reflect a fair correlation. Coefficients between  $\pm 0.1$  and  $\pm 0.2$  represent a poor correlation. A coefficient of 0 suggests no correlation, implying that the rank of one variable does not predict the rank of the other [23].

A critical element of the statistical approach was conducting sensitivity analyses using resampling methods, specifically bootstrap and Monte Carlo simulation, to validate the robustness of the findings. These methods were paramount given the small sample size in the Overweight group, raising concerns about the potential bias and the stability of the results. With 2,000 iterations, the bootstrap method was employed to estimate the distribution of the Kruskal-Wallis test statistic under the null hypothesis, providing a confidence interval for this statistic. The models were refitted to each bootstrap sample, and we calculated the mean statistics and their bias-corrected 95% confidence intervals (CI). Monte Carlo simulation, also set at 2,000 iterations, assessed the p-value distribution. Through 2,000 iterations, it assessed the distribution of p-values and test statistics beyond conventional analytical constraints, providing a more nuanced understanding of the data's statistical properties.

## **RESULTS**

Table 1 presents the general characteristics of all study participants. Of the 30 patients evaluated, 28 (93.3%) were male and 2 (6.7%) were female, with a median age of 40.4 years (interquartile range [IQR], 23.75-53 years). The median length of hospital stay was 24 days (IQR, 13.25-68.5 days). Traumatic brain injury and polytrauma were the primary reasons for hospitalization, accounting for 76.7% of the sample. None of the participants were obese or with severe malnutrition, and only two individuals (6.7%) were overweight. For the moderate malnutrition group, the QMT mean  $\pm$  SD were  $1.46 \pm 0.46$ cm ( $p=0.547$ ). The eutrophy group showed a QMT mean  $\pm$  SD of  $2.12 \pm 0.43$ cm

( $p=0.785$ ). The mild malnutrition group had a QMT median of 2.09cm (IQR, 2.03-2.23cm) ( $p=0.032$ ). The two observed QMT values for the Overweight group were 1.92cm and 2.11cm.

**Table 1** – Demographic, anthropometric, and clinical characteristics of the participants.

| Characteristics   | Total (n=30)      | $p^*$  |
|---|-------------------|--------|
| Sex, %  |                   |        |
| Men   | 93.3              |        |
| Women   | 6.7               |        |
| Age in years, median (IQR)                                    | 40.5 [23.75-53.0] | 0.042  |
| Length of stay in days, median (IQR)                          | 24.0 [13.25-68.5] | <0.001 |
| Reason for hospitalization, %                                 |                   |        |
| Brain cancer  | 3.3               |        |
| Exogenous intoxication  | 13.3              |        |
| Polytrauma  | 36.7              |        |
| Traumatic brain injury  | 40.0              |        |
| Spinal cord injury  | 6.7               |        |
| Patients on MV > 48h, %                                       | 90.0              |        |
| MUAC (cm), mean $\pm$ SD                                      | 28.8 $\pm$ 4.19   | 0.335  |
| MUACA (% $\pm$ SD)  | 91.0 $\pm$ 13.5   | 0.648  |
| Nutritional status (%)  |                   |        |
| Moderate malnutrition   | 23.3              |        |
| Mild malnutrition   | 20.0              |        |
| Eutrophy  | 50.0              |        |
| Overweight  | 6.7               |        |
| QMT (cm), mean $\pm$ SD                                       | 1.97 $\pm$ 0.46   | 0.798  |
| QMT (cm) by nutritional status, mean $\pm$ SD or median (IQR) |                   |        |
| Moderate malnutrition   | 1.46 $\pm$ 0.46   | 0.547  |
| Mild malnutrition   | 2.09 [2.03-2.23]  | 0.032  |
| Eutrophy  | 2.12 $\pm$ 0.43   | 0.785  |
| Overweight  | 1.92 and 2.11     | -      |

Note: \*All continuous variables were tested for normality using the Shapiro-Wilk test,  $p$ -value <0.05. MUAC: Mid-Upper Arm Circumference; MV: Mechanical Ventilation; QMT: Quadriceps Muscle Thickness; SD: Standard Deviation.


Upon applying the Kruskal-Wallis test to all groups (N=30), the statistics were as follows:  $H(3) = 7.635$ ,  $p=0.054$ . The moderate malnutrition group had a QMT median of 1.44 cm (IQR, 1.19-1.86cm). The eutrophy group showed a QMT median of 2.08cm (IQR, 1.82-2.47cm). The overweight group, limited in sample size, had a median of 2.01cm (based on only two observed values). As previously stated, the mild malnutrition group had the reported median and IQR values (Figure 3). Excluding the overweight group (N=28) adjusted the statistics to  $H(2)=7.532$ ,  $p=0.023$ , reaching statistical significance. Sensitivity analyses employing bootstrap and Monte Carlo methods were conducted to examine the stability of these findings, given the potential influence of the small sample size in the overweight group.

The 95% CI for the proportion of significant tests, based on bootstrapped  $p$ -values, ranged from approximately 0.00079 to 0.5414, indicating substantial variability in the data. A considerable proportion of these tests (59.95%) yielded significant results ( $p<0.05$ ) (Figure 4A). Furthermore, the mean  $p$ -value derived from the bootstrap method was 0.0898 (Figure 4B). The Monte Carlo simulations demonstrated that in 65.3% of instances, the observed differences were statistically significant ( $p<0.05$ ) (Figure 4C) and the mean  $p$ -value was 0.0649 (Figure 4D). Although these results do not meet the criterion for significance ( $p<0.05$ ), it denotes a moderate trend toward statistical significance. These findings suggest that while the null hypothesis cannot be categorically




rejected based on our data set, there is a preliminary indication of potential effects worthy of future investigation. The decision to include the overweight group was driven by clinical interpretability considerations.

**Figure 3** – Box and whisker plots of quadriceps muscle thickness by nutritional status.



Note: Box and whisker plots illustrate the quadriceps muscle thickness (QMT) (cm) across four nutritional status groups classified based on mid-upper arm circumference adequacy (%). Mild malnutrition displays the highest median QMT, followed by Eutrophy, Overweight and Moderate malnutrition. An outlier, represented by a dot, is evident below Mild malnutrition box. The interquartile range (IQR) provides the distribution dispersion about the median. The Overweight group demonstrates an absence of IQR, indicating a limited data set comprising two observed values.

**Figure 4** – Distribution of *p*-values from Kruskal-Wallis test via Monte Carlo simulation.




Note: Scatter plots (Figures A and C) and histograms overlaid with a density curve (Figures B and D) illustrating the distribution of *p*-values derived from Kruskal-Wallis test based on bootstrap resampling on quadriceps muscle thickness across four distinct nutritional statuses: Moderate malnutrition, Mild malnutrition, Eutrophy, and Overweight. The assessment utilized 2,000 iterations, consistent with the Monte Carlo simulation parameters. Figure A: Further dispersion of points across the entire *p*-value range indicates variability in the results. A significant proportion (59.95%) of these tests yielded results with *p*-values less than 0.05. A dashed red line, marking  $p=0.05$ , serves as a reference point for statistical significance. Figure B: Two dashed lines of significance are evident on the graph: a red line marking the adopted significance threshold of  $p=0.05$  and an orange line indicating the mean *p*-value of 0.0898 for our data. Figure C: A significant proportion (65.3%) of these tests yielded results with *p*-values less than 0.05. A dashed red line, marking  $p=0.05$ , serves as a reference point for statistical significance. Figure D: Two dashed lines of significance are evident on the graph: a red line marking the adopted significance threshold of  $p=0.05$  and a green line indicating the mean *p*-value of 0.0649 for our data.

A significant positive correlation was observed between the QMT and the MUAC ( $p < 0.05$ ). The Spearman’s correlation coefficient (*r*<sub>s</sub>) was 0.557, indicating a fair to moderately strong correlation



(Figure 5A). Our analysis also revealed a significant positive correlation between QMT and the MUACA ( $p < 0.05$ ). The  $r_s$  for this association was 0.531, further establishing a fair to moderately strong correlation (Figure 5B).

**Figure 5** – Scatter plots of quadriceps muscle thickness and mid-upper arm circumference.



Note: Figure A: Scatter plot illustrating the relationship between quadriceps muscle thickness (cm) and the mid-upper arm circumference (cm). Figure B: Scatter plot illustrating the association between quadriceps muscle thickness (cm) and the mid-upper arm circumference adequacy (%). Each data point represents an individual measurement. The significance level was set at  $p < 0.05$ .

## DISCUSSION

### Summary of findings

The differences between the groups, particularly the moderate malnutrition group showing a significantly lower QMT, are a cause for concern, highlighting the potential relationship between higher degrees of malnutrition and muscle mass. The discovery that even with a small sample, the overweight group does not follow the trend of the moderate malnutrition group is a valuable discussion point. It reflects the complexity of the relationship between body mass and muscle mass, especially in situations involving the hospitalization of critically ill patients, as in the present study, where the nutritional status of mild malnutrition, eutrophy, and overweight exhibited higher QMT

than moderate malnutrition. Therefore, our study's statistically significant positive correlations suggest that as QMT increases, there is a concordant rise in MUAC and its adequacy. This can be vital in clinical and nutritional settings, especially when evaluating an individual's muscle mass and its relation to overall nutritional status.

#### *Body composition assessment by ultrasound*

Dual-energy X-ray absorptiometry (DEXA) is a precise and clinically applicable method for body composition assessment. It is appropriate for research settings, offering detailed analysis of lean and fat tissues. However, it is less practical in ICU settings due to high costs, the need for specialized personnel, and patient transfer requirements. In contrast, US is non-invasive, readily available, and can be performed at the bedside, making it ideal for continuous monitoring of muscle mass and quality in critically ill patients [24]. In the context of nutritional assessment of critically ill patients the need for more accurate and non-invasive method to assess body composition is increasingly on the rise in health research. Ceniccola et al. [25] reported that US is superior to other anthropometric methods for estimating nutritional status, as it is a practical tool that allows for the non-invasive quantification of tissue composition, without requiring patient mobilization. Reid et al. [26] demonstrated that US measurements are not affected by fluid accumulation or edema, validating the method for bedside patient monitoring.

Another point that has yet to be standardized in the literature is the amount of pressure needed to correctly quantify the muscle mass when using a US transducer [27]. Toledo et al. [3] point out that among researchers, minimal pressure is preferable for standardization. However, Paris et al. [22] noted that greater pressure is necessary to facilitate identification of muscle mass when edema is present. In the present study, the analysis was performed by two trained researchers who applied minimal pressure to measure the quadriceps in most patients, although greater pressure was required in the presence of severe edema. However, in these cases, the individuals were excluded from the study (Figure 1) due to impaired US image viewing, which was attributed to intramuscular or subcutaneous edema by the authors. Thus, in this study, the authors adopted the minimal pressure standard, as suggested by other researchers [3]. It is worth emphasizing that the effectiveness of this technique has been previously demonstrated in edematous patients when accessing the quadriceps muscle [28], as well as in muscles of other limbs of the body [29,30].

#### *Correlations and implications of MUAC and QMT*

In the present study, a positive correlation was established between MUAC and QMT. The MUAC measurement is a simple, practical, and low-cost method that has been validated for many years and widely used in hospitals due to its prognostic value in relation to both nutrition and mortality, particularly in resource-limited situations. Nevertheless, MUAC is not used to monitor specifically muscle composition, an important parameter in the nutritional care of critically ill patient context.

In recent studies, a reduction in the measurements of MUAC and QMT has been observed during illness and hospitalization. El-Liethy and Kamal [31] conducted a study on sarcopenic patients with and without liver cirrhosis, where the MUAC and QMT values were found to progressively decrease with the severity of the disease. Similarly, Chapple et al. [32] reported a decrease in both measures in patients between hospital admission and discharge. However, their study was limited to descriptive data, and a correlation between MUAC and QMT was not performed.

Sanz-Paris et al. [33] performed a correlation between MUAC and QMT measurements in 101 sarcopenic patients diagnosed with malnutrition, who were hospitalized for a hip fracture. Using US, they found statistically significant negative and positive correlations between MUAC and QMT echogenicity and thickness, respectively. The QMT mean value was  $2.21 \pm 0.645$  cm, with only the rectus femoris and vastus intermedius muscles used as constituents of the quadriceps. In the present study, we also used only the rectus femoris and vastus intermedius muscles as constituents of the quadriceps.

In the present study, the QMT of critically ill patients, which have suffered trauma mostly, was determined and associated with each identified nutritional status in the sample. The findings demonstrated that moderate malnutrition patients tend to exhibit lower QMT, suggesting that adequate nutrition is a crucial factor in maintaining muscle mass. In a previous clinical study, Toledo et al. [3] examined critically ill yet well-nourished adult patients in relation to BMI and a lower cut-off point (1.64 cm) for quadriceps thickness depletion was reported in mechanically ventilated subjects. However, that lower QMT value compared to our study can be attributed to the fact that most of their patients had developed sepsis, which is known to directly affect muscle loss.

#### *The role of ultrasound in nutritional monitoring*

The presented studies suggest that several variables could have potentially interfered in the body composition of the participants, thereby influencing the QMT. These variables include disease severity, inflammatory profile, length of stay, nutritional support, associated comorbidities, among others, which could explain the different results observed. In this study, most of the participants suffered from severe trauma and were on respiratory support, requiring a prolonged hospitalization period, all of which indicate a greater muscle depletion in comparison to patients presenting a less accentuated catabolic profile.

Taking into consideration the most recent studies in which ultrasound was employed to assess muscle mass composition, including the present study, it is not the primary objective to determine QMT cutoff points for patient profile. Rather, the relevance of this study is to show that the US can detect differences in muscle integrity, which changes according to nutritional status of critically ill patients, and can be used as a tool to monitor the progression of patients during hospitalization. By utilizing this technique, periodic evaluation of the QMT could enable a more precise adjustment of protein supply and other nutrients to patients at risk of muscle depletion.

#### *Limitations of the study*

Some limitations exist in this study, including the lack of information regarding drugs administered that may be associated with acute muscle loss and random patient selection based on length of stay. The decision-making in our statistical approach was complex, requiring a balance between methodological rigor and interpretative caution due to the small sample size of the Overweight group. While the initial Kruskal-Wallis test suggested a marginal statistical significance when all groups were included, the reanalysis without the Overweight group revealed a clearer statistical significance. The sensitivity analyses provided further insights. The bootstrap results highlighted the potential instability of the findings related to the Overweight group, suggesting its influence could introduce variability. Both the bootstrap method and Monte Carlo simulations presented a significant number of p-values indicating potential statistical significance in the difference

between groups, including the Overweight group. The choice was made to proceed with the inclusion of the Overweight group from the primary analyses. This decision was based on following factors: the significant *p*-values when including the Overweight group in many iterations, and the clinical consideration regarding the interpretability. We suggest more studies in this field to evaluate other variables that could imply in muscle composition changes in ICU patients as well as the monitoring the same patient in different days of hospitalization. Specifically, further research is required to analyze the distinct outcomes in overweight and obese ICU patients compared to those with other nutritional statuses, ensuring a significant data interpretation.

## CONCLUSION

The ultrasound assessment of the quadriceps muscle thickness revealed changes in muscle mass depending on nutritional status, mainly when moderate malnutrition patients were compared to other nutritional statuses, and may constitute a valuable tool for monitoring muscle wasting in critically ill patients. Also, measurement of quadriceps muscle thickness using the ultrasound may serve as a criterion for defining a more assertive diet therapy plan for recovery and maintaining nutritional status.

## REFERENCES

1. Federal Council of Medicine (Brazil). Resolução CFM nº 2271/2020. Diário Oficial da União. 2020;77(1):90.
2. van Gassel RJJ, Baggerman MR, van de Poll MCG. Metabolic aspects of muscle wasting during critical illness. *Curr Opin Clin Nutr Metab Care*. 2020;23(2):96-101. <https://doi.org/10.1097/MCO.0000000000000628>
3. Toledo DO, Freitas BJ, Dib R, Pfeilsticker FJDA, Santos DMD, Gomes BC, et al. Peripheral muscular ultrasound as outcome assessment tool in critically ill patients on mechanical ventilation: An observational cohort study. *Clin Nutr ESPEN*. 2021;43:408-14. <https://doi.org/10.1016/j.clnesp.2021.03.015>
4. Herridge MS. Legacy of intensive care unit-acquired weakness. *Crit Care Med*. 2009;37(10):S457-61. <https://doi.org/10.1097/CCM.0b013e3181b6f35c>
5. Kondrup J. Nutrition risk screening in the ICU. *Curr Opin Clin Nutr Metab Care*. 2019;22(2):159-61. <https://doi.org/10.1097/MCO.0000000000000551>
6. Nienow MK, Susterich CE, Peterson SJ. Prioritizing nutrition during recovery from critical illness. *Curr Opin Clin Nutr Metab Care*. 2021;24(2):199-205. <https://doi.org/10.1097/MCO.0000000000000728>
7. McFall A, Peake SL, Williams PJ. Weight and height documentation: Does ICU measure up? *Aust Crit Care*. 2019;32(4):314-8. <https://doi.org/10.1016/j.aucc.2018.06.005>
8. Coltman A, Peterson S, Roehl K, Roosevelt H, Sowa D. Use of 3 tools to assess nutrition risk in the intensive care unit. *JPEN J Parenter Enteral Nutr*. 2015;39(1):28-33. <https://doi.org/10.1177/0148607114532135>
9. Lohman T. The use of skinfold to estimate body fatness on children and youth. *J Phys Educ Recreat Dance*. 1987;58(9):98-103. <https://doi.org/10.1080/07303084.1987.10604383>
10. Simpson F, Doig GS. Bedside nutrition evaluation and physical techniques assessment in critical illness. *Curr Opin Crit Care*. 2016;22(4):303-7. <https://doi.org/10.1097/MCC.0000000000000324>
11. Danziato-Neto MA, Alencar ES, Carioca AAF, Daltro AFCS, Lima Junior PCB. Use of ultrasound in the rectus femoris muscle to evaluate muscle mass in hospitalized patients: A systematic review. *Braspen J*. 2022;37(1):82-100. <https://doi.org/10.37111/braspenj.2022.37.1.13>
12. Vieira L, Rocha LPB, Mathur S, Santana L, Melo PF, Silva VZMD, et al. Reliability of skeletal muscle ultrasound in critically ill trauma patients. *Rev Bras Ter Intensiva*. 2019;31(4):464-73. <https://doi.org/10.5935/0103-507X.20190072>
13. Schueda MA, Cohen M, Bach Neto JA, Kulevicz GV, Ribeiro GR, Bellolio JIA. Quadriceps or multiceps? Bibliographic review on your muscle composition. *BJDV*. 2021;7(8):81100-14. <https://doi.org/10.34117/bjdv7n8-363>

14. Pardo E, El Behi H, Boizeau P, Verdonk F, Alberti C, Lescot T. Reliability of ultrasound measurements of quadriceps muscle thickness in critically ill patients. *BMC Anesthesiol.* 2018;18(1):205. <https://doi.org/10.1186/s12871-018-0647-9>
15. Scott JM, Martin DS, Ploutz-Snyder R, Matz T, Caine T, Downs M, et al. Panoramic ultrasound: a novel and valid tool for monitoring change in muscle mass. *J Cachexia Sarcopenia Muscle.* 2017;8(3):475-81. <https://doi.org/10.1002/jcsm.12172>
16. Magalhães LM, Rossato EV, Franco Filho JW, Nedel WL. Variability in the rectus femoris muscle area and its association with clinical outcomes in critically ill patients: A prospective cohort study. *Rev Bras Ter Intensiva.* 2020;32(1):156-8. <https://doi.org/10.5935/0103-507x.20200023>
17. Vallet H, Guidet B, Boumendil A, De Lange DW, Leaver S, Szczeklik W, et al. The impact of age-related syndromes on ICU process and outcomes in very old patients. *Ann Intensive Care.* 2023;13(1):68. <https://doi.org/10.1186/s13613-023-01160-7>
18. Blackburn GL, Thornton PA. Nutritional assessment of the hospitalized patient. *Med Clin North Am.* 1979;63(5):11103-15.
19. Cartwright MS, Kwayisi G, Griffin LP, Sarwal A, Walker FO, Harris JM, et al. Quantitative neuromuscular ultrasound in the intensive care unit. *Muscle Nerve.* 2013;47(2):255-9. <https://doi.org/10.1002/mus.23525>
20. Bielemann RM, Gonzalez MC, Barbosa-Silva TG, Orlandi SP, Xavier MO, Bergmann RB, et al. Estimation of body fat in adults using a portable A-mode ultrasound. *Nutrition.* 2016;32(4):441-6. <https://doi.org/10.1016/j.nut.2015.10.009>
21. Souza R, Donadio M, Heinzmann-Filho J, Baptista R, Pinto L, Epifanio M, et al. Use of ultrasonography to evaluate muscle thickness and subcutaneous fat in children and adolescents with cystic fibrosis. *Rev Paul Pediatr.* 2018;36(4):457-65. <https://doi.org/10.1590/1984-0462/2018;36;4;00015>
22. Paris MT, Mourtzakis M, Day A, Leung R, Watharkar S, Kozar, R et al. Validation of Bedside Ultrasound of Muscle Layer Thickness of the Quadriceps in the Critically Ill Patient (VALIDUM Study). *JPEN J Parenter Enteral Nutr.* 2016;41(2):171-80. <https://doi.org/10.1177/0148607116637852>
23. Chan YH. *Biostatistics 104: Correlational analysis.* Singapore Med J. 2003;44(12):614-9.
24. Lopez-Ruiz A, Kashani K. Assessment of muscle mass in critically ill patients: Role of the sarcopenia index and images studies. *Curr Opin Clin Nutr Metab Care.* 2020;23(5):302-11. doi:10.1097/mco.0000000000000673
25. Ceniccola GD, Castro MG, Piovacari SMF, Horie LM, Correa FG, Barrere APN, et al. Current technologies in body composition assessment: Advantages and disadvantages. *Nutrition.* 2019;62:25-31. <https://doi.org/10.1016/j.nut.2018.11.028>
26. Reid CL, Campbell IT, Little RA. Muscle wasting and energy balance in critical illness. *Clin Nutr.* 2004;23(2):273-80. [https://doi.org/10.1016/S0261-5614\(03\)00129-8](https://doi.org/10.1016/S0261-5614(03)00129-8)
27. Weinel LM, Summers MJ, Chapple LA. Ultrasonography to measure quadriceps muscle in critically ill patients: A literature review of reported methods. *Anaesth Intensive Care.* 2019;47(5):423-34. <https://doi.org/10.1177/0310057X19875152>
28. Özdemir U, Özdemir M, Aygencel G, Kaya B, Türkoğlu M. The role of maximum compressed thickness of the quadriceps femoris muscle measured by ultrasonography in assessing nutritional risk in critically-ill patients with different volume statuses. *Rev Assoc Med Bras.* 2019;65(7):952-8. <https://doi.org/10.1590/1806-9282.65.7.952>
29. Campbell I, Watt T, Withers D, England R, Sukumar S, Keegan M, et al. Muscle thickness, measured with ultrasound, may be an indicator of lean tissue wasting in multiple organ failure in the presence of edema. *Am J Clin Nutr.* 1995;62(3):533-9. doi:10.1093/ajcn/62.3.533
30. Mul K, Horlings CG, Vincenten SC, Voermans NC, van Engelen BG, van Alfen N. Quantitative muscle MRI and ultrasound for facioscapulohumeral muscular dystrophy: Complementary imaging biomarkers. *J Neurol.* 2018;265(11):2646-55. doi:10.1007/s00415-018-9037-y
31. El-Liethy NE, Kamal HA. Value of ultrasound in grading the severity of sarcopenia in patients with hepatic cirrhosis. *Egypt J Radiol Nucl Med.* 2021;52(1). <https://doi.org/10.1186/s43055-021-00638-3>
32. Chapple LA, Gan M, Louis R, Yaxley A, Murphy A, Yandell R. Nutrition-related outcomes and dietary intake in non-mechanically ventilated critically ill adult patients: A pilot observational descriptive study. *Aust Crit Care.* 2020;33(3):300-8. <https://doi.org/10.1016/j.aucc.2020.02.008>

33. Sanz-Paris A, González-Fernandez M, Hueso-Del Río LE, Ferrer-Lahuerta E, Monge- Vazquez A, Losfablos-Callau F, et al. Muscle thickness and echogenicity measured by ultrasound could detect local sarcopenia and malnutrition in older patients hospitalized for hip fracture. *Nutrients*. 2021;13(7):2401. <https://doi.org/10.3390/nu13072401>

## ACKNOWLEDGMENTS

The research team would like to express its gratitude to the medical, nursing, and physiotherapy teams at Doutor José Frota Institute for their support and assistance in the management of patients during the course of this study.

## CONTRIBUTORS

Conceptualization: MA DANZIATO-NETO, PSS CALDAS, and CRS CÂMARA. Formal Analysis: MA DANZIATO-NETO and AAF CARIOCA. Investigation: MA DANZIATO-NETO, PSS CALDAS, JMC RÉGO, and CRS CÂMARA. Methodology: MA DANZIATO-NETO, PSS CALDAS, JMC RÉGO, and CRS CÂMARA. Project administration: CRS CÂMARA. Visualization: MA DANZIATO-NETO and CRS CÂMARA. Writing–original draft: MA DANZIATO-NETO, PSS CALDAS, JMC RÉGO, and CRS CÂMARA. Writing–review and editing: MA DANZIATO-NETO, AAF CARIOCA, and CRS CÂMARA.