

FLORA DA RESERVA DUCKE, AMAZONAS, BRASIL: LABIATAE

Raymond M. Harley¹

Labiatae Martinov, Tekhno-Bot. Slovar. 355. 1820.

Epling, C. 1935-7. Synopsis of South American Labiatae. Repert. Spec. Nov. Regni Beih. 85: 1-341.

Epling, C. 1949. Revision del género *Hyptis* (Labiatae). Revista Mus. La Plata 7: 1-497.

Harley, R. M.; Atkins, S.; Budantsev, A.; Cantino, P. D.; Conn, B.; Grayer, R.; Harley, M. M.; Kok, R.; Krestovskaja, T.; Morales, A.; Paton, A. J.; Ryding, O. & Upson, T. 2004. Labiatae. In J. W. Kadereit (ed.). The families and genera of vascular plants (K. Kubitzki, ed. in chief), vol. 7. Pp. 167–275.

Herbs, shrubs or rarely **trees**, usually with glands or glandular or non-glandular hairs (these simple in species from Flora area) and aromatic, stems often 4-angled. **Leaves** exstipulate, usually simple and opposite. **Inflorescence** thyrsoid, the cymes borne in the axils of bracts or upper leaves and often much modified by reduction or contraction to form false whorls (verticillasters), or forming terminal spikes, heads or capitula, or, rarely, by reduction, racemose. Bracts similar to or often markedly different from the leaves, bracteoles usually present, in capitulate inflorescences bracts or bracteoles often forming an involucre. **Flowers** bisexual or functionally pistillate in gynodioecious plants, or, rarely, plants dioecious. Calyx usually 5-lobed, the lobes subequal to 2-lipped, with 3 posterior and 2 anterior lobes, but often modified in various ways and sometimes the upper or both lips entire; calyx 5–20-veined, corolla gamopetalous, zygomorphic and bilabiate, the posterior (upper) lip indistinctly 2-lobed to entire, falcate or straight, flat to concave or hooded, anterior (lower) lip usually \pm 3-lobed, longer or shorter than the posterior, the median lobe flat to concave or (in

Hyptidinae) cymbiform, compressed; sometimes the corolla \pm actinomorphic. Stamens 4, adnate to the corolla, didynamous, or 2 fertile with 2 staminodes usually present, included or exserted from the corolla and often oriented towards the upper or lower lip of the corolla. Anther-thecae 2- or 1-celled, parallel or divergent, sometimes confluent, the connective sometimes extended or greatly elongated (*Salvia*). Ovary superior, bicarpellate and 4-ovulate, deeply 4-lobed, each lobe with a single ovule. Style usually gynobasic, shortly bifid above. **Fruit** of 4 (rarely fewer) usually dry nutlets, mucilaginous or not when wetted. Seeds with scanty or no endosperm.

Recent studies indicate that many genera formerly included in the Verbenaceae, would be better placed within the Lamiaceae, a situation now generally adopted (Harley *et al.* 2004). However, the present work maintains the traditional delimitation of these two families.

Cosmopolitan, especially frequent in warm-temperate to tropical savanna and submontane regions; *ca.* 230 genera and over 5000 species, 2 genera and 3 species in the flora area.

Key to the genera of Labiatae in Reserva Ducke

1. Nutlets cymbiform, concave on the internal face with a fimbriate, involute margin. Calyx-lobes broadly triangular, strongly reflexed in fruit. Viscous herbs with violet-blue flowers in lax heads, bracteoles not in a distinct involucre *Marsypianthes*
- 1'. Nutlets \pm ovoid, never concave. Calyx-lobes not broadly triangular, subulate or filiform. Herbs with white flowers in compact capitula with a distinct involucre of bracteoles *Hyptis*

¹Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, U.K

1. *Hyptis*

Hyptis Jacq., Collectanea 1: 101. 1786 (1787).

Mesosphaerum P. Browne, Civ. Nat. Hist. Jamaica: 257. 1756.

Often aromatic **shrubs, sub-shrubs**, perennial or rarely annual **herbs** of various habit, stems erect to prostrate. **Leaves** simple to lobed or rarely pinnatifid. **Inflorescence** thyrsoid, of bracteate, cymose clusters of flowers, forming congested or rarely diffuse panicles, or verticillasters or capitula, these often surrounded by an involucre of bracteoles and sometimes aggregated to form pseudoracemes or spikes, bracteoles present. **Flowers** with calyces usually 10-veined, with 5 usually \pm subequal lobes, these sometimes unequal or obsolescent; calyx tube straight or curved, usually strongly accrescent in fruit, rarely inflated. Corolla often white, sometimes

pink-spotted on posterior lip, otherwise pink, purplish or blue, very rarely red, 2-lipped, the posterior lip 2-lobed or emarginate, the anterior lip 3-lobed, the outer lobes spreading, the median lobe cymbiform and usually compressed, with thickened hinge at base, at first enclosing the declinate stamens to form an explosive pollination mechanism, later becoming reflexed; corolla tube cylindrical, usually widening slightly above. Stamens 4, declinate; anthers 2-thecae, the thecae parallel, confluent. Stylopodium absent or rarely present. **Nutlets** usually ovoid, but sometimes elongate or flattened.

About 290 species extending from Southern U.S.A., Mexico southwards to the Caribbean and South America, a few species extending to Africa, Asia, Australasia and the Pacific. Two species in the flora area.

Key to the species of *Hyptis* in Reserva Ducke

1. Leaves ovate, membranous, subglabrous to sparsely hairy beneath, with spreading hairs; petioles 5–15 mm. Stems with spreading hairs. Style jointed above the developing nutlets, the basal part (stylopodium) persistent in fruit. Fruiting capitula *ca.* 1 cm diam., on peduncles up to *ca.* 2 cm long; calyx lobes filiform, delicate, to 1.5 mm long 1. *H. atrorubens*
- 1'. Leaves ovate-elliptic to rarely lanceolate, somewhat coriaceous, often densely hairy beneath, hairs appressed, sometimes restricted to the veins; petioles indistinct, very short. Stems uniformly appressed-hairy. Style not jointed, falling in entirety as fruit ripens. Fruiting capitula 15–20 mm diameter on peduncles 2–7 cm; calyx-lobes subulate, 1.5–3.5 mm long 2. *H. lantanifolia*

1.1 *Hyptis atrorubens* Poit., Ann. Mus. Natl. Hist. Nat. 7: 466, t. 27 fig. 3. 1806.

Mesosphaerum atrorubens (Poit.) Kuntze, Revis. Gen. Pl. 2: 525. 1891.

Herb, 15–50 cm high with stems semi-prostrate to erect, often growing through surrounding vegetation. **Leaves** to 3.5 cm, ovate, aromatic, thinly hairy, with apex obtuse, base cuneate, margin regularly crenate-serrate, petiole 5–15 mm. **Flowers** in shortly pedunculate capitula *ca.* 1 cm diam., surrounded at the base by an involucre

of broadly elliptic acuminate bracteoles, often reddish-tinged, peduncles to 2 cm long. Calyx-lobes filiform with broad truncate sinuses between them. Corolla white or rarely very pale lilac with small pink spots on upper lip. Style jointed above the developing nutlets, the basal part (stylopodium) persistent in fruit. **Nutlets** ovoid, *ca.* 1 mm long.

Mexico, Central America, West Indies, Tropical South America, West Africa.

Wet forest margins and by water.

10.II.1995 (fl) Costa, M. A. S. & Assunção, P. A. C. L. 139 (BM G IAN INPA K MBM UB US); 27.IV.1995 (fl) Costa, M. A. S. et al. 215 (INPA K MG MO NYR RB SPU); 26.IV.1981 (fl) Lowe, J. 4102 (INPA).

1.2 *Hyptis lantanifolia* Poit., Ann. Mus. Natl. Hist. Nat. 7: 468, t. 29. 1806.

Mesosphaerum lantanifolium (Poit.) Kuntze, Rev. Gen. Pl. 2: 525. 1891.

Perennial **herb**, stems decumbent to erect, often weak-stemmed, to 80 cm., appressed-hairy. **Leaves** ovate-elliptic to lanceolate, almost glabrous to densely hairy on the upper surface of the lamina, usually more densely so beneath, with conspicuous appressed hairs on the prominent veins, the base attenuate into a short petiole, longer on the lower leaves, the apex acute, margin serrulate. **Flowers** in often long-pedunculate capitula from the axils of the 2 or 3 uppermost pairs of leaf-like bracts. Involucral **bracteoles** firm, ovate-lanceolate, densely appressed-hairy. Calyx pale at first becoming ferruginous, with rigid subulate teeth, sinuses narrow. Corolla white with pale lilac lobes, the upper lip spotted. Style not jointed above the nutlets, caducous from the base. **Nutlets** ovoid, ca. 1.2 mm long.

Widespread in the Neotropics from Mexico and the Caribbean, southwards through Central America and the Guianas to Brazil, Peru and Bolivia.

Wet forest margins and by water.

27.IV.1995 (fl) Costa, M. A. S. et al. 214 (BM G IAN INPA K MBM UB UEC US); 13.V.1996 (fl) Costa, M. A. S. et al. 513 (COL INPA K MG MO NYR RB SPU); 17.IV.1997 (fl) Hopkins, M. J. G. et al. 1628 (IAN INPA K SPF VEN).

In both species, a range of material, from Amazonia and elsewhere, was used to complete the descriptions of fruiting material.

2. *Marsypianthes*

Marsypianthes Mart. ex Benth., Labiat. Gen. Spec. 64. 1833.

Marsypianthus in generic conspectus, but corrected in text and index.

Perennial **herbs** or **subshrubs**, usually viscid and aromatic with softly membranous leaves, stems erect or procumbent to ascending. **Inflorescences** axillary, in subglobose, cymose, many- or rarely few-flowered heads. **Flowers** shortly pedicellate and subtended by long, curved, linear to subulate bracteoles, scarcely forming a well-defined involucre. Calyx campanulate to broadly funnel-shaped; calyx-lobes sub-equal, broadly lanceolate to triangular, spreading in fruit, not densely hairy in throat. Corolla violet-blue, 2-lipped, with slender tube and ± spreading lobes, the posterior lip 2-lobed or emarginate; the anterior lip 3-lobed, the outer lobes spreading, the median lobe saccate, hinged at base and at first enclosing the declinate stamens to form an explosive pollination mechanism, later deflexed. Stamens 4, declinate: anthers 2-theous with parallel thecae. **Nutlets** ovate, smooth on outer surface, the inner concave-cymbiform with involute, fimbriate margin.

About 6 species, chiefly in Central Brazil, but with one species, extending throughout the Neotropics.

The genus can easily be recognized in fruit, the nutlets having the appearance of minute tortoises.

2.1 *Marsypianthes chamaedrys* (Vahl) Kuntze in Revis. Gen.Pl: 524. 1891.

Clinopodium chamaedrys Vahl, Symb., 3: 77. 1794.

Marsypianthes hyptoides Mart. ex Benth., Labiat. Gen. Spec.: 64. 1833.

Viscid **herb** to 75 cm with stems prostrate to erect, with spreading, weakly villous, often gland-tipped hairs. **Leaves** ovate to ovate-lanceolate, apex acute, base

cuneate to truncate and slightly attenuate along petiole, hairy with often gland-tipped hairs, petiole 3–9 mm. **Flowers** borne in pedunculate heads 1.2–2 cm diam., peduncles 5–17 mm, from the axils of leaf-like bracts, bracteoles very narrowly elliptic to linear or filiform, often bluish or purple-tinged; calyx-tube funnel-shaped, lobes acute, broadly triangular, corolla violet-blue or pale lilac, with very small lower lip.

Fruiting calyx with lobes connivent at first, later spreading and becoming strongly reflexed to expose nutlets. **Nutlets** 2–3 mm long, dorsal surface smooth, pale, ventral surface concave and with an incurved fimbriate margin.

Throughout tropical America.

A weedy species of disturbed habitats.

3.II.1995 (bd) *Costa, M. A. S. & Nascimento, J. R. 123* (INPA KMG NY SP).