

REVISÃO

PENTAVALENT ANTIMONIALS: OLD DRUGS FOR NEW DISEASES

Philip Davis Marsden

INTRODUCTION

Clinicians like myself feel a sense of pressure because things change so slowly for the patient in spite of the volume of current research. This is currently the case in the chemotherapy of leishmaniasis where the drug formulations used in the first line treatment have not changed for four decades. Neither is a practical alternative on the horizon. There is no money to be made in treating the poor people of rural areas suffering from this disease. The small efforts that have been extended in this direction have not met with success. Therefore we must try to use antimonials more efficiently for the benefit of the patient and this is the chief purpose of this review.

HISTORY

We are indebted to Dr. Edgard de Cerqueira Falcão²³ for a monograph summarising the contributions of Gaspar Vianna with a collection of his scientific papers. The original communication on the use of tartar emetic in cutaneous and mucosal leishmaniasis is reproduced⁷². Also, in a less well known paper, Vianna illustrates a very good result in a patient with severe mucosal disease⁷³. He used a 1% solution of tartar emetic although the origin of the drug is not stated. Early preparations of tartar emetic were made by allowing antimony goblets to stand with white wine and extracting the supernatant. Confirmation of Vianna's favourable results followed in several Brazilian publications^{6 22 69}. In 1915 the effectiveness of tartar emetic in the treatment of infantile kala-azar in Italy was reported¹⁹. The drug was then widely used during the epidemic of kala-azar on the tea estates in Assam⁴⁵.

Núcleo de Medicina Tropical e Nutrição, Universidade de Brasília - 70910 Brasília-DF, Brasil.

Recebido para publicação em 23/5/85.

The main support over the years for these clinical studies has been from a United States Public Service Grant (AI 16282), administered by the Department of International Medicine, Cornell Medical College, New York.

For those of us who have used tartar emetic it is obvious that its severe side effects made an alternative desirable. Bramachari in 1920 developed the first pentavalent antimonial: urea stibamine⁵⁵. A more stable pentavalent antimonial by Schmidt in Germany in 1936 under the name Solustibosan (sodium antimony (V) gluconate⁶³) had the advantage over the earlier Neostibosan in that it was stable in water and could be stored in solution until ready for use. Also it could be given intramuscularly, intravenously and subcutaneously since it was not irritant to tissues even in a hypertonic solution. It was first produced by Bayer and shown to be effective in kala-azar in India and China^{36 46}. The Wellcome product Pentostam is sodium antimony gluconate. An alternative pentavalent antimonial, Glucantime (N-methyl-glucamine antimoniate) was developed at the end of the Second World War. Early studies of its efficacy in both cutaneous and visceral leishmaniasis appeared in French literature^{21 57}. Its value in patients with mucosal lesions was reported from Brazil in 1952¹⁵. Today Glucantime is produced by Rhodia in Brazil and is widely used. Pentostam is not available here.


The mode of action of these drugs in leishmaniasis is unknown.

Glucantime and Pentostam

These are the only two pentavalent antimonials available commercially. Both are initially manufactured as white powders (approximately 34% antimony content) and are readily soluble in water. Both solutions are best kept in the dark since they may undergo chemical change if exposed to the light.

Since they don't have defined chemical formulae they are difficult to standardise. Steck⁶⁷ gives the formulae for the two compounds illustrated in Figure 1. It has been suggested that toxicity is not necessarily related to antimony content³¹. Activity and toxicity may vary from batch to batch³. The author has personal experience of this with both drugs. It is worth citing a comment on the pentavalent antimonials³².

FIGURE 1



"These drugs, under whatever name, are simply solutions of antimony pentoxide in a highly hydroxylated organic medium and making them is cookery, not chemistry. One can make such a drug to contain any amount of antimony, and to have any degree of toxicity; one has to choose a method of making it which provides a reasonable product, with reasonable activity and reasonably low toxicity. When one puts it into a patient about 80 or 90% of the antimony it contains is excreted by the kidneys in the first few hours, so one is relying on what is left in the animal, soaked into the tissue cells, which later reduce it to trivalent antimony (Goodwin & Page 1943). This is the killer, hopefully, of the parasite but unfortunately sometimes of the patient." A variation in anti-leishmanial activity in different ampoules of Glucantime has been noted in an *in vitro* system⁴³. A reevaluation of this problem is long overdue since pentavalent antimonials remain the drugs of choice¹. There appears to be little difference in action between Pentostam and Glucantime⁸ although a good comparative trial has yet to be done.

Glucantime is marketed in 5 ml ampoules containing 1.5 grams of the drug meglumine antimoniate and 425 mg pentavalent antimony (the equivalent of 85 Sb^v mg per ml). Pentostam is presented in 100 ml flasks each ml containing 100 mg Sb^v. The ampoule has an advantage over the flask for treatment in the field since ampoules are less liable to contamination by local people applying repeated injections for patients. These applicators usually can only give intramuscular injections. The volume using Pentostam is 15% less, reducing local pain a little when large doses are applied intramuscularly.

It is difficult to get accurate information on sales of these drugs from the respective companies but it can't be that good, since both companies have tried to terminate manufacture, possibly because of the difficulties of standardising these products. Only an outcry from the medical profession, who have no real alternative for treatment, prevented this measure. Certainly in our field area the price of one ampoule of Glucantime to the patient has been as high as two American dollars after various intermediaries have taken their cut. In June 1984 a bottle of Pentostam cost £63 over the counter in England (RA Neal: personal communication). This explains why patients forced to self treatment buy so little Glucantime; often less than ten ampoules. They simply cannot afford more. Our recommendation for a patient with mucosal disease weighing 60 kilograms in the initial treatment course is 120 ampoules of Glucantime as a total dose. Governments are aware of this problem of cost. India and China faced with epidemics of kala-azar made their own pentavalent antimonial. Brazil recently completed the last steps of Glucantime fabrication at a government level but the resulting product was not satisfactory. In Brazil and Venezuela the ministry of health provides Glucantime free to applying physicians. We benefit greatly from this arrangement in our field clinic in Três Braços, Bahia, but even for us, a field project with government support, often Glucantime is not available. The cost buying hundred of thousands of ampoules must raise difficulties when considering other health priorities. An attempt has been made to map in Brazil the importance of leishmaniasis based on notifications and requests for Glucantime but this certainly underestimates the size of the problem²⁵.

In 1984 SUCAM bought 392,329 ampoules of Glucantime. The Ministry of Health has calculated it will buy 600,000 ampoules in 1985 (JBF Vieira: personal communication). Rhodias' 1985 price to SUCAM per ampoules is 6,732 cruzeiros (about 1.2 American dollars). How much profit Rhodia makes per ampoule is not known. Wellcome have no interest in marketing Pentostam in Brazil. The author knows of two attempts by the Brazilian government to produce the drug neither of which has been successful. India is producing a satisfactory sodium stibogluconate (RA Neal: personal communication) and this might be a cheaper source of drug, if demand continues to increase.

Absorption, distribution and excretion

Some aspects have been reviewed by Beveridge³. Many antimony drugs like tartar emetic are too irritant to be given by mouth but Panayotatou⁵³ cured a case of infantile kala-azar by instillation per rectum of a trivalent salt. In our current situation perhaps her observations should be followed up and the value of the oral route for pentavalent antimonials reviewed in suitable animal experiments.

Parenteral pentavalent antimony achieves peak blood levels within an hour. Intramuscular injection results in a slightly lower, later peak than intravenous application but this makes little difference¹². The drug is rapidly excreted in the urine^{28 51}. After 6 hours blood levels have fallen to less than 1% of peak values and more than 80% has been excreted in the urine⁵⁹. Thus patients are voiding the greater part of the drug they receive. This raises the interesting possibility of interfering with this rapid excretion and therefore using smaller doses to achieve the same effect. Because of the potential toxicity of antimonials this implies another series of animal studies. The hamster is not a good model for such studies since the kidney is composed almost entirely of long loop nephrons. The dog would be a more suitable model to mimic the human situation.

About 12% of the antimony dose is retained with a half-life of 32.8 ± 3.8 hours¹². With repeated injections a rise in the base line level of retained antimony can be observed over the initial five days of treatment. Some of this antimony is thought to be reduced to the trivalent state³⁰ and trivalent antimony compounds may be less effective than pentavalent in killing *Leishmania*³⁵. Much the retained antimony is concentrated in the liver and spleen^{27 58}. These is no

information on how much of this antimony is present in the skin or mucous membranes although it is probably a very small amount. Yet in skin or mucosal lesions it is this fraction that must be important. There is some rationale then for attempting local antimony treatment in skin ulcers^{1 65}. Some success in animals has been noted with Pentostam cream⁴² which is now being tried in man. Obviously such local treatment is insufficient for organisms causing systemic disease or those which produce metastatic lesions.

It is the antimony retained in the tissues which is thought to be responsible for toxic effects. Trivalent compounds bind more to tissues (*eg.* red cells) and their toxicity is marked in comparison to pentavalent compounds that are rapidly excreted in the urine. For this reason pentavalent antimonials are recommended today in higher uninterrupted doses than in the past.

Complications of antimony therapy

While the toxicity of pentavalent antimonials is much less than trivalent compounds it is not negligible. Most side effects appear at the end of courses of therapy suggesting they are the result of cumulative effects of tissue antimony. However there are exceptions. We have seen a single intravenous dose of 5 ml Pentostam given to a man with cutaneous leishmaniasis result in profound cardiovascular collapse with marked fall in blood pressure, sweating, and clouding of consciousness. A similar dose the next day gave the same result. There was no history of previous antimony therapy especially Pentostam. For this reason we now advise a 1 ml intravenous test dose before commencing a course of therapy. We have reported in the literature anuria following a total dose of only 33 grams of Glucantime drug (9.3 grams Sb^v)⁶¹. Recently we have seen a woman with mucosal disease who developed such severe shoulder arthralgia after 8 days Pentostam therapy she could not raise her arms. This improved within a day of cessation of therapy. Further attempts to commence antimony therapy resulted in her developing this symptom after a single dose of antimony and treatment with this drug had to be abandoned⁷.

An occasional complication occurring during the first week of treatment is oedema and erythema of granulomas, associated sometimes with haemorrhage, and we have illustrated and reported this finding in mucosal disease⁶⁰. This is a particularly serious complication if the laryngeal aperture is already reduced. For this reason we recommended that a

Table 1 – Detailed side effects using 5 different treatment schedules in mucocutaneous leishmaniasis.

Side effects	A	B	C	D	E
	30 cases	20 cases	18 cases	21 cases	7 cases
Arthralgia	17	18	2	10	
Myalgia	8	9		4	
Anorexia	7	6	8	5	4
Nausea	3		8	2	1
Vomiting	2	1	1	1	
Fullness	1				
Epigastric pain	1				
Waterbrash		1			
Abdominal pain		1			
Itching	1				1
Fever	2				
Weakness	2				1
Headache	3		2	2	4
Dizziness			1		
Palpitations	1			1	
Insomnia			2		
Nervousness		1			
Pyrogenic shock					4
Facial oedema		1			
Herpes zoster	1		2		

A) Glucantime 28mg/kg/day Sb^V for 10 to 12 days in series with intervals of 15 days.

B) Glucantime or Pentostam 20mg/kg/day of Sb^V during a mean of 30 days.

C) Glucantime and Nifurtimox (Lampit) in doses of 29mg/kg/day Sb^V for 15 days and 10mg/kg/day for 30 days respectively.

D) Pentostam 10mg/kg/day of Sb^V for 30 days.

E) Amphotericin B in a mean of 2.3 grams total dose.

tracheostomy set by the bed of the patient during the first week of therapy with facilities to do an emergency tracheostomy. A similar syndrome occurs with amphotericin B therapy so it is not a drug effect. One possibility is that the death of the tissue *Leishmania* promotes this reaction in a manner similar to the Jarisch – Herxheimer reaction. Although corticosteroids have been recommended to control such reactions we have had no occasion to use these to date and they could have adverse effects^{5 35}. Simply stopping the drug for a few days is sufficient as this phenomenon never recurs.

Sampaio⁶² has recently analysed the occurrence of symptomatic side effects with four different antimonial schedules (Table 1) among patients admitted to our university hospital in Brasilia. Common symptomatic side effects are arthralgia,

myalgia, anorexia, nausea, vomiting and headache¹. These side effects are commonest with high continuous doses of antimony (Table 2). Herpes infection of the skin is a recognised complication of heavy metal therapy and occurred in our earlier series⁶².

A granulocytosis has been noted⁴ and we have seen low white blood cell counts develop while on pentavalent antimonial therapy.

In patients receiving antimonials the biochemical side effects mentioned in Table 2 were rises in the transaminases and alkaline phosphatase suggesting hepatocellular dysfunction⁵². Only in two patients taking antimonials was a rise in blood urea the biochemical abnormality. Glomerular lesions due to toxicity are very rare in our experience but a tubular defect is common with high dose antimony⁷⁰. We have

studied tubular function after antimony therapy and it recovers rapidly in nearly all patients⁷¹. The tubular defect appears to be based on two mechanisms, interference with the action of antidiuretic hormone and a direct toxic effect on the tubular cells²⁶. If lower doses of antimony are used this renal complication is uncommon³⁴.

Electrocardiographic abnormalities are also listed in Table 2 and appear to be more frequent with schedule A. In our early hospital studies we tried to give this dose schedule continuously but reported electrocardiographic changes in all three patients after 20 days continuous therapy⁶⁰. 28 mg Sb^v/kg/day is approaching the toxic single dose of 30 mg Sb^v/kg/day¹. The frequency of cardiac arrhythmias is related to the total daily dose of antimony¹⁴. The common ECG changes are T wave inversion and a prolonged QT interval but rarely severe even fatal arrhythmias have occurred^{14 66}. It must be emphasised that all these side effects occur at high dosage in the patient with normal organ function; at lower doses electrocardiographic effects are not seen⁴⁴.

Many of these side effects occur together with maximal therapy. An illustration is our patient who resisted 30 days continuous Pentostam therapy (10 mg Sb^v/kg/day) and 2.5 grams amphotericin B but was eventually cured by 85 days therapy at a daily dose of 20 mg Sb^v/kilogram per day. On the 85th day he had

Table 2 – Side effects of 5 different treatment schedules in mucosal leishmaniasis.

	A	B	C	D	E
% Symptomatic	57	80	66	66	100
% Biochemical	23	26	37	26	40
% Abnormal ECGs	40	30	20	20	0

Treatment schedule A – 28 mg Sb^v/kg/day for 10-12 days repeated 3 times with an interval of 15 days.

Treatment schedule B – 20mg Sb^v/kg/day for a mean of 30 days.

Treatment schedule C – 28mg Sb^v/kg/day for 15 days plus oral Nifurtimox 10mg/kg/day for 30 days.

Treatment schedule D – 10mg Sb^v/kg/day for 30 days.

Treatment schedule E – Amphotericin B mean total dose 2.3 grams.

arthritis with effusion of the left wrist and shoulder, a palpable liver and raised transaminase and alkaline phosphatase levels, diffuse alterations of repolarisation on the electrocardiogram and a renal tubule concentrating defect⁴¹.

Toxic encephalitis associated with heavy metal therapy we have not seen nor can I trace a reference in the literature associated with pentavalent antimonials. One elderly lady in the field area who was given a daily dose in error exceeding 30 mg Sb^v/kg/day of Glucantime developed peripheral neuritis and exfoliative dermatitis. Fortunately both improved after cessation of therapy. Perhaps it is worth mentioning that we have never permanently prejudiced or lost a patient as a result of pentavalent antimony therapy in spite of using relatively high doses in over 1,000 patients.

Really the situation with regard to the use of antimonials, specially in the field, has not changed from the observation of Pessoa and Barretto⁵⁴ quoted below with relation to tartar emetic.

“Clinical experience in both visceral and cutaneous American leishmaniasis shows that the most satisfactory results are obtained using large doses in the shortest periods of time even if certain symptoms of intolerance are produced. We wish to call the attention of clinicians in general to this point for our experience shows that, many clinicians in the interior of the state, where the disease is prevalent, use, for fear of producing accidents due to intolerance to tartar emetic, excessively small doses of the product resulting in low treatment efficiency and a prolonged treatment time”⁵⁴.

Drug delivery in man of pentavalent antimonials

All drug doses are now expressed in milligrams of pentavalent antimony per kilogram per day so that schemes are standardised irrespective of the preparation¹.

As might be expected this is a most controversial area. Not only clinicians vary widely in their opinions as to the dose of pentavalent antimonials that should be used but even the two principal drug companies have different recommendations. For Pentostam the Wellcome Laboratories Ltd, Beckenham, England, recommend that adults (60 kg) be given intravenous or intramuscular injections of 10 mg Sb^v/kg/day for 7-10 days and that 2-3 cycles be separated with ten day rest intervals if additional treatment is necessary. For Glucantime Rhodia SA,

São Paulo, Brazil, recommend adults be given intramuscular injections of 17-28 mg Sb^v/kg/day for 10-20 days followed by a 15 day rest interval and a second cycle if needed. Both recommendations are now obsolete since the work done in Kenya showed that such rest periods are pharmacologically unsound since both drugs are rapidly excreted in the urine. This work which I have reviewed elsewhere³⁹, since it shortens treatment time, is the only practical advance for the patient in the treatment of leishmaniasis to appear in the last decade.

Yet one can sympathise with the drug companies since these recommendations are an enormous problem as evidence grows of differing susceptibilities to antimonials of various leishmanial species and the clinical syndromes caused by these parasites. Unfortunately comparative studies of different taxonomically defined leishmanial infections in man where similar doses have been used do not exist in the literature although a study is being prepared from Panama¹³. Many workers can't mount such studies. For example, in our work in Central Brazil *Leishmania braziliensis braziliensis* (Lbb) infection is so common we have very few other species identifications in patients with mucocutaneous disease. Recently I have tried to design a simple table

as a guide for antimonial use for Conn's *Current Therapy*, a didactic treatment text, and this is shown in Table 3. It represents an oversimplification and the fact is that the clinician has to exercise considerable skill in the use of antimonials taking into account the nature of the lesion, the probable infecting organism (this information is not available at the time of the treatment decision), the age and weight of the patient as well as any evidence of organ disfunction.

The smallest adult dose of Glucantime recommended in Brazil is one ampoule of Glucantime two times a week⁴⁹. While this seems homeopathic to some and cannot be recommended it is possibly the only dose available, for reasons of cost, for a child with advanced kala-azar, anaemic and malnourished who has to be treated at home in one of the *favelas* in the cities of Brazil where kala-azar is currently endemic. The fact that Brazilian kala-azar responds to such a dose shows it has a high sensitivity although the relapse rate must be high. The highest daily dose advocated in Brasil is six ampoules of Glucantime a day¹⁸. This is well over the toxic limit mentioned by the World Health Organization¹ for a single dose and it is curious that few toxic effects were observed. It cannot be recommended. A Brazilian Ministry of

Table 3 - Guide to the treatment response of different leishmanial species causing clinical syndromes with pentavalent antimonials.

Treatment response	Easy to treat Group A	More difficult Group B	Most difficult Group C
Clinical form			
Cutaneous	Limited cutaneous lesions <i>L. tropica</i> <i>L. major</i> <i>L. m. mexicana</i> <i>L. m. amazonensis</i> <i>L. peruana</i>	<i>L. b. panamensis</i> <i>L. b. guyanensis</i> <i>L. b. braziliensis</i> Post Kala-azar Dermal leishmaniasis (<i>L. donovani</i>)	Diffuse cutaneous leishmaniasis <i>L. aethiopica</i> <i>L. m. pifanoi</i> <i>L. m. amazonensis</i> Leishmaniasis recidivans Skin granuloma due to <i>L. aethiopica</i>
Visceral or mucosal forms		Visceral leishmaniasis (<i>L. donovani</i>)	Mucosal leishmaniasis <i>L. b. braziliensis</i> <i>L. b. panamensis</i> <i>L. b.?</i>

Key. L = *Leishmania* L. m. = *Leishmania mexicana* group L. b. = *Leishmania braziliensis* group.

Note. Oversimplified, this guide does not take into account the exceptions to the rule. Relapse can occur in individual patients in all categories.

Treatment Group A - 10mg Sb^v/kg/day for 10-15 days.

" Group B - 10mg Sb^v/kg/day for 30 days or 20 mg Sb^v/kg/day for 15 days.

" Group C - 20mg Sb^v/kg/day for 30 days or more 20mg Sb^v/kg/twice daily for 15 days or more.

Health recommendation is one ampoule of Glucantime daily for 7-10 days; the course being repeated if necessary. In our recent experience it appears that this will close many skin ulcers due to Lbb; but will it prevent mucosal disease? Since in the past year we have had little Glucantime, we have many patients with Lbb who had to be treated with such a schedule (7 mg Sb^v/kg/day). They are under longitudinal observation. We have used four times this dose in the endemic area of Três Braços for years (28 mg/kg/day) and in our recently published studies mucosal relapse was almost unknown on short term follow-up³⁷. The practical difficulties of giving such high doses under field conditions are considerable (20 cm³ daily intramuscular dose) but they cannot be discussed here. Prata has never seen a kala-azar patient relapse in Brazil using similar doses in over 100 patients (AR Prata: personal communication) but they do rarely occur^{20 68}.

On the whole in terms, of frequency of treatment problems in hospitals in Brazil, it is mucosal leishmaniasis which causes most concern. Table 4 shows our results with the four antimony schedules previously mentioned in relation to mucosal disease in our hospital in Brasília⁶³. Our analysis of the results of treatment with cutaneous lesions failed to show a difference between the four schedules but there is a difference between these in mucosal patients. Such data is extremely difficult to collect due to follow-up problems and as yet it does not lend itself to statistical analysis. However it suggests that 20 mg/Sb^v/kilogram body weight for a mean of 30 days is the best recommendation. Bryceson in London has recently started to use this dose twice a day to further reduce hospital stay (ADM Bryceson: personal communi-

cation). Evidently we still have to define optimal schedules for antimony use.

Both Pentostam and Glucantime are best given by slow intravenous infusion over at least three minutes. No diluent is usually necessary unless the patient develops a venous thrombosis at an inoculation site. Intramuscular injection is painful but often necessary under field conditions where no doctor can be present for daily injection. One lot of Glucantime used in Três Braços produced several muscle abscesses. Pharmacokinetic studies show little difference in peak serum levels after application by these two routes¹². An interesting study carried out at the Walter Reed Army Medical Center compared three methods of drug delivery in American soldiers with leishmaniasis. While all subjects were given the same total dose of 600 mg Sb^v/day this was administered either as a continuous 24 hours infusion, in three divided eight hourly doses or as a single intravenous dose⁵⁰. The latter method was shown to be more effective in terms of cure and incidence of subjective complaints.

In patients with evidence of renal, cardiac or hepatic disease an alternative may have to be sought depending on the clinical assessment of the patient. Patients with renal insufficiency become lethargic, anorexic, drowsy and weak after antimonial therapy⁵⁹. Alternatives are the recommended second line drugs¹ namely Pentamidine which has few renal side effects or amphotericin B which is free of hepatic toxicity and cardiac effects are only related to potassium imbalance.

Finally even high dose pentavalent antimonial will not cure all patients. We have reported a girl who

Table 4 – Cure rate of five treatment schedules in mucosal leishmaniasis.

Nº failed/ total number	A	B	C	D	E
	11/26	1/12	3/7	7/18	2/8
% Cure	58%	92%	57%	61%	75%
Mean follow up period (months)	18	8	19	12	12

Treatment schedule A – 28mg Sb^v/kg/day for 10-12 days repeated 3 times with an interval of 15 days.

” ” B – 20mg Sb^v/kg/day for a mean of 30 days.

” ” C – 28mg Sb^v/kg/day for 15 days plus oral Nifurtimox 10mg/kg/day for 30 days.

” ” D – 10mg Sb^v/kg/day for 30 days.

” ” E – Amphotericin B mean total dose 2.3 grams.

developed mucosal granulomas during antimonial treatment for her skin lesion due to Lbb⁴⁰.

Unresponsiveness

In brief recent work regarding disease response to total antimony dose suggests that high continuous antimony therapy is the best approach in recalcitrant disease for a favourable response. Syndromes such as are mentioned in group C of Table 3 require this approach. This is not to say that high dose antimony will always meet with success particularly in patients with mucosal disease^{24 40}. Anergic Hansenoid leishmaniasis will fail to respond since the patient can't mount an immune response. The importance of unresponsiveness is difficult to assess since high enough doses have not been widely employed to date.

The term unresponsive is useful since resistance is often taken to imply parasite resistance which is not necessarily the case. Patients with fatal antimony unresponsive kala-azar had parasites which showed a normal response to antimonials in animal models indicating a host defect^{20 32}. On the other hand at least in Kenyan kala-azar there is evidence in cell cultures² and in animal models of parasite resistance to antimony³³. Unfortunately similar experiments have not been possible with Lbb since it cannot be induced to grow in *in vivo* cell lines. However Moreira⁴³ has examined one of our most suspect Lbb isolates in terms of resistance and it shows diminished sensitivity in her promastigote culture system.

The World Health Organization¹ suggests we consider primary unresponsiveness as no clinical or parasitological improvement on initial therapy and secondary unresponsiveness where relapse occurs after one or more courses of apparently successful treatment. To illustrate how difficult it is to discuss this subject with the recent changes in dosage schedules India which recorded a 13% relapse rate in kala-azar is said to have reduced it to zero by doubling the antimony dose¹. Certainly a risk of minimal dose therapy is the emergence of parasite resistance to antimonial treatment.

Concluding remarks

The last two decades have seen great advances in our understanding of the taxonomy of *Leishmania* using biochemical²⁹ and immunological⁵⁶ methods. Today this is seen as an essential first step to unravelling the complexities of the epidemiology and

clinical picture of human disease. At the University of Brasília we were the first to use the monoclonal technology developed at Harvard in such field studies¹⁷. We are dealing in Três Braços, Bahia, with what appears to be virtually a monotransmission to man of Lbb. We are still defining the clinical picture based on this taxonomy. Similar data on other important parasites in Brazil such as *Leishmania braziliensis guyanensis* will doubtless be available soon. New clinical entities are being defined; for example we are preparing a report on disseminated cutaneous leishmaniasis caused by Lbb¹⁶.

In contrast to this exciting field in epidemiology, prospects for advances in the chemotherapy of leishmaniasis are gloomy. I never imagined when I qualified in medicine 25 years ago that I would review antimonial therapy in 1985. Heavy metal therapy was the mainstay of 19th century medicine. The experts in this field today are physicians treating leishmaniasis. There has been no advance in drug formulations for primary therapy that has benefited the patient for forty years. Parental pentavalent antimonials are still the drugs of choice for most of the forms of leishmaniasis.

I will not discuss why this is so, although I have definite views after confronting hundreds of patients with Lbb infections and often not having even antimony to offer them. Let us adopt a more positive approach. The fact is that there is only one pharmaceutical company and one research institute seriously involved in developing new drugs; namely the Wellcome Research Laboratories in America and England and the Walter Reed Army Institute of Medical Research in Washington. Two drugs, an 8 aminoquinoline (WR 6026)¹⁰ from the latter and Allopurinol riboside from the former⁴⁷ have entered clinical trials. Preliminary information suggests that neither may be suitable for our problems with Lbb, but even if this is not so, I may well be an old man before I see such oral preparations used in our field clinic. This is because clinical evaluation is a lengthy process and involves testing initially in kala-azar and limited skin leishmaniasis before it can be used for an organism causing metastatic disease. Also we ourselves routinely do a trial in hospital in Brasília before a drug can be taken to the field. The evaluation of each of these trials will take several years.

Another possibility is the use of liposome linked antimony which has given promising result in animals¹¹. There is some evidence that this is more effective in cutaneous lesions than the free drug⁹. The

one patient with Kenyan kala-azar unresponsive to free antimony treated with liposomes failed to respond³⁵. Both the research groups mentioned are involved in this development. Although if it ever came to human trial this approach would still imply injection therapy it would abolish the need for these painful, prolonged, injection schedules. I have no doubt that the single most important advance in the control of leishmaniasis is the development of more efficient drug therapy³⁸ in terms of an oral inexpensive treatment of short duration. Such a drug would be equally useful both in kala-azar and in the mucocutaneous infections of rural areas. With ministry support we are currently testing ketoconazole but this hardly falls into this category.

The last word is perhaps better said by the father of one of our patients in the endemic area of Três Braços where they are as familiar with Glucantime as we are with aspirin⁴⁸. He asked me why if we had been working there for ten years we were still using this drug and had not found a more convenient therapy. I explained that we were dependent in the field, at the end of the line as it were, on the research of other colleagues and nothing had appeared to date. "But will it", he insisted "so that I don't have to walk 5 kilometres with my daughter to get her injection?"

REFERENCES

1. Anonymous. The leishmaniasis. World Health Organization Technical Report Series nº 701, Geneva, 1984.
2. Berman JD. In vitro susceptibility of antimony resistant *Leishmania* to alternative drugs. *Journal of Infectious Diseases* 145: 279, 1982.
3. Beveridge E. Chemotherapy of leishmaniasis. In: Schnitzer RJ, Hawking F (ed) *Experimental chemotherapy*, vol. 1, Academic Press, New York, 1963.
4. Brahmachari PN, Maiti CR, Kor CC. Agranulocytosis following antimony therapy in kala-azar. *Journal Indian Medical Association* 25: 408-410, 1955.
5. Bryceson ADM. Therapy in man. In: Peters W, Killick Kendrick R (ed) *The leishmaniasis in biology and medicine*, Academic Press, London, in press.
6. Carini A. Três casos de leishmaniose. *Archivos da Sociedade de Medicina e Cirurgia de São Paulo* 5: 8-9, 1914.
7. Castro CN, Costa JML, Sampaio RNR, Marsden PD. Unusual side effects following pentavalent antimonial therapy. *Revista da Sociedade Brasileira de Medicina Tropical*, in press.
8. Castro RM. Tratamento da leishmaniose tegumentar americana. *Anais Brasileiro de Dermatologia* 55: 87-89, 1980.
9. Chance ML, New RRC. The use of liposomes in the treatment of experimental cutaneous and visceral leishmaniasis. In: Van de Bossche (ed) *The Host Parasite Interplay*, p. 659. Elsevier/North Holland Biomedical Press, Amsterdam, 1980.
10. Chapman WL, Hanson WL, Waits VB, Kinnamon KE. Antileishmanial activity of selected compounds in dogs experimentally infected with *Leishmania donovani*. *Revista do Instituto de Medicina Tropical de São Paulo* 21: 189-193, 1979.
11. Chapman WL, Hanson WL, Alving CR, Hendricks LD. Antileishmanial activity of liposome encapsulated meglumine antimonate in the dog. *American Journal of Veterinary Research* 45: 1028-1030, 1984.
12. Chulay J. Clinical pharmacokinetics of currently used compounds and their side effects. Informal meeting on Chemotherapy of Visceral Leishmaniasis. Mimeographed document. Nairobi, 1982.
13. Chulay JD, Oster CN, McGreevy PB, Kreutzer RD, Hendricks LD. American cutaneous leishmaniasis. Clinical presentation and problems of patient management. *Journal of Infectious Disease*, in press.
14. Chulay JD, Spencer HC, Mugambi M. Electrocardiographic changes during treatment of leishmaniasis with pentavalent antimony (Sodium stibogluconate). *American Journal of Tropical Medicine and Hygiene* 34: 702-709, 1985.
15. Corrêa A, Elizabetsky M. Ação de antimonato de N-methyl glucamina sobre as lesões mucosas de leishmaniose tegumentar americana. *Publicações Médicas de São Paulo* 22: 13-21, 1952.
16. Costa JML, Marsden PD, Llanos-Cuentas EA, Netto EM, Carvalho EM, Barral A, Rosa AC, Cuba CC, Magalhães AV, Barreto AC. Disseminated cutaneous leishmaniasis in a field clinic. Report of 8 cases. *Journal of Tropical Medicine and Hygiene*, in press.
17. Cuba Cuba CA, Marsden PD, Barreto AC, Roitman I, Vexenat A, De Lima LMP, De Sá MHP. Identification of human stocks of *Leishmania* spp. isolated from patients with mucocutaneous leishmaniasis in Três Braços, Bahia, Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 78: 708-710, 1984.
18. Cuce LC, Freitas THP, Oliveira ZNP, Sampaio SAP.

- Tratamento da leishmaniose tegumentar americana pelo antimônio pentavalente (esquema rápido). *Anais Brasileiros de Dermatologia* 57:107-108, 1982.
19. Di Christina G, Caronia G. Sulla terapia della leishmaniosi interna. *Bulletin de la Société de Pathologie Exotique* 8: 63-66, 1915.
 20. Dietze R, Araújo RC, Lima MLR, Vexenat JA, Marsden PD, Barreto AC. Ensaio terapêutico com glucantime em sagüis (*Callithrix jacchus*) infectados com uma cepa de *Leishmania donovani* aparentemente resistente ao tratamento. *Revista da Sociedade Brasileira de Medicina Tropical* 18: 39-42, 1985.
 21. Durand P, Benmussa S, Caruana M. Traitement du kala-azar par un nouveau composé stibié 2, 168 RP (premiers résultats). *Bulletin Société Medicine Hôpital Paris* 62: 399-409, 1946.
 22. Dutra e Silva O. Sobre a leishmaniose tegumentar e seu tratamento. *Memórias do Instituto Oswaldo Cruz* 7: 213-248, 1915.
 23. Falcão EC, Opera Omnia de Gaspar Vianna. Empresa Gráfica da Revista dos Tribunais, São Paulo, 1962.
 24. Furtado TA. An appraisal of chemotherapy of American leishmaniasis. *Dermatologia Ibérica Latino-Americana* 3: 257-268, 1968.
 25. Furtado T, Vieira JBF. Geografia da leishmaniose tegumentar americana no Brasil. *Anais Brasileiro de Dermatologia* 57: 135-140, 1982.
 26. Gagliardi ART, Veiga JPR, Rosa TT, Marsden PD. Pentavalent antimony inhibition of water transport by the toad bladder. *Brazilian Journal of Medical and Biological Research*, in press.
 27. Gelhorn A, Van Dyke HB. The correlation between distribution of antimony in tissues and chemotherapeutic effect in experimental leishmaniasis. *Journal of Pharmacology and Experimental Therapeutics* 88: 162-172, 1946.
 28. Gelhorn A, Tupikova NA, Van Dyke HB. The tissue distribution and excretion of four organic antimonials after single or repeated administration to normal hamsters. *Journal of Pharmacology and Experimental Therapeutics* 87: 169-180, 1945.
 29. Goldfrey DG. Molecular biochemical characterization of human parasites. In: Gilles HM (ed) *Recent Advance in Tropical Medicine*. Churchill Livingstone, Edinburgh, p. 289-319, 1984.
 30. Goodwin LG, Page JE. A study of the excretion of organic antimonials using a polargraphic procedure. *Biochemical Journal* 37: 198-209, 1943.
 31. Goodwin LG. The toxicity and trypanocidal activity of some organic antimonials. *Journal of Pharmacology and Experimental Therapeutics* 81: 224-234, 1944.
 32. Goodwin LG. Commentary. In: *Trypanosomiasis and Leishmaniasis*. Ciba Foundation Symposium 20. Elsevier, North Holland, Amsterdam, p. 331, 1974.
 33. Hanson WL, Waits VB. Relative insensitivity of a Kenyan strain of *Leishmania donovani* to pentavalent antimony therapy in hamster. *Journal of Parasitology* 69: 446-447, 1983.
 34. Jolliffe DS. Nephrotoxicity of pentavalent antimonials. *Lancet* 1: 584, 1985.
 35. Kager PA, Rees PH. Clinical aspects of kala-azar in Kenya ICG Printing Dordrecht, 1983.
 36. Kikuth W, Schmidt H. Contribution to the progress of antimony therapy of kala-azar. *Chinese Medical Journal* 52: 425-432, 1937.
 37. Llanos-Cuentas EA, Marsden PD, Lago EL, Barreto AC, Cuba CC, Johnson WD. Leishmaniose mucocutânea humana em Três Braços, Bahia. Uma área de transmissão de *Leishmania braziliensis braziliensis*. II Doença cutânea: apresentação e evolução. *Revista da Sociedade Brasileira de Medicina Tropical* 17: 169-177, 1984.
 38. Marsden PD. Selective primary health care: strategies for control of disease in the developing world. XIV Leishmaniasis. *Reviews of Infectious Disease* 6: 736-744, 1984.
 39. Marsden PD. New light on pentavalent antimonials in the treatment of leishmaniasis. *Revista da Sociedade Brasileira de Medicina Tropical* 16: 172-174, 1983.
 40. Marsden PD, Llanos-Cuentas EA, Lago EL, Cuba CC, Barreto AC, Costa JM, Jones TC. Human mucocutaneous leishmaniasis in Três Braços, Bahia-Brazil. An area of *Leishmania braziliensis braziliensis* transmission. III Mucosal disease: presentation and initial evolution. *Revista da Sociedade Brasileira de Medicina Tropical* 17: 179-186, 1984.
 41. Marsden PD, Sampaio RNR, Carvalho EM, Veiga JPR, Costa JLM, Llanos-Cuentas EA. High continuous antimony therapy in two patients with unresponsive mucosal leishmaniasis. *American Journal of Tropical Medicine and Hygiene*, 34: 710-713, 1985.
 42. McGreevy PB, Marsden PD. American trypanosomiasis and leishmaniasis in the chemotherapy of parasitic diseases. Ed. Campbell W.C., Rew R. S., Plenum Press, in press.

43. Moreira ESA. Ação do antimoníaco de N-metilglucamina sobre formas promastigotas de *Leishmania*. Masters thesis, Universidade Federal de Minas Gerais, Belo Horizonte, 1984.
44. Muller RH, Valera M, Carrasco H, Fuenmayor A, Benedetti JR, Torrealba W, Mendonza E. Valoración electrocardiográfica de la cardiotoxicidad del antimoníaco de N-metil glucamina (R Glucantime). *Arquivos do Instituto de Cardiologia Mexicana* 52: 155-9, 1982.
45. Napier LE, Muir E. Kala-azar. London, Oxford University Press, 1923.
46. Napier L. E. Chaudhuri R, Rai Chaudhuri M. N. A stable solution of antimony for the treatment of kala-azar. *Indian Medical Gazette* 72: 462-465, 1937.
47. Nelson DJ, Lafon SW, Tuttle JV, Miller WH, Miller RL, Krenitsky TA, Elion GB, Berens RL, Marr JJ. Allopurinol riboside as an antileishmanial agent. *Journal of Biological Chemistry* 254: 11544-11549, 1979.
48. Netto EM, Tada MS, Golightly L, Kalter DC, Lago E, Barreto AC, Marsden, PD. Conceitos de uma população local a respeito da leishmaniose mucocutânea em uma área endêmica. *Revista da Sociedade Brasileira de Medicina Tropical* 18: 33-37, 1985.
49. Neves J. Tratamento da leishmaniose visceral americana pelo antimoníaco de N-metil glucamina. Tese, Universidade Federal de Minas Gerais, Belo Horizonte, 1963.
50. Oster CN, Chulay JD, Hendricks LD, Pamplin CL, Ballow WR, Berman JD, Takafuji EJ, Tramont EC, Canfield CJ. American cutaneous leishmaniasis: a comparison of three sodium stibogluconate treatment schedules. *American Journal of Tropical Medicine and Hygiene* 34: 856-860, 1985.
51. Otto GF, Maren TH, Brown HW. Blood levels and excretion rates of antimony in persons receiving trivalent e pentavalent antimonials. *American Journal of Hygiene* 46: 193-211, 1947.
52. Pablos Gallego JM, Torres AC, Almagro M, De Puerta S, Garrido PL, Morales MG, Esquivias JJ. Kala-azar y anemia hemolítica autoimune. A propósito de un caso de evolución fatal por hepatotoxicidad del antimoníaco de N-metil glucamina. *Revista Clínica Española* 164: 417-420, 1982.
53. Panayotatou A. Infantile kala-azar: a case cured at Alexandria by a new method. *British Journal of Childhood Diseases* 26: 112-115, 1929.
54. Pessoa SB, Barreto MP. Leishmaniose tegumentar americana. Imprensa Nacional, Rio de Janeiro. 1948.
55. Peters W. The treatment of kala-azar – new approaches to an old problem. *Indian Journal of Medical Research (Supplement)* 73: 1-18, 1981.
56. Pratt DM, Bennett e, David JR. Monoclonal antibodies that distinguish subspecies of *Leishmania braziliensis*. *Journal of Immunology* 129: 926-927, 1982.
57. Raju S, Schneider J. Traitement d'une leishmaniose cutanée por l'antimoníaco de N-methyl glucantime (2168 RP). *Bulletin Société Française Dermatologie et Syphilis* 2: 38, 1947.
58. Rao LS, Hardy JG, Wilson CG. Tissue distribution and rate of free and liposome – encapsulated (125 Sb) sodium stibogluconate by gamma scintigraphy. *International Journal of Pharmaceutics* 17: 283-290, 1983.
59. Rees, PH, Kager PA, Keating MI, Hockmeyer WT. Renal clearance of pentavalent antimony (sodium stibogluconate). *Lancet* 2: 226-229, 1980.
60. Rocha RAA, Sampaio RN, Guerra M, Magalhães A, Cuba CC, Barreto AC, Marsden PD. Apparent glucantime failure in five patients with mucocutaneous leishmaniasis. *Journal of Tropical Medicine and Hygiene* 83: 131-139, 1980.
61. Sampaio RNR, Rocha RAA, Marsden PD, Cuba CC, Barreto AC. Leishmaniose tegumentar americana. Casuística do Hospital Escola da UnB. *Anais Brasileiro de Dermatologia* 55: 69-76, 1980.
62. Sampaio RNR. Tratamento hospitalar da leishmaniose cutâneo mucosa. Master's thesis, Universidade Federal de Minas Gerais, Belo Horizonte, 1984.
63. Sampaio RNR, Sampaio JHD, Marsden PD. Pentavalent antimonial treatment in mucosal leishmaniasis. *Lancet* 1: 1097, 1985.
64. Schimdt H. An advance in the therapeutics of pentavalent antimony: solustibosan (sodium antimony gluconate). *Journal of Tropical Medicine and Hygiene* 53: 95-102, 1950.
65. Solano AE, Hidalgo HH, Zeledon RA. Tratamiento intralesional exitoso de la leishmaniasis por *Leishmania braziliensis panamensis* com glucantime. *Medicina Cutis ILA* 12: 19-24, 1984.
66. Soto JM, Urbina A, Convit J. Comunicación preliminar sobre toxicidad cardiaca de antimonial pentavalente. *Revista Venezolana de Sanidade e Assistência Social* 39: 269-278, 1974.
67. Steck EA. The leishmaniasis. In: Jucker E (ed) *Progress in drug research*. Birkhäuser Verlag, Basel, p. 289-351, 1974.

68. Teixeira R. Experiências vividas com a leishmaniose visceral. 1954-1980. (Aspectos epidemiológicos, sorológicos e evolutivos). Doctoral thesis, Universidade Federal da Bahia, Salvador, 1980.
69. Terra F. Leishmaniose tegumentaire au Brèsil. *Boletim da Sociedade Brasileira de Dermatologia* 2-3: 58-67, 1913.
70. Veiga JPR, Wolff ER, Sampaio RN, Marsden PD. Renal tubular dysfunction in patients with mucocutaneous leishmaniasis treated with pentavalent antimonials. *Lancet* 2: 569, 1983.
71. Veiga JPR, Rosa TT, Kimachi T, Wolf ER, Sampaio RN, Gagliardi RT, Juqueira LF, Costa JML, Marsden PD. Função renal em pacientes com leishmaniose mucocutânea tratados com antimoniais pentavalentes. *Revista do Instituto de Medicina Tropical de São Paulo*, in press.
72. Vianna G. Tratamento da leishmaniose tegumentar por injeções intravenosas de tartar emético. *Anais do 7º Congresso Brasileiro de Medicina e Cirurgia* 4: 426-428, 1912.
73. Vianna G. Sobre o tratamento de leishmaniose tegumentar. *Anais Paulista de Medicina e Cirurgia* 2: 167-169, 1914.