

**LEPTOCHLOA FUSCA CULTIVATION FOR UTILIZATION OF SALT –  
AFFECTED SOIL AND WATER RESOURCES IN CHOLISTAN DESERT**

**Cultivo de Leptochloa Fusca para a utilização de solos afetados por  
sais e recursos hídricos no Deserto de Cholistan**

Farooq Ahmad

Department of Geography, University of the Punjab  
Lahore, PAKISTAN  
farooq@geog.pu.edu.pk

Artigo recebido para publicação em 15/07/2009 e aceito para publicação em 08/02/2010

**ABSTRACT:** *In the Cholistan Desert, 0.44 million hectares are salt – affected low lying and clayey in nature locally known as ‘dhars’, where rainwater as well as saline groundwater could be utilized for growing salt tolerant grasses like Leptochloa fusca as forage during summer. L. fusca is a promising candidate grass for economic utilization and better management of sodic, high pH, saline soil and saline water resources of the Cholistan desert. L. fusca is known to be a versatile, halophytic, primary colonizer, easily propagatable, perennial, nutritive and palatable forage plant species. The grass has the good biomass producing potential and can grow equally well both under upland and submerged saline soil environment.*

**Keywords:** Cholistan, *Leptochloa fusca*. Saline irrigation. Salt – affected. Soil reclamation.

**RESUMO:** *No Deserto de Cholistan, 0,44 milhões de hectares são afetados por condições de solos rasos, salinos e argilosos por natureza, sendo localmente conhecidos como ‘dhars’, onde a água da chuva, bem como as águas subterrâneas salinas poderiam ser utilizadas para o cultivo de gramíneas tolerantes a sais, como a Leptochloa fusca, como forragem durante o verão. A L. fusca é uma gramínea promissora para a utilização econômica e melhor manejo dos solos salinos, sódicos e com alto pH e dos recursos hídricos salinos do Deserto de Cholistan. A gramínea é conhecida por ser versátil, halófila, colonizadora primária, de fácil propagação, perene, nutritiva e por ser uma espécie de planta forrageira palatável. Além disso, possui um bom potencial de produção de biomassa e pode crescer igualmente tanto em ambientes de sequeiro quanto em ambientes de solos salinos inundáveis.*

**Palavras – Chave:** Cholistan. *Leptochloa fusca*. Irrigação salina. Solos afetados por sais. Recuperação de solos degradados.

---

## 1. INTRODUCTION

The Cholistan Desert (FIGURE 1) lies within the southeast quadrant of Punjab Province between

27° 42' and 29° 45' North latitude and 69° 52' and 73° 05' East longitude (FAO/ADB 1993; ARSHAD *et al.* 1995; JOWKAR *et al.* 1996; AHMAD *et al.* 2004; AHMAD 2005; 2007; 2008) and covers an area of 2.6

million hectares (AHMAD 2002), out of which 1.13 million ha comprising stable as well as non – stable sand dunes, 0.95 and 0.06 million ha consist of sandy and loamy soils respectively, while 0.44 million ha are clayey in nature, locally known as ‘dhars’. About 17% of Cholistan consist of such ‘dhars’ (TABLE 1) having flat and hard surface with salt incrustation and surrounded by sand dunes. Dhars are shallow to moderately deep, poorly drained with low vegetation, calcareous and having saline sodic fine to medium textured clayey soils. Except *Haloxylon recurvum*, other plant species can’t survive due to salinity, compaction of soil and complete inundation during rainy season. The ponded rainwater in ‘dhars’ stagnates for a period until the water evaporates (KHAN *et al.*, 1990). It is judicious to utilize the land using ground saline and surface rainwater resources for growing palatable grasses. Biological approach

for economic utilization of salt – affected soil are feasible and is the only viable method when the soil is sodic and sweet water is unavailable for irrigation (ABDULLAH, 1985).

*Leptochloa fusca* is highly tolerant of saline and sodic conditions even when irrigated by saline groundwater or ponded rainwater. After the successful cultivation of *L. fusca* in Cholistan, other palatable grasses like para grass (*Brachiaria mutica*), Rhodes grass (*Chloris gayana*), Bermuda grass (*Cynodon dactylon*) and *Sporobolus* grass species can be tested (ABDULLAH *et al.* 1990). The cultivation of salt tolerant grasses would not only provide much needed palatable forage for livestock but also improve the soil physical properties due to biological activity of grass roots.



FIGURE 1: Location map of Cholistan desert

TABLE 1: Soil Types

Soil Types	Extent (Ha)	Percentage
Sand dunes	1,133,900	44.0
Sandy soils	945,500	37.0
Loamy soils	58,700	2.0
Saline sodic clayey soils ( <i>Dhars</i> )	441,900	17.0
Total	2,580,000	100.0

Source: PADMU. – Pakistan Desertification Monitoring Unit.

## 2. GROWTH CHARACTERISTICS OF LEPTOCHLOA FUSCA

*Leptochloa fusca* is also known as *Diplachne fusca*, and is widely spread in salt affected regions of Pakistan. This forage plant is locally known as “*Kallar grass*” (salt grass). Being a tropical grass, the plant follows the photosynthetic CO<sub>2</sub> fixation process of C<sub>4</sub> – NAD – malice enzyme metabolism (Zafar and Malik 1984). It is native of saline soil which gives clear indication of its halophytic character; the plant is perennial or biennial in nature. It has been regarded as good quality forage especially in salt – affected and waterlogged areas where other superior forage species may not grow successfully.

*Leptochloa fusca* can be easily propagated and established through seed, stem cutting, root stumps or rhizomes. The grass can grow to a height of 1 – 1.5 meters with a high leaf production rate and can be grazed directly or cut for stall – feeding. This fodder is highly palatable to sheep, goats, buffaloes and cattle alike and no toxic effects of this grass during long – term consumption have been diagnosed. Moreover, it is similar to other conventional fodder regarding its nutritional status and 3 – 4 cutting within 3 months may be easily harvested, producing 20 – 40 tonnes of green fodder per ha per year or 5 – 10 tonnes per ha per cutting in salt – affected soils (SANDHU *et al.* 1981; QURESHI *et al.* 1982; SANDHU 1993). The grass grows well during the hot season from March to September with peak yields during rainy season *i.e.* July and August in Pakistan, indicating a strongly thermophilic character. The development of extensive and dense fibrous root system has been observed even in highly sodic soils (JOSHI *et al.*

1981). The penetration of roots in such soils can enhance hydraulic conductivity, microbial activity, organic matter and ultimately leaching of salts. This grass can make better growth under normal situation than under stress conditions, but it is a common observation that, in normal soils, the grass could not compete with other species and is soon eliminated due to the growth of other vegetation. Joshi *et al.* (1981) noted a decline of *L. fusca* growth due to decrease of soil sodicity, while Haq and Khan (1971) observed that *L. fusca* has a general tendency to decrease EC<sub>e</sub> (electrolyte conductivity), SAR (sodium adsorption ratio), pH (soluble ions) and even ESP (exchangeable sodium percentage) of artificially salinized soils. Malik *et al.* (1986) confirmed the utility of *L. fusca* not only as a primary colonizer of salt – affected lands, but also as ameliorative plant for the soil.

## 3. NUTRITIONAL REQUIREMENTS

It has been observed that 3 – 4 cuttings of this grass could be easily taken without the addition of nitrogen (N) fertilizer in salt – affected and less fertile soils. Malik *et al.* (1980) demonstrated a high activity of N, which indicates strong associative symbiotic relationship of N<sub>2</sub> – fixing bacterium (*Bacillus gram negative*) in the rhizosphere of *L. fusca*. Moreover, the N fixation through the growth of blue – green algae and *Azolla* under flooded conditions may partly contribute to the N supply and economy of the species. It is also observed that *L. fusca* contributes more stable organic matter fraction due to its slow decomposition as compared to succulent plant species like *Sesbania aculeata*. Kumar *et al.* (1980) reported an abrupt increase in the yield of *L. fusca* from 24 – 26 tonnes per ha per year,

without N application to 41 – 46 tonnes per ha per year when only 40 kg N per ha was applied in a sodic soil. Abdullah (1985) showed a definite ameliorative effect of phosphorus (P) on the growth of *L. fusca* under saline environment. The application of P at the rate of 50 kg per ha gave significantly higher fresh and dry matter yield at  $EC_e$  10  $dS\ m^{-1}$  than all other treatments, which was followed by 75 and 25 kg P per ha at  $EC_e$  20  $dS\ m^{-1}$ . The synergistic P ´ salinity effect was obvious at the highest P level of 75 kg per ha. Thus, the species responded favourably to P application at all salinity levels studied (*i.e.*  $EC_e$  3.5 to 30  $dS\ m^{-1}$ ), indicating higher P requirements. In general, the species is capable of accumulate trace elements (Zn, Cu, Fe, Mn) in sufficient amounts to meet the dietary requirements of the livestock under saline soil conditions (Abdullah *et al.* 1990).

#### 4. ROLE IN SOIL RECLAMATION

*Leptochloa fusca* behaved as a typical Crypno – eu – halophyte having both accumulating and excreting properties (Abdullah 1986; Abdullah *et al.* 1990). The efficient salt excretion from the shoot makes it a useful plant to deplete excessive salt from the root – zone and to provide a better root – environment for the growth of other plants. The extensive and fibrous roots of grass can open soil, increase air exchange, organic matter and hydraulic conductivity, decrease rhizosphere pH, stimulate biological activity, dissolve native  $CaCO_3$ , enhance leaching of salts, lower the water table of waterlogged soils, release plant nutrients and the shoot foliage can increase organic matter, humus and soil mulching, decrease surface evaporation and progressively improve soil physical propertiese (HAQ and KHAN 1971; JOSHI *et al.* 1981; ABDULLAH *et al.* 1986; MALIK *et al.* 1986 and AKHTAR *et al.* 1988).

#### 5. USE OF GROUND SALINE WATER FOR IRRIGATION

Although the groundwater is saline, it can be used for saline agriculture to grow salt tolerant trees, vegetables, crops and fodder grasses in non – saline – non – sodic coarse textured soils with minimum

adverse effects. This is due to rapid leaching of salts beyond the root zone and flushing of salts from root zone by rains (ABDULLAH *et al.* 1990). Furthermore, dense saline – sodic soils can also be used for growing such palatable grasses, which are very salt tolerant and capable of surviving in soils with poor properties. The sandy and loamy soil that is about 1 million ha can be brought under agriculture by using underground saline water and harvested rainwater. Experiments showed that under certain conditions plant could not only survive but also even vast areas of land could be irrigated with water of such high concentration. The soil is either sandy gravel or dune sand. Moderately saline irrigation water stimulates vegetation, assists the benevolent bacteria of the soil and improves yield and quality. Furthermore, use of brackish water reduces soil evaporation, transpiration of plants and increases resistance to drought (ABDULLAH *et al.* 1990). The solution of the adverse effects is suggested:

- Identification and selection of species and varieties tolerant of high salinity,
- The use of brackish water of such a degree of salinity only as is compatible with help of such species and with the nature of the soil,
- The selection of irrigation with such water in areas in which soils permeable, well drained and rich in calcium and the hydrates of iron and aluminium.

The Pakistan Council of Research in Water Resources (PCRWR) has planted *Eucalyptus*, *Acacia*, *Parkinsona*, *Zizyphus* (Beri), *Tamarix*, *Prosopis*, *Asparagus*, Date palm, Pomegranate, Jojoba and Iple Iple (*Leucaena leucocephala*). The saline water of concentration TSS (total soluble salts) is 2800 ppm (part per million) and SAR (sodium adsorption ratio) is 14 and being used for irrigation along with harvested rainwater to flush salts at certain intervals (Abdullah *et al.* 1990). The calcium sulphate fertilizer is also used to neutral the adverse effects of sodium salts. The growth of some of them is given in TABLES 2, 3 and 4.

TABLE 2: Biomass of fodder grasses per hectare grown at Dingarh, Cholistan Desert using saline water

Fodder grass	Biomass Fresh (kg)	Biomass Dry (kg)	Carrying capacity per year			
			Camel	Goat	Sheep	Cattle
<i>Cenchrus ciliaris</i>	16811	15012	2	14	16	3
<i>Panicum antidotale</i>	22191	12407	1	11	14	3
<i>Lasirus indicus</i>	25217	18247	2	17	20	4
<i>Napier Bajra</i>	43710	38780	4	35	42	9
<i>Leptochloa fusca</i>	13449	11445	1	10	13	3

Forage requirement (Dry matter per day):

1 sheep = 2.5 K

1 goat = 3.0 K

1 camel = 25.0 K

1 cattle = 12.5 K

Source: Based on field survey, June 2000.

TABLE 3: Biomass potential of some salt tolerant forages

Species	Green matter (kg/plant)	Dry matter (kg/plant)	Green matter yield (tonnes/hectare)	Dry matter yield (tonnes/hectare)	Plants/hectare
<i>Atriplex amnicola</i> 949	4.31	1.99	2.7	1.24	625
<i>Atriplex amnicola</i> 971	5.37	2.39	3.4	1.49	625
<i>Atriplex amnicola</i> 573	6.73	3.43	4.2	2.14	625
<i>Atriplex amnicola</i> □ <i>Atriplex nummularia</i>	5.13	2.15	3.2	1.34	625
<i>Atriplex buburyana</i> 1205 (Carnarvan)	3.11	1.6	7.8	4.0	2500
<i>Atriplex buburyana</i> 1200 (Leonora)	2.0	1.2	5.0	3.0	2500
<i>Atriplex cinerea</i> 524	5.0	2.35	3.1	1.46	625
<i>Atriplex lentiformis</i>	5.45	3.19	3.4	2.0	625
<i>Maireana aphylla</i> 1062	2.53	1.2	6.3	3.0	2500

Source: PADMU. – Pakistan Desertification Monitoring Unit (1986).

TABLE 4: Survival percentage and canopy cover of some plants grown in Cholistan Desert

Name of Tree / Shrub / Bush	Age (months)	Survival Percentage	Height (Cm)			Canopy Cover (Cm)		
			Min.	Mean	Max.	Min.	Mean	Max.
<i>Eucalyptus (Camddulensis)</i>	24	76	90	156	223	66	113	161
<i>Tamarix</i>	24	48	59	106	154	52	112	173
<i>Acacia</i>	24	67	66	125	193	55	126	197
Beri ( <i>Zizyphus</i> )	24	43	55	118	181	38	82	126
Jojoba ( <i>Simmondsia chinensis</i> )	18	76	20	60	110	08	48	89
<i>Atriplex halimus</i> (Local)	11	65	--	77	---	--	45	---
<i>Atriplex amnicola</i> 573	11	40	--	48	---	--	08	---
<i>Atriplex amnicola</i> 197	11	80	--	74	---	--	15	---
<i>Atriplex amnicola</i> 223	11	25	--	47	---	--	09	---
<i>Atriplex bunburyana</i> 1041	11	60	--	51	---	--	07	---
<i>Atriplex bunburyana</i> 1036	11	70	--	66	---	--	13	---
<i>Atriplex cincerea</i>	11	40	--	28	---	--	02	---
<i>Atriplex lintiformis</i>	11	60	--	76	---	--	26	---
<i>Atriplex commercial</i>	11	25	--	44	---	--	13	---
<i>Maireana aphylla</i>	11	85	--	35	---	--	30	---

Source: PADMU. – Pakistan Desertification Monitoring Unit (1986).





FIGURE 2: Wild oats grown by highly saline irrigation at PCRWR research station at Cholistan desert (Pakistan).  
Ahmad, Farooq 2008.



FIGURE 3: Frost trees grown by highly saline irrigation at PCRWR research station at Cholistan desert (Pakistan).  
Ahmad, Farooq 2008.



FIGURE 4: Innovative approach for mustard crop cultivation with saline irrigation on sandy desert at Cholistan (Pakistan). Ahmad, Farooq 2008.



FIGURE 5: Innovative approach for development of grassland with saline irrigation on sandy desert at Cholistan (Pakistan). Ahmad, Farooq 2008.

The germination capacity of different varieties of tomato, ladyfinger (*bhindi*), spinach (*palak*), cowpea and zucchini (*tori*) at different levels  $EC_e$  3 to 18 mmho/cm was studied in sand culture (PAMDU 1986). The germination was delayed and decreased with increased salinity. Significant

vegetables were found to fall in the order of salt tolerance: Spinach > Zucchini > Cowpea > Tomato > Ladyfinger (Abdullah *et al.* 1990). The list of some salt tolerant grasses and forages cultivated in the Cholistan Desert by using saline water is given in TABLE 5.

TABLE 5: Salt tolerant plants cultivated in Cholistan Desert by using saline water

Forage Crop/Grass/Bush	Tolerance (EC <sub>e</sub> × 10 <sup>6</sup> ) 4000 – 18000
Alkali grass ( <i>Puccinellia airoides</i> )	High
Bermuda grass ( <i>Cynodon dactylon</i> )	High
Kallar grass ( <i>Leptochloa fusca</i> )	High
Salt grass ( <i>Distichlis stricta</i> )	High
Desert wheat grass ( <i>Agropyron cristatum</i> )	High
Barley ( <i>Hordium vulgare</i> )	High
Rape ( <i>Brassica napus</i> )	Medium
Clover ( <i>Melilotus</i> )	Medium
Alfalfa ( <i>California common</i> )	Medium
Oats ( <i>Hay</i> )	Medium
<i>Atriplex spp.</i>	High

Source: PADMU. – Pakistan Desertification Monitoring Unit (1986).

## 6. CONCLUSION

The growth factors such as easy propagation, high – spreading rate, colonizing ability, vigorous growth, yield, palatability, nutritional value, long – term survival and high adaptability to environmental stress make *L. fusca* an excellent and versatile species that can be cultivated by using brackish water and salt – affected land of Cholistan for economic exploitation. The species has great promise for the economic utilization of sodic, high pH, waterlogged and saline soils. Similarly, high saline – sodic water can be used for successful cultivation of *L. fusca*.

## ACKNOWLEDGEMENTS

The author wishes to thank Dr. Mohammad Arshad, Cholistan Institute of Desert Studies, Islamia University Bahawalpur, Pakistan for providing technical assistance during the field study of Cholistan desert and valuable comments on a draft – version of this paper.

## REFERENCES

ABDULLAH, M. *Salt tolerance studies on Leptochloa Fusca*, Ph.D. 1985. Dissertation (unpublished), University of Agriculture, Faisalabad, Pakistan, 1985. p. 286.

ABDULLAH, M., AKRAM, M., KHAN, A.D.,

QURESHI, R.H. Internal water resources management by plants under various root environment stresses with special reference to kallar grass *Leptochloa Fusca*, In: NATIONAL SEMINAR ON WATER RESOURCES DEVELOPMENT AND ITS MANAGEMENT IN ARID AREAS, 6 – 8 October 1990, Quetta, Pakistan. *Anais...* Quetta, Pakistan, 1990.

ABDULLAH, M., QURESHI, R.H., AHMAD, N. Responses of *Leptochloa Fusca* to various types of substrate salinities. In: PAK. – US. BIOSALINE RESEARCH WORKSHOP, 22 – 26 September 1985, Karachi, Pakistan, 1986. *Anais...* Karachi, Pakistan, 1986. p.115 – 123.

AHMAD, F. *Socio – economic dimensions and ecological destruction in Cholistan*, Ph.D. 2002. Dissertation (unpublished), Department of Geography, University of Karachi, Pakistan, 2002.

AHMAD, F. Agro – pastoral systems in Cholistan, *Pakistan Geographical Review*, Vol.60 (2), 2005. p. 65 – 69.

AHMAD, F. Geoinformatics application to investigate agricultural potential in Cholistan desert, *Journal of Food, Agriculture & Environment*, WFL Publisher, Finland, Vol.5(2), 2007. p. 310 – 314.

AHMAD, F. Runoff farming in reducing rural poverty in Cholistan desert. *Soc. nat. (Online)*,



- Uberlândia, v. 20, n. 1, jun. 2008 . Disponível em <[http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S1982-45132008000100012&lng=pt&nrm=iso](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1982-45132008000100012&lng=pt&nrm=iso)>. Acesso em 18 fev. 2010. DOI: 10.1590/S1982-45132008000100012.
- AHMAD, F.; GULZAR, F.; SHIRAZI, S. A.; FAROOQ, S.; ALI, Z. *Leptochloa Fusca*: A high yielding grass for utilization of salt – affected soil and water resources in Cholistan, In: FOURTH INTERNATIONAL CONFERENCE ON LAND DEGRADATION, 12 – 17 September 2004, Murcia, Spain. *Anais...* Murcia, Spain, 2004. p.434.
- AKHTAR, J., R.A. WAHEED, K.A. MALIK, M.I. Haq. Effect of kallar grass growth on water transmission characteristics of salt – affected lands, In: FIRST NATIONAL CONGRESS ON SOIL SCIENCES, 6 – 8 October 1985, Lahore, Pakistan. *Anais...* Lahore, Pakistan, 1985. p.393 – 400.
- ARSHAD, M., ALTAF – UR – REHMAN RAO, AKBAR, G. Cholistan desert in a state of flux, UNEP: *Desertification Control Bulletin*, 26, 1995. p. 55 – 58.
- FAO/ADB. *Cholistan area development project*, Report No. 59/53 ADB – PAK 58 (Final version), Food and Agriculture Organization (FAO), Rome, Italy. 1993.
- HAQ, M., KHAN, M.F.A. Reclamation of saline and alkaline soil by growing kallar grass, *The Nucleus*, Vol.8 (4), 1971. p. 139 – 144.
- JOSHI, Y.C., QADIR, A., SHARMA, S.K. *Sequential changes in germination, root and shoot growth and anatomical characters of halophytes and glycophytes*, Annual report for 1980, Central Soil Salinity Research Institute, Karnal, India, 1981. p.94 – 97.
- JOWKAR, F., KHAN, M. AJMAL. *Socio – economic dimensions of resource management in Cholistan*, Institute for Development Anthropology (IDA), Binghamton, New York. 1996.
- KHAN, A.D., AKRAM, M., ABDULLAH, M. *Rainwater harvesting in Cholistan desert*, In: NATIONAL SEMINAR ON WATER RESOURCES DEVELOPMENT AND ITS MANAGEMENT IN THE ARID AREAS, 6 – 8 October 1990, Quetta, Pakistan. *Anais...* Quetta, Pakistan, 1990. p. 59 – 80.
- KUMAR, A.D., ABROL, I.P. DARGAN, K.S. *Karnal grass grows well in sodic soils*, Indian Farming, Vol.30 (3), 1980. p. 14 – 15.
- MALIK, K.A.; ZAFAR, Y.; HUSSAIN, A. (1980). *Nitrogenase activity in the rhizosphere of kallar grass Diplachne Fusca*, Biologia, Vol.26 (1 – 2), 1980. p. 107 – 112.
- MALIK, K.A., ASLAM, Z., NAQVI, M. *Kallar grass – a plant for saline lands*, Nuclear Institute for Agriculture and Biology (NIAB), Faisalabad, Pakistan, 1986. p.93.
- PAMDU. *Desertification process in Cholistan desert*, PCRWR Publication No. PADMU 7/86, 1986. p.155.
- QURESHI, R.H., ASLAM, M., ABDULLAH, M., PITMAN, M.G. *Diplachne Fusca*: An Australian salt tolerant grass used in Pakistan, *Journal of Australian Institute for Agricultural Science*, Vol.48 (4), 1982. p. 195 – 199.
- SANDHU, G.R. *Sustainable agriculture: A Pakistan National Conservation Strategy Sector Paper No. 2*, Environment and Urban Affairs Division, Government of Pakistan and IUCN – The World Conservation Union, Pakistan. 1993.
- SANDHU, G.R., ASLAM, Z., SATTAR, A., QURESHI, R.H., N. AHMAD, R.G. Wyn Jones. *The effect of salinity on the yield and composition of Diplachne Fusca (kallar grass) plant*, Cell Environ, Vol.4, 1981. p. 171 – 181.
- ZAFAR, Y.; MALIK, K.A. *Photosynthetic system of Leptochloa Fusca (L.) Kunth*, Pakistan Journal of Botany, Vol.16 (2), 1984. p. 109 – 116.