

SHORT COMMUNICATION

## Vultures and others scavenger vertebrates associated with man-sized pig carcasses: a perspective in Forensic Taphonomy

Caroline Demo<sup>1</sup>, Edison Rogério Cansi<sup>1</sup>, Cecília Kosmann<sup>1</sup> & José Roberto Pujol-Luz<sup>1,2</sup>

<sup>1</sup> Núcleo de Entomologia Forense, Departamento de Zoologia, Universidade de Brasília. 70910-900 Brasília, DF, Brazil.

<sup>2</sup> Corresponding author. E-mail: jrpujol@unb.br

**ABSTRACT.** The activity of vertebrates that feed on corpses can modify the chronology of the decomposition process and interfere with postmortem interval estimates. Further, by destroying the soft parts of the cadaver, scattering, burying or causing the disappearance of bones, it can entirely change the crime scene. In this study, we simulated a clandestine cemetery in an area of Cerrado located inside a farm in Brasília, Distrito Federal. Three domestic pigs of the size of a human of about 60 kg were placed on the ground in different periods of 2010 and 2011. We recorded four species of birds and one of mammal eating the carcasses: 1) Cathartidae: *Coragyps atratus* (Bechstein, 1973), *Cathartes aura* (Linnaeus, 1758), *Sarcoramphus papa* (Linnaeus, 1758); 2) Falconidae: *Caracara plancus* (Miller, 1777); and 3) Felidae: *Leopardus pardalis* (Lund, 1840). The behavior of these animals interfered in the decomposition process and resulted in the dispersion and loss of bony parts.

**KEY WORDS.** Cerrado; chronothanatognosis; clandestine cemeteries; decomposition; necrophagy.

Forensic Taphonomy is applied in several situations where the causes of death, or evidence, cannot be explained through routine methods (e.g., mass accidents, summary and mass executions, clandestine cemeteries) (UN 1991). The activities of scavengers animals, specially vultures, can modify the crime scene and generate several artifacts (MORTON & LORD 2006, CARTER *et al.* 2007, URURAHY-RODRIGUES *et al.* 2008, REEVES 2009), as the concealment of the body and/or its dismemberment (HAGLUND & SORG 1997, SPRADLEY *et al.* 2012). Either opportunistically or by lack of food, some predators such as big cats (RIPPLEY *et al.* 2012), rodents, dogs, and coyotes (HAGLUND 1997a,b) may occasionally feed on the carcasses, dismantling and moving it. The activity of these animals can modify the rate of decomposition and the post-mortem events, as well as the estimative of the post-mortem interval (PMI) (SPRADLEY *et al.* 2012).

During 2010 and 2011, three man-sized pigs (60 kg) were placed on the ground of a savannah-like formation (Cerrado) in a farm near to the urban area of Brasília, Distrito Federal, in order to simulate a clandestine cemetery. With the aim to record the vertebrates that fed on the carcasses, we installed a camera trap near them and left it there for 24 hours/day, from the moment of death until skeletonization.

The camera took a total of 831 photos. Five species of vertebrates, four birds and one mammal (Figs 1-6), were recorded: 1) Cathartidae: *Coragyps atratus* (Bechstein, 1973), *Cathartes aura* (Linnaeus, 1758), *Sarcoramphus papa* (Linnaeus, 1758); 2)

Falconidae: *Caracara plancus* (Miller, 1777); and 3) Felidae: *Leopardus pardalis* (Lund, 1840). The presence of the scavenger vertebrates was associated with the five stages of decomposition: initial, bloated, decay, postdecay, and dry (Table I).

Vultures, one of the most versatile scavenger birds, were the only vertebrates observed during the experiments that are known to have a diet composed almost entirely of carcasses (RUXTON & HOUSTON 2004). In Brazil, they are represented by six species: *C. atratus*, *C. aura*, *Cathartes burrovianus* (Cassin, 1845), *Cathartes melambrotus* (Wetmore, 1964), and *S. papa*. These birds are usually the first vertebrate scavenger to access the body (DEVVAULT *et al.* 2004, KJORLIEN *et al.* 2009, O'BRIEN *et al.* 2010), and can eat meat in advanced stage of putrefaction. The most remarkable interaction we observed was the lack of competition for food between *C. plancus* and *C. atratus*. The southern

Table I. Species of vertebrates visiting the pigs carcasses at the different stages of decomposition.

Species	Decomposition Stages				
	Initial	Bloated	Decay	Postdecay	Dry
<i>Coragyps atratus</i> (Bechstein, 1793)	■	■	■	■	■
<i>Cathartes aura</i> (Linnaeus, 1758)	■	■	■	■	■
<i>Sarcoramphus papa</i> (Linnaeus, 1758)	■	■	■	■	■
<i>Caracara plancus</i> (Miller, 1777)	■	■	■	■	■
<i>Leopardus pardalis</i> (Lund, 1840)	■	■	■	■	■



Figures 1-6. Scavenger vertebrates photographed by the camera trap using pigs carcasses as a food resource: (1) *Sarcoramphus papa*; (2) *Cathartes aura*; (3) *Caracara plancus*; (4) *Coragyps atratus*; (5) *Caracara plancus* and *Coragyps atratus* in a social behavior; (6) *Leopardus pardalis*.


crested caracara (*C. plancus*) is not limited to consuming carcasses, it can also eat the remains around it, beetle larvae and other insects that move away from the body. As a result of the intense activity of *C. atratus* on the carcasses, large bones, for instance the scapula and femur, were taken about six meters away from the site of decomposition. The desarticulation and transport of flesh and skeletonized remains is a well-known phenomenon attributed to scavenger birds, especially vultures (REEVES 2009, SPRADLEY *et al.* 2012).

Despite the fact that big cats are not commonly found feeding on carcasses, RIPLEY *et al.* (2012) reported bob cats *Lynx rufus* (Schreber, 1777) feeding on human corpses. The ocelot

(*L. pardalis*) has a solitary and terrestrial habits, predominantly nocturnal, and feeds on small vertebrates (DiBITETTI *et al.* 2006). Nevertheless, in our experiment, when *L. pardalis* fed on a carcass, it displaced it from its original position (Figs 7 and 8). Mechanical movements, spontaneous or deliberate, become important events for the interpretation of the crime scenes. Allied to this, moving the carcasses facilitates the dispersion of the body parts, as observed by URURAHY-RODRIGUES *et al.* (2008).

Studies involving the reconstruction of crime scenes are important and must consider the activity of scavengers in order to separate the peri-mortem and post-mortem events in the investigation (UBELAKER 1997).





Figures 7-8. *Leopardus pardalis* accessing the carcass and changing the angle of it by 90°: (7) August 31, 2011 at night; (8) September 1, 2011 at morning, where is possible to observe the movement of the carcass.

### ACKNOWLEDGEMENTS

The authors thank Miguel A. Marini (UnB) for the identification of the birds; José M. da S. Diogo, head of Fazenda Água Limpa (UnB), for the logistic help with the experiments; Hélio R. da Silva (UFRRJ) for reviewing the manuscript. We are also grateful to Hélio Spindula for his help in the field work and Karine B.B. Cordeiro for the images edition. To FAPDF and CNPq for the grants.

### LITERATURE CITED

- CARTER, D.O.; D. YELLOWLEES & M. TIBBETT. 2007. Cadaver decomposition in terrestrial ecosystems. *Naturwissenschaften* **94**: 12-24.
- DEVAULT, T.; I. BRISBIN & O. RHODES JR. 2004. Factors influencing the acquisition of rodent carrion by vertebrate scavengers and decomposers. *Canadian Journal of Zoology* **89**: 502-509.
- DI BITETTI, M.S.; A. PAVIOLO & C. DE ANGELO. 2006. Density, habitat use and activity patterns of ocelots (*Leopardus pardalis*) in the Atlantic Forest of Misiones, Argentina. *Journal of Zoology* **270**: 153-163.
- HAGLUND, W.D. 1997a. Rodents and Human Remains, p. 405-413. *In*: W.D. HAGLUND & M.H. SORG (Eds). **Forensic Taphonomy: The Post Mortem Fate of Human Remains**. Boca Raton, CRC Press, XXVI+636p.
- HAGLUND, W.D. 1997b. Dogs and Coyotes: Postmortem involvement with Human Remains, p. 367-379. *In*: W.D. HAGLUND & M.H. SORG (Eds). **Forensic Taphonomy: The Post Mortem Fate of Human Remains**. Boca Raton, CRC Press, XXVI+636p.
- HAGLUND, W.D. & M.H. SORG. 1997. Introduction to Forensic Taphonomy, p. 1-26. *In*: W.D. HAGLUND & M.H. SORG (Eds). **Forensic Taphonomy: The Post Mortem Fate of Human Remains**. Boca Raton, CRC Press, XXVI+636p.
- KJORLIEN, Y.P.; O.B. BEATTIE & A.E. PETERSON. 2009. Scavenging activity can produce predictable patterns in surface skeletal remains scattering: Observations and comments from two experiments. *Forensic Science International* **188**: 103-106.
- MORTON, R.J. & W.D. LORD. 2006. Taphonomy of child-sized remains: a study of scattering and scavenging in Virginia, USA. *Journal of Forensic Sciences* **51**: 475-479.
- O'BRIEN, R.C.; S.L. FORBES; J. MEYER & I. DADOUR. 2010. Forensically significant scavenging guilds in the southwest of Western Australia. *Forensic Science International* **198**: 85-91.
- REEVES, M.A. 2009. Taphonomic effects of vulture scavenging. *Journal of Forensic Sciences* **54**: 523-528.
- RIPPLEY, A.; N.C. LARISON; K.E. MOSS; J.D. KELLY & J.A. BYTHERMAY. 2012. Scavenging Behavior of *Lynx rufus* on Human Remains During the winter Months of Southeast Texas. *Journal of Forensic Sciences* **57**: 699-705.
- RUXTON, G.D. & D.C. HOUSTON. 2004. Obligate scavengers must be soaring fliers. *Journal of Theoretical Biology* **228**: 431-436.
- SPRADLEY, M.K.; M.D. HAMILTON & A. GIORDANO. 2012. Spatial patterning of vulture scavenged human remains. *Forensic Science International* **219**: 57-63.
- UBELAKER, D.H. 1997. Taphonomic Applications in Forensic Anthropology, p. 77-90. *In*: W.D. HAGLUND & M.H. SORG (Eds). **Forensic Taphonomy: The Post Mortem Fate of Human Remains**. Boca Raton, CRC Press, XXVI+636p.
- UN. 1991. **Manual sobre la prevención e investigación eficaz de las ejecuciones extralegales, arbitrarias o sumarias**. New York, United Nations, OCHA, 72p
- URURAHY-RODRIGUES, A.; J.A. RAFAEL; R.F. WANDERLEY; H. MARQUES & J.R. PUJOL-LUZ. 2008. *Coprophanæus lancifer* (Linnaeus, 1767) (Coleoptera, Scarabaeidae) activity moves a man-size pig carcass: relevant data for forensic taphonomy. *Forensic Science International* **182**: 19-22.

Submitted: 18.X.2012; Accepted: 13.IV.2013.

Editorial responsibility: Kleber Del Claro