

RESEARCH ARTICLE

## Assessing the population density of the spotted paca, *Cuniculus paca*, (Rodentia: Cuniculidae) on an Atlantic Forest island, southeastern Brazil

Átilla C. Ferreguetti<sup>1</sup>, Bruno C. Pereira<sup>2</sup>, Helena G. Bergallo<sup>1</sup>

<sup>1</sup>Departamento de Ecologia, Universidade do Estado do Rio de Janeiro. Avenida São Francisco Xavier 524, Maracanã, 20550-013 Rio de Janeiro, RJ, Brazil.

<sup>2</sup>Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis. Avenida Prefeito Mello Reis 1500, Aeroporto, 36033-560 Juiz de Fora, MG, Brazil.

Corresponding author: Átilla Colombo Ferreguetti ([atilla.ferreguetti@gmail.com](mailto:atilla.ferreguetti@gmail.com))

<http://zoobank.org/E92572BC-C948-44A8-B384-6773FB58CA7A>

**ABSTRACT.** The spotted paca, *Cuniculus paca* (Linnaeus, 1766), is a Neotropical, opportunistic, frugivorous caviomorph rodent, that inhabits primarily broadleaf forests. We aimed to provide the first estimates of density of *C. paca* for the Ilha Grande, an island located in the Atlantic Rain Forest biome of Brazil. Density and population size were estimated using the total number of individuals observed along each trail through the program DISTANCE 7. Our estimates of density and population size reinforces the importance of the Ilha Grande as an important reservoir of the species. Therefore, the results presented herein can be a starting point to support future action plans for the species, making predictions regarding the ecosystem and management and conservation of the spotted paca. Furthermore, the results can be used as a surrogate for other regions in which the species occurs.

**KEY WORDS.** Abundance, Distance sampling, Ilha Grande State Park, Line-transect.

### INTRODUCTION

Estimates of density and population size are fundamental to any conservation effort of endangered species. These estimates allow to evaluate a great range of impacts such as loss of habitat, identifying priority areas for conservation, evaluating the viability of isolated populations, determine the species conservation status, and has utility in several ecological studies in addition to serving as an important tool for decision makers (e.g., Jathanna et al. 2003, Tomas et al. 2004, Cardillo et al. 2006, Cunha and Loyola 2011, Buckland et al. 2016). However, such information is rare for most taxa, including endangered species, especially in the Neotropics (Cardillo et al. 2006, Cunha and Loyola 2011).

The spotted paca, *Cuniculus paca* (Linnaeus, 1766), is a Neotropical, opportunistic, frugivorous caviomorph rodent, that inhabits primarily broadleaf forests (Collett 1981, Pérez 1992, Aquino et al. 2009, Goulart et al. 2009). The species presents nocturnal habits, being able to eventually present crepuscular and dawn activity (Gómez et al. 2005, Michalski and Norris 2011). The spotted paca is listed as Least Concern both in the global and Brazilian red lists in view of its wide distribution, presumed large population, occurrence in a number of protected areas (Ministério do Meio Ambiente 2014, Emmons 2016). However,

local extinctions have occurred in the southeast of its range due to habitat fragmentation and poaching (Emmons 2016). Despite its conservation status, the spotted paca is of conservation and management interest throughout their geographic range, because it is a popular game species for people, an important prey of large carnivores, and an important seed disperser (Dubost and Henry 2006, Aquino et al. 2009, Foster et al. 2010). Currently, estimates of density and abundance of the spotted paca are limited to few studies that used the most frequently methods employed, such as sightings; indirect records, and camera traps (e.g. Collett 1981, Beck-King et al. 1999, Asprilla-Perea and Lopez-Perea 2011, Santos-Moreno and Perez-Irriego 2013). Spotted pacas may be locally abundant but estimated population parameters can be laborious and results are likely to be unsatisfying (Beck-King et al. 1999). Few studies employed nocturnal surveys (e.g., Glanz 1990, Malcolm 1990, Beck-King et al. 1999, Oversluijs-Vásquez 2003, Aquino et al. 2009), and a great part of knowledge about the species natural history comes from studies that focus on the species as a feline prey.

In this context, we aimed to provide the first estimates of density of *C. paca* for the Ilha Grande, an island located in the Atlantic Rain Forest biome of Brazil. We also discuss the importance of performing nocturnal samplings for estimating density of species such as *C. paca*.

## MATERIAL AND METHODS

Our study was carried out in Ilha Grande State Park (PEIG), an island (Ilha Grande) located in the Southwestern coast of Rio de Janeiro State, Brazil (Fig. 1). PEIG is the second largest insular strictly protected park in Brazil managed by Brazilian authorities and covers some 120 km<sup>2</sup>, over half (62%) of the 193 km<sup>2</sup> area of Ilha Grande. The climate is tropical, hot and humid, without a dry season. Ilha Grande is the top of a submerged mountain and has two dominant types of topography, mountain and coastal plain. Mountain peaks occur in the center of the island. Almost half of the area (47%) is covered by dense, relatively pristine Atlantic rainforest. Secondary forest, in an advanced successional stage, is the second major habitat type (43%). The remaining areas comprise rocky outcrops with herbaceous vegetation (7%), salt marsh, mangroves and beaches (2%), and 1% human settlement (Alho et al. 2002). Although the area occupied by human settlement is small, it is concentrated around the northern coastline of Ilha Grande Bay and in settlements such as Abraão village.

During the period between December 2003 and May 2005, 128 transects were performed, totaling 401.3 km walked in 382 hours of effort. The transects were surveyed in five existing dirt trails in Ilha Grande to minimize the impact of opening new trails (Fig. 1, Table 1). These five trails covered all vegetation types found in the Ilha Grande and crossed several streams. Two of those trails are located in the northern part, connecting the largest settlement on the island, the Abraão Village, the beaches of Palmas and Feiticeira (respectively T01 and T02). The other three trails are on the south side, connecting the Dois Rios Village to Caxadaço and Parnaióca beaches, and the place known as Jararaca (respectively T03, T04 and T05). Transects ranged in length from 2.1 to 6.7 km. Transects were walked at times when the species are at increased activity: at dawn and twilight/night following sunrise and sunset hours along the year (Gómez et al. 2005, Michalski and Norris 2011). Thus, they were divided into two categories: dawn and twilight/nocturnal transects. Transects were walked at an average speed of 1.1 km/h ( $\pm 0.5$ ).

For each sighting, we recorded the perpendicular distance of the animal from transect (measured using measuring tape),




Figure 1. Ilha Grande in the state of Rio de Janeiro, Brazil and location of the transects. Black triangle representing the Abraão village.

Table 1. Characteristics of transects (length, altitude and region), number of times each transect was walked (N samples), total kilometers walked, and the encounter rates of *Cuniculus paca* per transect.

Transect	T01	T02	T03	T04	T05	Total
Coordinates	23°07'37.4"S 44°10'47.3"W	23°08'43.3"S 44°08'53.5"W	23°10'07.9"S 44°10'24.9"W	23°11'26.2"S 44°13'43.5"W	23°09'25.4"S 44°14'19.6"W	
Extension (km)	2.1	2.3	2.7	6.7	2.1	15.9
N samples	27	23	27	24	27	128
Total Km walked	56.7	52.9	72.9	160.8	56.7	401.3
<i>Cuniculus paca</i>	0	2.26	0.41	0.18	0.18	0.62

the length of transect walked to that point, date, and timing of the record. Individuals of spotted paca were detected using a head-flashlight. Density and population size were estimated using the total number of individuals observed along each trail through the program DISTANCE 7 (Buckland et al. 2001). DISTANCE uses the perpendicular distances to the observed animals to estimate effective strip width (ESW) in the study area and model the detection function that best suits the probability of detection of an animal at a given distance (Buckland et al. 2001, Thomas et al. 2010). The best detection model is selected by the Akaike Information Criterion (AIC; Akaike 1973).

## RESULTS

Throughout the surveys, *C. paca* was only recorded at night and we obtained 42 observations. Estimated density for *C. paca* was  $35 \pm 3.2$  individuals/km<sup>2</sup> and estimated population size for the entire Ilha Grande was  $6755 \pm 617$  individuals. The effective strip width (ESW) was  $3.40 \pm 0.58$  m with records obtained from 0 to 9 m from the line of the transect (Fig. 2). The model which best fitted our data was a Hazard rate with cosine adjustment. The coefficient of variation for both density and population size was 18.22%.

## DISCUSSION

Spotted paca density, estimated in the Ilha Grande through transects, is the first estimate for the area and showed a coefficient of variation below 20%, which is the maximum value recommended to obtain reliable estimates (Buckland




Figure 2. Plot of the detection function for spotted pacas based on the AIC selected Conventional Distance Sampling (CDS) model. Histogram represents the probability of detection for each distance interval. The curved line is the detection function, showing the probability that a spotted paca is observed as a function of distance from the transect line.

et al. 2001). This indicate precision in the estimates and can assist in the species' conservation as starting point for monitoring the species in the Ilha Grande. Density estimated for the species in the Ilha Grande is within the range observed in Neotropics forests, with a population density range of 25-70 adults per square km depending on the habitat (Patton 2015). In fact, estimated densities range from 3.5 individuals/km<sup>2</sup> in Peru (Emmons 1987) to 93 ind/km<sup>2</sup> in Puente Gloria, Colombia (Collett 1981). This intermediary density can be explained by the absence of larger predators such as *Panthera onca* (Linnaeus, 1758) and *Puma concolor* (Linnaeus, 1771) in Ilha Grande (Lessa et al. 2017) which allows to increase the spotted paca density in the island. In fact, prey species such as spotted pacas are thought to have lower densities in areas in which top predators are present (Terborgh and Winter 1980, Wright et al. 1994). However, despite the absence of large predators, the region is impacted by both poaching and invasive species such as *Canis lupus familiaris* Linnaeus, 1758 (Lessa et al. 2017), which could be reducing species density. Thus, highlighting the importance to monitor the population of the species in the long term that despite being listed as Least Concern by the global and Brazilian red lists have become locally extinct in some locations due to poaching (Emmons 2016).

Although some authors avoid collecting data in nocturnal transects (Duckworth 1998), this study has shown that they provide a good amount of data for the spotted paca that is lacking density estimations in the Atlantic Forest. Since most of the Neotropical mammals have predominantly nocturnal habits (Eisenberg and Redford 1999, Paglia et al. 2012), a higher sampling effort in this time range is required to obtain sufficient data to estimate the density of these species. It is worth mentioning that in the zero-distance perpendicular to the transect, the probability of detection is lower than in the first meter as showed in Fig. 2. This is due to the fact that for nocturnal animals, eye brightness is the most important detection method (Pereira et al. 2017). Therefore, if the animal is too close, the angle may not allow you to have the vision of the eye glow. This must be taken into account when performing the analysis to avoid bias in the density estimation as suggested by Buckland et al. (2016).

The estimate presented here is the first for the Atlantic Forest using distance sampling. Our estimates of density and population size reinforces the importance of the Ilha Grande as an important reservoir of the species. Therefore, the results presented herein can be a starting point to support future action plans for the species, making predictions regarding the ecosystem and management and conservation of the spotted paca. Furthermore, the results can be used as a surrogate for other regions in which the species occurs.

## ACKNOWLEDGEMENTS

We thank the Centro de Estudos Ambientais e Desenvolvimento Sustentável (CEADS) and the Universidade do Estado

do Rio de Janeiro (UERJ) for the logistical support during the execution of this research. We also thank the Instituto Biomas for financial support. BCP is also grateful to Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes) for a graduate scholarship. HGB thanks grants from FAPERJ (E26/201.267/2014), CNPq (307781/2014-3) and Prociência/UERJ.

## LITERATURE CITED

- Akaike H (1973) Maximum likelihood identification of gaussian autoregressive moving average models. *Biometrika* 60: 255–265. <https://doi.org/10.1093/biomet/60.2.255>
- Alho CJR, Schneider M, Vasconcellos LA (2002) Degree of threat to the biological diversity in the Ilha Grande State Park (RJ) and guidelines for conservation. *Brazilian Journal of Biology* 62(3): 375–385. <https://doi.org/10.1590/S1519-69842002000300001>
- Aquino R, Gil D, Pezo E (2009) Aspectos ecológicos y sostenibilidad de la caza del majás (*Cuniculus paca*) en la cuenca del río itaya, Amazonía peruana. *Revista Peruana de Biología* 16: 67–72.
- Asprilla-Perea J, López-Perea JJ, Viveros-Riveros JA, Jiménez-Ortega AM (2011) Relación entre abundancia relativa y el aprovechamiento de *Cuniculus paca* (guagua, tepezcuintle) en comunidades negras de la Cuenca Del Atrato, Colombia. *Mastozoología Neotropical* 18: 301–306.
- Beck-King H, Helversen OV, Beck-King R (1999) Home range, population density, and food resources of *Agouti paca* (Rodentia: Agoutidae) in Costa Rica: a study using alternative methods. *Biotropica* 31: 675–685. <https://doi.org/10.1111/j.1744-7429.1999.tb00417.x>
- Buckland ST, Anderson DR, Burnham KP, Laake JL, Borchers DL, Thomas L (2001) *Introduction to Distance Sampling: estimating abundance of biological population*. Oxford University Press, Oxford, 446 pp.
- Buckland ST, Rexstad E, Thomas L, Borchers DL (2016) Distance sampling surveys of population size: enabling better decision-making by wildlife managers. In: Buckland ST (Ed.) *UK success stories in industrial mathematics*. Springer International Publishing, 45–51. [https://doi.org/10.1007/978-3-319-25454-8\\_6](https://doi.org/10.1007/978-3-319-25454-8_6)
- Cardillo M, Mace GM, Gittleman JL, Purvis A (2006) Latent extinction risk and future battlegrounds of mammal conservation. *Proceedings of the National Academy of Sciences* 103: 4157–4161. <https://doi.org/10.1073/pnas.0510541103>
- Collett SF (1981) Population characteristics of *Agouti paca* (rodentia) in Colombia. *Publications of the Museum, Michigan State University, Biological Series* 5: 485–602.
- Cunha FMA, Loyola RD (2011) Spatial priorities for the conservation of threatened mammals in the Neotropics. *Bioikos* 25: 75–90.
- Dubost G, Henry O (2006) Comparison of diets of the Acouchy, Agouti and Paca, the three largest terrestrial rodents of French Guianan forests. *Journal of Tropical Ecology* 22: 641–651. <https://doi.org/10.1017/S0266467406003440>
- Duckworth JW (1998) The difficulty of estimating population densities of nocturnal forest mammals from transect counts of animals. *Journal of Zoology* 246: 443–486. <https://doi.org/10.1017/S0952836998281210>
- Eisenberg JF, Redford KH (1999) *Mammals of the Neotropics. The Central Neotropics: Ecuador, Peru, Bolivia, Brazil*. University of Chicago Press, Chicago, 609 pp.
- Emmons LH (1987) Comparative feeding ecology of felids in a Neotropical Rainforest. *Behavioral Ecology and Sociobiology* 20: 271–283. <https://doi.org/10.1007/BF00292180>
- Emmons L (2016) *Cuniculus paca*. The IUCN Red List of Threatened Species 2016: e.T699A22197347. <https://doi.org/10.2305/IUCN.UK.2016-2.RLTS.T699A22197347.en> [Accessed: 25/08/2018]
- Foster RJ, Harmsen BJ, Valdez B, Pomilla C, Doncaster CP (2010) Food habits of sympatric jaguars and pumas, across a gradient of human disturbance. *Journal of Zoology* 280: 309–318. <https://doi.org/10.1111/j.1469-7998.2009.00663.x>
- Glanz WE (1990) Neotropical mammal densities: how unusual is the community on Barro Colorado island, Panama? In: Gentry AH (Ed.) *Four Neotropical Rainforest*. Yale University Press, New Haven, 287–311.
- Gómez H, Wallace RB, Ayala G, Tejada R (2005) Dry season activity periods of some Amazonian mammals. *Studies on Neotropical Fauna and Environment* 40(2): 91–95. <https://doi.org/10.1080/01650520500129638>
- Goulart FVB, Cáceres NC, Graipel ME, Tortato MA, Ghizoni IR, Oliveira-Santos LGR (2009) Habitat selection by large mammals in a southern Brazilian Atlantic Forest. *Mammalian Biology* 74: 182–190. <https://doi.org/10.1016/j.mambio.2009.02.006>
- Jathanna D, Karanth KU, Johnsingh AJT (2003) estimation of large herbivore densities in the tropical forest of southern india using distance sampling. *Journal of Zoology* 261: 285–290. <https://doi.org/10.1017/S0952836903004278>
- Lessa ICM, Ferreguetti AC, Kajin M, Dickman CR, Bergallo HG (2017) You can't run but you can hide: the negative influence of human presence on mid-sized mammals on an Atlantic island. *Journal of Coastal Conservation* 21(6): 829–836. <https://doi.org/10.1007/s11852-017-0544-2>
- Malcolm JR (1990) Estimation of mammalian densities in a continuous forest north of Manaus. In: Gentry AH (Ed.) *Four Neotropical Rainforest*. Yale University Press, New Haven, 339–357.
- Michalski F, Norris D (2011) Activity pattern of *Cuniculus paca* (Rodentia: Cuniculidae) in relation to lunar illumination and other abiotic variables in the southern Brazilian Amazon. *Zoologia (Curitiba)* 28(6): 701–708. <https://doi.org/10.1590/S1984-46702011000600002>
- Ministério do Meio Ambiente (2014) *Lista Nacional Oficial das Espécies da Fauna Brasileira Ameaçadas de Extinção*. Portaria n. 444, de 17 de dezembro de 2014. *Diário Oficial da República Federativa do Brasil, Brasília, Seção 1*, 245, 121–126.
- Oversluijs-Vásquez MR (2003) Animales de caza en la zona reservada Allpahuayo-Mishana. *Folia Amazónica* 14: 7–16. <https://doi.org/10.24841/fa.v14i1.149>

- Paglia AP, Da Fonseca GA, Rylands AB, Herrmann G, Aguiar LM, Chiarello AG, Leite YLR, Costa LP, Siciliano S, Kierulff MCM, Mendes SL, Tavares VC, Mittermeier RA, Patton JL (2012) Lista Anotada dos Mamíferos do Brasil / Annotated Checklist of Brazilian Mammals. Conservation International, Arlington, Occasional Papers in Conservation Biology, no. 6, 2<sup>nd</sup> ed., 76 pp.
- Patton JL (2015) Family Cuniculidae G.S. Miller & Gidley, 1918. In: Patton JL, Pardiñas UFJ, D'Elía G (Eds) Mammals of South America. University of Chicago Press, Chicago, 726–733. <https://doi.org/10.7208/chicago/9780226169606.001.0001>
- Pereira BC, Ferregueti AC, Bergallo HG (2017) Factors affecting mammalian encounter rates in transect surveys: a case study in Ilha Grande State Park, State of Rio de Janeiro, Brazil. *Oecologia Australis* 21(4): 1–9. <https://doi.org/10.4257/oeco.2017.2104.06>
- Pérez EM (1992) *Agouti paca*. *Mammalian Species* 404: 1–7. <https://doi.org/10.2307/3504102>
- Santos-Moreno A, Pérez-Irineo G (2013) Abundancia de tepezcutintle (*Cuniculus paca*) y relación de su presencia con la de competidores y depredadores en una selva tropical. *Therya* 4: 89–98. <https://doi.org/10.12933/therya-13-97>
- Terborgh J, Winter B (1980) Some causes of extinction. In: Soulé ME, Wilcox BA (Eds) *Conservation Biology: an evolutionary-ecological perspective*. Sunderland, Sinauer Associates, 119–133.
- Thomas L, Buckland ST, Rexstad EA, Laake JL, Strindberg S, Hedley SL, Bishop JRB, Marques TA, Burnham KP (2010) Distance software: design and analysis of distance sampling surveys for estimating population size. *Journal of Applied Ecology* 47(1): 5–14. <https://doi.org/10.1111/j.1365-2664.2009.01737.x>
- Tomas WM, Rodrigues FHG, Fusco R (2004) Técnicas de levantamento e monitoração de populações de carnívoros. Embrapa, Corumbá, 34 pp.
- Wright SJ, Gompper ME, Deleon B (1994) Are large predators keystone species in Neotropical forests? The evidence from Barro Colorado island. *Oikos* 71: 279–294. <https://doi.org/10.2307/3546277>
- 
- Submitted: December 19, 2017  
Accepted: February 15, 2018  
Available online: November 22, 2018  
Editorial responsibility: Diego Astúa de Moraes
- 
- Author Contributions:** ACF, BCP and HGB contributed equally to this article.  
**Competing Interests:** The authors have declared that no competing interests exist.  
© 2018 Sociedade Brasileira de Zoologia. Published by Pensoft Publishers at <https://zoologia.pensoft.net>