

SHORT COMMUNICATION

Distribution extension of *Phimophis guerini* (Serpentes: Dipsadidae: Xenodontinae) in the Brazilian Amazon

Alfredo P. Santos-Jr¹, Danilo A. Almeida-Santos², Síría Ribeiro¹,
 Isla C.M. Carmargo¹, Ana Lúcia da Costa Prudente³

¹Laboratório de Ecologia e Comportamento Animal, Universidade Federal do Oeste do Pará. Rua Vera Paz, Salé, 68035-110 Santarém, PA, Brazil.

²Independent Researcher currently not affiliated. Rua Hortênciã 261, Jardim Santarém, 68030-310 Santarém, PA, Brazil.

³Coordenação de Zoologia, Laboratório de Herpetologia, Museu Paraense Emílio Goeldi. Avenida Perimetral 1901, Caixa Postal 399, 66077-830 Belém, PA, Brazil.

Corresponding author: Alfredo P. Santos-Jr (alphredojr@hotmail.com)

<http://zoobank.org/C8D91102-A2E8-466A-B1C0-519730B30BC9>

ABSTRACT. *Phimophis guerini* Duméril, Bibron & Duméril, 1854 is a Xenodontinae snake distributed in Argentina, Brazil and Paraguay. In Brazil, the species is broadly distributed, occurring mainly in open areas of the Cerrado, but also in the Amazon, Atlantic forest and Caatinga. We provide a new record for this species from the municipality of Santarém in the western portion of the state of Pará (Brazil). Five specimens were collected in a small area covered with Amazonian Savanna vegetation. We also provide the description of the morphological variation for the collected specimens. The new record extends the northern limit of the distribution by some 640 km (from Floresta Nacional de Carajás, Parauapebas municipality, eastern Pará). The record from Santarém provides a third locality for *P. guerini* within the Amazon biome and supports the hypothesis of a past ecological corridor linking the Cerrado and the open habitats within the Amazon.


KEY WORDS. Amazonian savanna, morphological data, Pará state, Pseudoboini.

Phimophis and 11 other genera compose the South American tribe Pseudoboini (Zaher et al. 2009, Grazziotin et al. 2012). Currently, the genus is represented by three known species: *Phimophis guerini* (Duméril, Bibron & Duméril, 1854), *Phimophis guianensis* (Troschel, 1848), and *Phimophis vittatus* (Boulenger, 1896). Within the genus, the species with the widest known distribution is *P. guerini* with confirmed records for Argentina, Paraguay and Brazil (Peters and Orejas-Miranda 1970, Tavares et al. 2012, Atkinson et al. 2017). In Brazil, *P. guerini* has a broad distribution, occurring mainly in open areas of the Cerrado, but also in the Mata Atlântica, Caatinga and Amazonian Savanna (Costa et al. 2010, Pereira-Filho et al. 2012, França et al. 2006, 2013, França et al. 2012, Marques et al. 2012, Maschio et al. 2012, Santos and Vaz-Silva 2012, Tavares et al. 2012).

The Amazonian savanna is composed of isolated patches of open vegetation within the Amazon forest. The floristic composition of the Amazonian savannas is less rich than the savannas in central Brazil. However, both formations present several species of herbaceous and woody plants in common (Magnusson et al. 2008). The areas are occasionally affected

by fire, but the vegetation is adapted to recover in a cycle that includes fire as an intrinsic and regular phenomenon (Faria et al. 2004). Compared to other Amazonian ecosystems (e.g., 'terra-firme' Forest), Amazonian savannas are poorly studied (Carvalho and Mustin 2017). Biogeographic studies have proposed past ecological corridors interconnecting these fragments with the central region of the South American's open diagonal, including the Caatinga, Cerrado and Chaco biomes (Werneck et al. 2012). Here we present a new geographic record and morphological data for five specimens of *P. guerini* collected in the central Amazonian Savanna.

The study was undertaken in a small area of Amazonian Savanna (Fig. 1) in the community of Tapari, municipality of Santarém, western portion of the state of Pará, Brazil (2°26.827'S; 54°53.577'W), where four specimens of *P. guerini* were collected in pitfall traps (see a photo of one of the specimens alive in Fig. 2). The study was carried out from October 2012 to September 2013, as part of a broader inventory project to catalog the species of snakes present in the Amazonian savanna of the western Pará. An additional specimen (Figs 3–5) was collected in May 2015



Figures 1–5. Habitat and specimens of *Phimophis guerini* collected in an amazon savanna area in Santarém, Pará, Brazil: (1) specimen collection environment; (2) living specimen MPEG 2638; (3–5) dorsal, lateral and ventral views, respectively, of the head of specimen UFOPA-H 0249. Scale bars: 5 mm.

on an unpaved road near one of our sampling sites (2°28.546'S; 54°53.827'W). The Amazonian Savanna in the studied area consists largely of herbaceous vegetation, varying in height and density, accompanied by shrub and tree strata, presenting abrupt edges between the open areas and the forests, with a small ecotone area where soil is predominantly sandy (Magnusson et al. 2008). All collected specimens were deposited in the collections of the Museu Paraense Emílio Goeldi (MPEG 26538, MPEG 26539, MPEG 26540) and of the Universidade Federal do Oeste do Pará (UFOPA-H 0249, UFOPA-H 1237).

The extent and nature of the morphological variation of *P. guerini* is not very well known. Therefore, we present here the morphometric and meristic data from the collected specimens (Table 1). Records of *P. guerini* from western Pará extends the previously known distribution of the species by approximately 640 km from the nearest locality (Floresta Nacional de Carajás, Parauapebas municipality, eastern Pará state; Maschio et al. 2012). This record from Santarém provides a third locality for *P. guerini* within the Amazonian biome (França et al. 2006, Maschio et al. 2012) (Fig. 6). All Amazonian records are restricted to open




Figure 6. Updated geographic distribution of *Phimophis guerini* in Brazil, showing the new record (star) in western Pará, Brazil. Previous records (circles) were taken from the following publications: Costa et al. (2010), Pereira-Filho et al. (2012) França et al. (2006, 2013), França et al. (2012), Marques et al. (2012), Maschio et al. (2012), Santos and Vaz-Silva (2012) Tavares et al. (2012).

Table 1. Morphometric and meristic data of specimens of *Phimophis guerini* collected in the community of Tapari, Santarém Pará, Brazil. The snout-vent length and tail length are given in millimeters. Bilateral variation is reported as "right/left".

Characters	Specimens				
	MPEG 26538	MPEG 26539	MPEG 26540	UFOPA-H 1237	UFOPA-H 0249
Sex	Male	Male	Female	Female	Female
Snout-vent length	465	540	560	587	669
Tail length	115	135	101	121	127
Supralabials	8/8	8/8	8/8	8/8	8/9
Infralabials	09/09	10/10	10/09	10/09	09/09
Oculars	1+2/1+2	1+2/1+2	1+2/1+2	1+2/2+2	1+2/1+2
Temporals	3+3/2+3	2+3/2+3	2+3/2+3	2+3/3+3	3+3/2+3
Dorsal scales	21+19+17	21+19+17	21+19+17	21+19+17	21+19+17
Ventrals	195	195	207	205	207
Subcaudals	73	73	63	67	61

habitats. The specimen UFOPA-H 0249 was collected at night while foraging and had the third-most part of its body inserted in a lizard hole in the sand, probably excavated by *Ameiva ameiva* (Linnaeus, 1758), which can indicate the saurophagous habit of the species (Sawaya et al. 2008).

Phimophis guerini is a relatively common species in the cerrado of Central Brazil, although the species seems to be rare in the Amazon domain. We present here the second record for this snake in the state of Pará, reinforcing the importance of inventorying biological diversity of the poorly known Amazonian Savanna. These areas can be considered particularly important for the conservation of the Amazonian fauna, since they can encompass communities composed of forest and savanna dwellers, and even present endemics species (Carvalho and Mustin 2017). The presence of *P. guerini* in Amazonian savannas supports the idea that past open corridors may have connected central Brazil to the extreme north of the Amazon during the Last Glacial Maximum (c. 21 ka) (Werneck et al. 2012). Such corridors were most likely temporary and the transitory connection among these regions is also supported by the disjunct distribution of other species of snake, including *Taeniophallus occipitalis* (Jan, 1863) and *Oxyrhopus rhombifer septentrionalis* Vellard, 1943 (Peters and Orejas-Miranda 1970, França et al. 2006, Santos-Jr et al. 2008).

ACKNOWLEDGMENTS

This research was supported by grant from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Universal 14/2011, process 478347/2011-1). We thank Maiume Silva da Silva and Joanderson Martins for helping during the field work, and Antônio Figueira for logistical support. Adrian Barnett reviewed the English of the first draft. We thank Felipe Grazziotin and an anonymous reviewer for comments and suggestions. DAAS was supported by scientific initiation grant from Universidade Federal do Oeste do Pará; and ICMC by master grant from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior.

LITERATURE CITED

- Atkinson K, Smith P, Sarvary J (2017) New and noteworthy snake species records (Colubridae and Dipsadidae) for the Reserva Natural Laguna Blanca, eastern Paraguay. *Check List* 13(1): 1–5. <https://doi.org/10.15560/13.1.2027>
- Carvalho WD, Mustin K (2017) The highly threatened and little known Amazonian savannas. *Nature Ecology and Evolution* 1 (100): 1–3. <https://doi.org/10.1038/s41559-017-0100>
- Costa GC, Nogueira C, Machado RB, Colli GR (2010) Sampling bias and the use of ecological niche modeling in conservation planning: a field evaluation in a biodiversity hotspot. *Biodiversity and Conservation* 19(3): 883–899. <https://doi.org/10.1007/s10531-009-9746-8>
- Faria AS (2004) The effects of fire on behaviour and relative abundance of three lizard species in an Amazonian savanna. *Journal of Tropical Ecology* 20: 591–594. <https://doi.org/10.1017/S0266467404001798>
- França FGR, Mesquita DO, Colli GR (2006). A checklist of snakes from Amazonian savannas in Brazil, housed in the Coleção Herpetológica da Universidade de Brasília, with new distribution records. *Occasional Papers of the Oklahoma Museum of Natural History* 17: 1–13.
- França FGR and Braz VS (2013) Diversity, activity patterns, and habitat use of the snake fauna of Chapada dos Veadeiros National Park in Central Brazil. *Biota Neotropica* 13(1): 74–85. <http://www.biotaneotropica.org.br/v13n1/en/abstract?article+bn01313012013>
- França RC, Germano CES, França FGR (2012) Composition of a snake assemblage inhabiting an urbanized area in the Atlantic Forest of Paraíba State, Northeast Brazil. *Biota Neotropica* 12(3): 1–13. <http://www.biotaneotropica.org.br/v12n3/en/abstract?inventory+bn00612032012>.
- Grazziotin FG, Zaher H, Murphy RW, Scrocchi G, Benavides MA, Zhang Y, Bonatto SL (2012) Molecular phylogeny of the New World Dipsadidae (Serpentes: Colubroidea): a reappraisal. *Cladistics* 28(5): 437–459. <https://doi.org/10.1111/j.1096-0031.2012.00393.x>
- Magnusson WE, Lima AP, Albernaz ALKM, Sanaiotti TM, Guillaumet J (2008) Composição florística e cobertura vegetal das savanas na região de Alter do Chão, Santarém – PA. *Revista Brasileira de Botânica* 31(1): 165–177. <https://doi.org/10.1590/S0100-84042008000100015>
- Marques R, Tinôco MS, Browne-Ribeiro HC, Fazolato CP (2012) *Phimophis guerini* (Duméril, Bibron and Duméril, 1854) (Squamata, Colubridae): Distribution extension in the northeast coast of the state of Bahia, Brazil. *Check List* 8 (5): 963–965.
- Maschio GF, Galatti U, Oliveira SN, Gordo M, Bitar YOC (2012) Répteis de Carajás. In: Martins FD, Castilho AF, Campos J, Hatano FM, Rolim SG. (Orgs) *Fauna da Floresta Nacional de Carajás: Estudos sobre Vertebrados Terrestres*. São Paulo, Nitro Imagens, 82–97.
- Pereira-Filho GA, Santana GG, Vieira WLS, Alves RRN, Montenegro PFGP, Freitas MA (2012) *Phimophis guerini* (Duméril, Bibron and Duméril, 1854) (Serpentes: Dipsadidae): Distribution extension in Paraíba, Brazil. *Check List* 8(5): 966–967. <https://doi.org/10.15560/8.5.966>
- Peters JA, Orejas-Miranda B (1970) Catalogue of the Neotropical Squamata: Part I. Snakes. *Bulletin of American Museum of Natural History* 297(1): 1–347.
- Santos DL, Vaz-Silva W (2012) Predation of *Phimophis guerini* and *Ameiva ameiva* by *Erythrolamprus aesculapii* (Snake: Colubridae). *Herpetology Notes* 5: 495–496.
- Santos-Jr AP, Di-Bernardo M, Lema T (2008) New species of the *Taeniophallus occipitalis* Group (Serpentes, Colubridae) from Eastern Amazonia, Brazil. *Journal of Herpetology* 42(3): 419–426.
- Sawaya RJ, Marques OAV, Martins M (2008) Composição e história natural das serpentes de Cerrado de Itirapina, São

- Paulo, sudeste do Brasil. *Biota Neotropica* 8(2): 127–149. <http://www.biotaneotropica.org.br/v8n2/en/abstract?inventory+bn01308022008>
- Tavares JR1, Melo CE, Campos VA, Oda FH, Strüssmann C (2012) Snakes from Canoa Quebrada Hydroelectric Power Plant, state of Mato Grosso, Brazil. *Herpetology Notes* 5: 543–546.
- Werneck FP, Nogueira C, Colli GR, Sites Jr JW, Costa GC (2012) Climatic stability in the Brazilian Cerrado: implications for biogeographical connections of South American savannas, species richness and conservation in a biodiversity hotspot. *Journal of Biogeography* 39: 1695–1706. <https://doi.org/10.1111/j.1365-2699.2012.02715.x>
- Zaher H, Grazziotin FG, Cadle JE, Murphy RW, Moura-Leite JC, Bonatto SL (2009) Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American Xenodontines: a revised classification and descriptions of new taxa. *Papéis Avulsos de Zoologia* 49(11): 115–153. <https://doi.org/10.1590/S0031-10492009001100001>
-
- Submitted: June 21, 2018
Accepted: April 7, 2019
Available online: July 5, 2019
Editorial responsibility: Felipe Grazziotin
-
- Author Contributions: APJS, SR and ALCP planned the project; APSJ, DAAS and SR conducted field works; APSJ, ICMC and ALCP analyzed the data; all authors contributed to write and review the paper.
- Competing Interests: The authors have declared that no competing interests exist.
- © 2019 Sociedade Brasileira de Zoologia. Published by Pensoft Publishers at <https://zoologia.pensoft.net>